

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

GENFARM
OXYFLUORFEN 240EC
HERBICIDE

ACTIVE CONSTITUENT: 240 g/L OXYFLUORFEN
SOLVENTS: 100 g/L N-METHYL PYRROLIDONE
605 g/L LIQUID HYDROCARBONS

GROUP **G** HERBICIDE

For selective weed control of broadleaf weeds and some grasses as specified in the Directions for Use table.

IMPORTANT: Read this booklet before use

Nutrien Ag Solutions Limited
Level 5, Building A, 26 Talavera Road, Macquarie Park, NSW, 2113
Tel: (02) 9889 5400 • Product Support Tel: 1800 44 88 92

DIRECTIONS FOR USE

CROP USE OR SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE
Treefruit, Nuts and Vines not less than 3 years old DORMANT APPLICATION Peaches Plums Apricots Almonds Grape Vines Olive Trees Apples Pears Macadamias Pecans Walnuts	Refer to Weeds Controlled list	Weed free soil	3 or 4 L/ha
		Emerged weeds present	
Peaches Plums Apricots Almonds Grapevines Apples Pears Macadamias Pecans Walnuts	Refer to Genfarm Panzer 360 label	Refer to Genfarm Panzer 360 label	75 mL/ha plus Genfarm Panzer 360 at recommended label rates.
Duboisia	Refer to Weeds Controlled list	Weed free soil	4 or 8 L/ha
		Weeds at the 4 to 6 true-leaf stage	

CRITICAL COMMENTS
DO NOT apply GENFARM OXYFLUORFEN 240EC Herbicide once bud swell has occurred. Apply to freshly cultivated weed free soil. Use the higher rate when longer residual control is required (up to 4 months). Where grass weeds are expected to be a major problem, or when control of a wider weed spectrum is required, mix the lower rate with 4.5L SURFLAN ² 500 per treated hectare. Read and follow all label directions.
DO NOT apply GENFARM OXYFLUORFEN 240EC Herbicide once bud swell has occurred. Use the higher rate when longer residual control is required (up to 4 months). When light infestations of weeds are present, apply as a tank mix with GENFARM PANZER 360 to produce both knockdown and residual control. Read and follow all label directions. Where weed growth is large and dense, weeds must be eliminated prior to application of GENFARM OXYFLUORFEN 240EC Herbicide, using GENFARM PANZER 360 or mechanical means.
Addition of GENFARM OXYFLUORFEN 240EC Herbicide to GENFARM PANZER 360 will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with GENFARM PANZER 360 applied alone), and give control of annual Nettles, <i>Urtica</i> spp. and Marshmallow, <i>Malva parviflora</i> . For rates of GENFARM PANZER 360, refer to the appropriate label. Read and follow all label directions. DO NOT apply the tank mix of Genfarm Panzer 360 and GENFARM OXYFLUORFEN 240EC near trees or vines less than 3 years old unless they are effectively shielded from spray and spray drift.
Apply after transplanting of young seedlings or after harvest of mature plants. Use the higher dosage where longer residual control (up to 6 months) is required. "over-the-top" application will be tolerated.
Best results are achieved when applied to moist soil free of weeds. However, some recently germinated small seedling grasses and broadleaf weeds (4 to 6 true leaves) will be controlled with these rates. Established larger weeds must be eliminated by mechanical means prior to application of GENFARM OXYFLUORFEN 240EC Herbicide. Add a non-ionic surfactant (at its recommended rate) to enhance activity.

CROP USE OR SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE
Tropical and Subtropical Fruit (inedible peel) Mango, Cherimoya Durian, Feijoa, Guava, Jackfruit, Longan, Mangosteen, Avocado Passionfruit Kiwi Fruit Custard Apple Lychee Pawpaw Longan Rambutan Star apple	Refer to Weeds Controlled list	Weed free soil Emergед weeds present	4 L/ha
Brassica Crops Cabbages Cauliflower Broccoli	Refer to Weeds Controlled list	Weed free soil	1.5 to 2 L/ha
Onions (seeded)	Blackberry nightshade, Dock, Crow'sfoot, Fumitory (Pinkweed), Hogweed (Wireweed), Plantain, Sorrel, Volunteer Potato, Potato Weed, Milk Thistle, Deadnettle, Common Cotula, Groundsel	Hook leaf stage 1½ to 2½ true leaf stage	50 – 100 mL/ha 350 – 500 mL/ha

CRITICAL COMMENTS
Best results are obtained when applied to moist soil free of weeds and followed by rainfall or irrigation.
If weeds are present, GENFARM OXYFLUORFEN 240EC Herbicide should be applied as a tank mix with recommended rates of GENFARM PANZER 360 , Genfarm Di-Par 250 ^o or Genfarm Paraquat 250 ^o . Read and follow all label directions.
Apply GENFARM OXYFLUORFEN 240EC Herbicide to prepared ground 4 to 7 days prior to transplanting. If soil is dry, irrigation or rainfall is required prior to transplanting for activation of GENFARM OXYFLUORFEN 240EC Herbicide. Utilise transplanting techniques which cause minimal soil disturbance. Excessive soil disturbance will lessen herbicidal activity. Use the higher rate in situations where weed pressure is known to be heavy.
Do not exceed 100 mL/ha at the hook leaf stage as excessive crop damage may occur.
Best results are obtained when the hook leaf stage treatment is followed by a further application of GENFARM OXYFLUORFEN 240EC Herbicide when onions are at the 1½ to 2½ true leaf stage. Apply GENFARM OXYFLUORFEN 240EC Herbicide in at least 100 L/ha water at pressures below 150 kPa. DO NOT exceed 150 kPa pressure because excessive crop damage could result. Multiple treatments may be applied provided the total dose does not exceed 2 L/ha in one season. For control of other weeds, use in a program with other registered herbicides. NOTE: Can cause flecking, twisting or stunting of onion plants although such symptoms are normally short-lived. DO NOT apply in conditions of very high humidity. DO NOT apply to weeds or crops which are stressed by prevailing weather conditions, disease or mechanical damage. DO NOT use wetting agents or other adjuvants.

CROP USE OR SITUATION	WEEDS CONTROLLED	TIME OF APPLICATION	RATE
Tobacco	Refer to Weeds Controlled list	Weed free soil	4 L/ha
Coffee	Refer to Weeds Controlled list	Weed free soil	2 to 4 L/ha
		Weeds at the 2 to 4 true leaf stage	
Forestry Trees	Broadleaf weeds and grasses	Weeds at the 4 to 6 true leaf stage	4 L/ha or 4 mL/10 m ²
Pyrethrum – as bare rooted transplants or seedlings	Refer to Weeds Controlled list (except chickweed)	Pre-plant incorporated into weed free soil worked to a fine tilth	4 or 6 L/ha
PRIOR TO SOWING Winter Cereals	Refer to GENFARM PANZER 450 or GENFARM PANZER 360 label	Refer to GENFARM PANZER 450 or GENFARM PANZER 360 label	75 mL/ha plus GENFARM PANZER 450 or GENFARM PANZER 360 at recommended label rates
PRIOR TO SOWING Cotton			

When using GENFARM Oxyfluorfen 240EC Herbicide in combination with other herbicides, refer to the appropriate label and read and follow all label directions.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THE LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS:

GRAZING DO NOT GRAZE TREATED WEEDS.
HARVEST NOT REQUIRED WHEN USED AS DIRECTED.

CRITICAL COMMENTS
Use to control weeds along spray lines only. Do NOT apply to tobacco crop. Apply to soil after solid-set irrigation system has been laid out in the field. GENFARM OXYFLUORFEN 240EC Herbicide should be applied to a moist soil. Where very small weeds (2-3 leaf) emerge prior to spraying, the addition of a wetting agent to the spray mixture is necessary for effective control. Should the weeds be more advanced, the addition of 2 L Diquat 200 is required. Avoid spray drift.
Apply after transplanting coffee plants or under established bushes. Use the higher rate where longer residual activity (up to 4 months) is required. Best results are achieved when applied to moist soil, free of weeds with follow-up irrigation.
When seedlings are present, apply as a tank mix with Paraquat 250 to produce both knockdown and residual control. AGRAL 600 should be used in the spray mixture at 250 mL/100 L. Apply as a directed spray to avoid contact with coffee plants. Mature established weeds must be eliminated by mechanical or chemical means prior to application.
For the establishment of trees for approved farm practices such as wind breaks, erosion control, woodlots and forestry plantings. When applying as a post-plant spray, ensure spray is directed to the base of seedlings, or that seedlings are protected. Do not apply under hot or windy conditions.
Apply prior to final soil preparation. The preferred implements for final soil preparation would be either a multiple tyne cultivator or rotary harrows. Use the 6 L/ha rate for heavy black clay soils only (as found in the Derwent and Coal River Valleys, Tasmania). GENFARM OXYFLUORFEN 240EC Herbicide will not provide consistent control of chickweed.
Addition of GENFARM OXYFLUORFEN 240EC Herbicide to GENFARM PANZER 450 or GENFARM PANZER 360 will improve knockdown and increase the speed at which treated weeds develop visible symptoms of phytotoxicity (compared to results achieved with GENFARM PANZER 450 or GENFARM PANZER 360 applied alone), and give control of annual Nettles, <i>Urtica</i> spp. and Marshmallow, <i>Malva parviflora</i> .

Weeds Controlled by GENFARM OXYFLUORFEN 240EC Herbicide include the following:

BEFORE GERMINATION

Amsinckia *Amsinckia* spp.
Barley grass *Hordeum leporinum*
Barnyard grass *Echinochloa* spp.
Blackberry nightshade *Solanum nigrum*
Bladder ketmia *Hibiscus trionum*
Burrgrass *Cenchrus australis*
Caltrop *Tribulus terrestris*
Capeweed *Arctotheca calendula*
Chickweed *Stellaria media*
Crowsfoot grass *Eleusine indica*
Deadnettle *Lamium amplexicaule*
Fat hen *Chenopodium album*
Giant pigweed *Trianthema portulacastrum*
Liverseed grass *Urochloa panicoides*
Lovegrass *Eragrostis* spp.
Pigeon grass *Setaria* spp.
Pigweed *Portulaca oleracea*
Prickly lettuce *Lactuca* spp.
Red Natal grass *Rhynchelytrum repens*
Redshank *Amaranthus cruentus*
Ryegrass *Lolium* spp.
Sesbania pea *Sesbania cannabina*
Shepherd's purse *Capsella bursa-pastoris*
Small Flowered Mallow *Malva parviflora*
Soursob *Oxalis pes-caprae*
Sowthistle *Sonchus oleraceus*
Starburr *Acanthospermum hispidum*
Stinkgrass *Eragrostis cilianensis*
Summer grass *Digitaria* spp.
Thornapple *Datura stramonium*
White eye *Richardia brasiliensis*
Wild mustard *Sisymbrium* spp.
Wild radish *Raphanus raphanistrum*
Wireweed *Polygonum aviculare*

SEEDLINGS (less than 4 true leaf)

Amsinckia *Amsinckia* spp.
Bellvine *Ipomoea* spp.
Capeweed *Arctotheca calendula*
Common cotula *Cotula australis*
Growsfoot grass *Eleusine indica*
Deadnettle *Lamium amplexicaule*
Groundsel *Senecio vulgaris*
Liverseed grass *Urochloa panicoides*
Small Flowered Mallow *Malva parviflora*
Pigweed *Portulaca oleracea*
Potato weed *Galinoga parviflora*
Redshank *Amaranthus cruentus*
Shepherd's purse *Capsella bursa-pastoris*
Sowthistle *Sonchus oleraceus*
Stinging nettle *Urtica urens*
Stinkgrass *Eragrostis cilianensis*
Wild radish *Raphanus raphanistrum*

GENERAL INSTRUCTIONS

RESISTANT WEEDS WARNING

GROUP G HERBICIDE

GENFARM OXYFLUORFEN 240EC Herbicide is a member of the Diphenyl ether group of herbicides. The mode of action of GENFARM OXYFLUORFEN 240EC Herbicide is to inhibit protoporphyrinogen oxidase. For weed resistance management, GENFARM OXYFLUORFEN 240EC Herbicide is a Group G herbicide.

Some naturally-occurring weed biotypes resistant to GENFARM OXYFLUORFEN 240EC Herbicide and other inhibitors of protoporphyrinogen oxidase herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These weeds will not be controlled by GENFARM OXYFLUORFEN 240EC Herbicide or other inhibitors or protoporphyrinogen oxidase herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Nutrien Ag Solutions Limited accepts no liability for any losses that may result from the failure of GENFARM OXYFLUORFEN 240EC Herbicide to control resistant weeds.

TIMING

For optimum residual weed control, GENFARM OXYFLUORFEN 240EC Herbicide should be applied to the soil surface prior to weed emergence after all other agricultural operations, such as mechanical cultivation and re-shaping of irrigation furrows, have been completed. The area should be left undisturbed during the time period for which weed control is desired. When applied to seedling weeds, they should be young and actively growing. Weed control for up to 6 months is expected but spot treatment, with knockdown herbicides, for escape weeds and perennial grasses may be necessary.

MIXING

When using GENFARM OXYFLUORFEN 240EC Herbicide alone, fill the spray tank at least one-third full with clean water, add the recommended amount of GENFARM OXYFLUORFEN 240EC Herbicide while the pump and agitator are running, then complete filling the spray tank.

A non-ionic surfactant, if required by label directions, should be added near the end of the filling process to minimize foaming.

When tank mixing with Surflan 500, add to one-third filled tank and then add the GENFARM OXYFLUORFEN 240EC Herbicide during the filling operation. Maintain agitation during mixing and until spraying is completed.

When tank mixing with GENFARM PANZER 450 or GENFARM PANZER 360, add these after GENFARM OXYFLUORFEN 240EC Herbicide during the filling operation.

When using GENFARM OXYFLUORFEN 240EC Herbicide in combination with GENFARM PANZER 450, GENFARM PANZER 360 or Surflan 500, refer to the appropriate label and read and follow all label directions.

APPLICATION

Spray equipment should be calibrated carefully before use.

GENFARM OXYFLUORFEN 240EC Herbicide should be applied uniformly as a directed treatment to the base of tree and vine crops using flat fan or off-centre nozzles. Complete coverage of seedling weeds is required for maximum knockdown effect.

A water volume of 250 to 500 litres per hectare is recommended for treatments on bare soil.

A spray volume of 100 to 1350 litres per hectare is recommended where seedling weeds (4 to 6 leaf stage) are present. Ensure both weed foliage and the soil surface are sprayed. Use the higher volumes where weed density is high.

Tank mixtures of 75 mL/ha of GENFARM OXYFLUORFEN 240EC Herbicide with GENFARM PANZER 450 or GENFARM PANZER 360 herbicides, should be applied in 30 to 200 litres spray volume per hectare. For maximum residual control, GENFARM OXYFLUORFEN 240EC Herbicide should NOT be incorporated or disturbed after application.

CROP SAFETY

GENFARM OXYFLUORFEN 240EC Herbicide may be applied as directed around dormant peach, plum, apricot, almond, apple and pear trees and grapevines of all ages when applied at rates of less than 1.0 L/ha.

When applied at 3.0 L/ha and above, the trees and grape vines should be at least 3 years of age.

DO NOT apply GENFARM OXYFLUORFEN 240EC Herbicide once bud swell has occurred when using rates greater than 1.0 L/ha.

Duboisia seedlings and mature plants will tolerate "over-the-top" applications of GENFARM OXYFLUORFEN 240EC Herbicide. When using the 75 mL/ha rate, GENFARM OXYFLUORFEN 240EC Herbicide may be applied as directed prior to sowing winter cereals.

SAFE SOWING INTERVALS

GENFARM OXYFLUORFEN 240EC Herbicide at up to 75mL/ha may be safely applied 1 day prior to planting broadacre crops such as cereals (wheat, barley, oats, triticale), canola, pulses (lupins, faba beans, field peas) and undersown pastures (lucerne, clover, medics, ryegrass, phalaris, cocksfoot) and 7 days minimum prior to plating cotton or soybean, provided minimum tillage planting equipment is used with minimal soil disturbance. Inversion, mixing of soil surface with that in the planting zone or covering seed with treated soil may result in injury to the emerging crop seedlings. **Avoid covering the seed with soil treated with Genfarm OXYFLUORFEN during the planting operation to minimize crop injury.**

Genfarm OXYFLUORFEN has residual soil activity, especially when applied at rates greater than 75mL/ha and on small seeded horticultural crops. Plantback intervals in the following table for horticultural crops must be observed if more than 75mL/ha Genfarm OXYFLUORFEN has been applied.

Safe sowing Intervals for Horticultural Crops (Days)

Oxyfluorfen Herbicide rate	Up to 75mL/ha	1L/ha
Beans	7	60
Brassicac	14	90
Capsicums	14	90
Carrots	14	90
Cucurbits	14	60
Lettuce	14	90
Onions	21	180
Potatoes	7	60
Tomatoes	14	60

COMPATIBILITY

GENFARM OXYFLUORFEN 240EC Herbicide is compatible with Genfarm Panzer 360, Genfarm Panzer 450, Genfarm Panzer 510, Oryzalin 500, Genfarm Paraquat 250 Herbicide, Genfarm Di-Par 250, Diquat 200, Touchdown and Basta.

RE-ENTRY PERIOD

Do not allow entry into treated areas until the spray has dried. If prior entry is necessary, wear chemical resistant footwear.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

DO NOT apply under weather conditions, or from spraying equipment, that may cause spray to drift onto nearby susceptible plants/crops, cropping lands or pastures.

PROTECTION OF LIVESTOCK

Use with care when applying in areas frequented by stock.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

This product is highly toxic to wildlife and fish.

DO NOT contaminate streams, rivers or waterways with the chemical or used containers. Use care when applying in areas frequented by wildlife or adjacent to any body of water.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank.

Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

Empty containers and product should not be burnt.

SAFETY DIRECTIONS

Will damage the skin. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container and preparing spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist and a washable hat, elbow-length butyl rubber gloves and goggles.

If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre (Phone Australia: 13 11 26).

SAFETY DATA SHEET

Additional information is listed in the Safety Data Sheet, which is available from the supplier.

CONDITIONS OF SALE

The use of this product is beyond the control of Nutrien Ag Solutions Limited. No warranty is expressed or implied regarding the suitability or efficiency for any purpose for which it is used by the buyer. Nutrien Ag Solutions Limited accepts no responsibility for any consequences resulting from the use of this product. Nutrien Ag Solutions Limited will not be held liable for any loss, injury or damage arising from the sale, supply or use of this product, whether through negligence or otherwise. No responsibility will be accepted for any consequences whatsoever resulting from the use of this product.



Nutrien Ag Solutions Limited

Level 5, Building A, 26 Talavera Road,
Macquarie Park, NSW, 2113 • Tel: (02) 9889 5400

Product Support Tel: 1800 44 88 92