

Bungalow™ 0-29-30

For Use on Turf and Ornamentals

GUARANTEED ANALYSIS

Available Phosphate (P_2O_5)	29.0%
Soluble Potash (K_2O)	30.0%
Derived from: Potassium Phosphate	◆◆◆ Contains 4.3 lbs of Phosphorus Acid per Gallon

BEFORE USING THIS PRODUCT, READ ALL PRECAUTIONARY STATEMENTS, DIRECTIONS FOR USE AND WARRANTY AND LIMITATION OF DAMAGES.

KEEP OUT OF REACH OF CHILDREN

WARNING

Harmful if swallowed. Avoid inhalation of spray mist. May cause irritation of nose, throat, or skin. Avoid contact with eyes, skin, and clothing. In case of contact with skin or eyes, flush affected area with water; for eyes seek medical attention. Wear suitable gloves and face shield when handling this product. Do not apply this product in such a manner as to directly expose workers or other persons. If product is being mixed with pesticides and/or spray adjuvants, follow all precautionary statements on the accompanying product's labeling.

STORAGE AND DISPOSAL

Store product in a cool, dry, area, out of reach of children and livestock. Keep away from food, drink and fodder. Avoid storage in extreme temperatures. Store Bungalow in its original container. Keep container closed. Do not contaminate water sources by cleaning of equipment or disposal of spray waste and rinse water. Triple rinse empty containers, puncture and discard in accordance with applicable local, state, and federal regulations. Do not allow to freeze (store above 32F).

WARRANTY AND LIMITATION OF DAMAGES

CONDITIONS OF SALE: To the extent consistent with applicable law, Sipcarn Agro USA, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with the directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Sipcarn Agro USA, Inc. Sipcarn Agro USA, Inc. disclaims all other warranties, express or implied. To the extent consistent with applicable law, Sipcarn Agro USA, Inc. shall not be liable for consequential, special, or indirect damages resulting from the use or handling of this product, and Sipcarn Agro USA, Inc.'s sole liability and Buyer's and User's exclusive remedy shall be limited to the refund of the purchase price. To the extent consistent with applicable law, Buyer and User acknowledge and assume all risks and liability resulting from the handling, storage, and use of this product. Sipcarn Agro USA, Inc. does not authorize any agent or representative to make any other warranty, guarantee or representation concerning this product.

Bungalow™ is a trademark of S.I.P.C.A.M. SOCIETA' ITALIANA PRODOTTICIMICI E PER L'AGRICOLTURA MILANO S.p.A.

Manufactured For:
SIPCARN AGRO USA, INC.
2525 Meridian Parkway, Suite 350
Durham, NC 27713



**READ THE LABEL CAREFULLY
BEFORE OPENING THE CONTAINER
SHAKE WELL BEFORE USING**

Density: 12.62 lb / gal. @ 68° F (5.72 kg)

F002220

NET CONTENTS: 2.5 Gallons (9.46L) NET WEIGHT: 31.55 lb (14.31 kg)

Bungalow™ 0-29-30

For Use on Turf and Ornamentals

GENERAL INFORMATION

Bungalow is a highly concentrated soluble nutrient solution intended to be used as a supplemental foliar treatment. It is readily absorbed through leaf or root tissue and quickly corrects deficiencies and enhances plant vigor, crop development, quality, and yield.

Information regarding the contents and levels of metals in this product is available on the internet at:

<http://www.aapfco.org/metals.htm>

DIRECTIONS FOR USE

TURFGRASS: Warm and Cool Season - Apply 1.5 to 3 ounces per 1000 sqft and repeat every 7 to 14 days. During Times of Stress – Apply 4 - 6 ounces per 1000 sqft and repeat every 7-14 days. Fairways, Sod & Finely Maintained Turf – Apply 2 to 8 quarts per acre in a minimum 100 gallons of water to ensure proper coverage. Overseeding, Sprigging, or Sod – Apply 2 to 8 quarts per acre at the time of germination, sprigging, or laying of sod and repeat in 14 day intervals thereafter in a minimum of 100 gallons of water until turf is well established.

ORNAMENTAL PLANTS: Transplanting/Propagule - Apply as a 0.5% v/v solution as a foliar spray or immerse the propagule or transplant tray temporarily in a 0.5% v/v solution before transplanting. Plant Vigor - Apply a 0.5% v/v solution to foliage every 10 to 15 days as needed starting at bud break through senescence.

For Professional Use Only

Do not mix with fixed copper or copper-based fungicides if the pH of the spray solution is below pH 6.

In Florida follow BMP's found at

<http://www.dep.state.fl.us/water/nonpoint/pubs.htm>

MIXING INSTRUCTIONS

Prior to any fertilizer or pesticide application, all spray mixing and application equipment must be cleaned. Fill the spray or mix tank one half full of water and begin agitation. Add pesticides and/or fertilizers as directed by labeling or in the following sequence:

1. Dry flowables or water dispersible granules
2. Wettable powders
3. Flowables
4. Emulsifiable concentrates
5. Water-based solutions
6. Compatibility agents
7. Micronutrients and fertilizers
8. Spray adjuvants.

FIRST AID

If Swallowed: If conscious, give water to drink. Immediately call a doctor.

Eye Contact: Flush eyes with clean water for 15 minutes, holding eyelids open. Obtain medical attention immediately.

Skin Contact: Remove contaminated clothing and wash skin thoroughly with soap and water. Obtain medical attention immediately.

If Inhaled: Move to fresh air and call a doctor if irritation develops or persists. If there is contact made with the spray solution containing pesticides and/or spray adjuvants, follow the First Aid Statement on the product's labeling.

Wear suitable gloves and face shield when handling this product. Do not apply this product in such a manner as to directly expose workers or other persons. If product is being mixed with pesticides and/or spray adjuvants, follow all precautionary statements on the accompanying product's labeling.