

LIQUID COPPER 5%

GENERAL INFORMATION

CNI LIQUID COPPER 5% is formulated for tree nut crops as well as field, vegetable, row, fruit, tree, vine, ornamental and turf

CNI LIQUID COPPER 5% may be applied through drip systems. Wt per gal: 9.50 lb. at 68°F Grams per liter: 1139 g @20°C aircraft, or by ground sprayers (high-volume or low-volume).

CNI LIQUID COPPER 5% may be applied alone or with Always add this product to spray tank before adding a fertilizers or pesticides. It can be applied either to the plant foliage or to the soil.

CNI LIQUID COPPER 5% is compatible with most pesticides Foliar Application: and fertilizers. However, a compatibility test is recommended if the desired combination has not been used in the past.

SHAKE WELL BEFORE USING

CONDITIONS OF SALE

Seller warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Weather, crop conditions, timing and methods Soil Application: of applications, mixtures with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict Aerial Application: accordance with directions given herewith.

Information regarding contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.html

GUARANTEED ANALYSIS

Sulfur(S)	3.7%
COPPER (Cu)	5.0%
Derived from Copper lignosulfonate	

DIRECTIONS FOR USE

pesticide.

RATES FOR ALL CROPS

Apply 1 to 4 quarts/ acre as needed on a 10- 14 day schedule.

When rates of over 2 quarts are required, split applications are recommended.

Avoid foliar applications of CNI LIQUID COPPER 5% when plants are under stress due to lack of moisture. AVOID FOLIAR APPLICATION ON DECIDUOUS TREE CROPS **DURING BLOOM PERIOD.**

For fertilizer maintenance apply 1-2 quarts per acre For mild deficiency apply 2-4 quarts per acre For high deficiency apply 1-2 gal per acre.

Apply 1 quart of this product per 5 to 10 gallons of water per acre.

Drip Irrigation Application:

Apply the suggested rate of this product in multiple applications 10 -14 days apart throughout the growing season of the crop.

KEEP OUT OF REACH OF CHILDREN



WARNING

WARNING Hazardous to humans and animals. Causes eye and skin irritations. When mixing or applying agricultural sprays, always wear spray goggles and impervious gloves plus protective clothing as recommended on pesticide

FIRST AID If this product gets into the eyes, immediately wash the eves with large amounts of water for at least 15 minutes, occasionally lifting the upper and lower lids. Obtain immediate medical treatment. If this product gets on the skin, immediately wash the contaminated area. Seek medical attention if skin irritation persists. If this product has been swallowed and the person is conscious. immediately give large volumes of water. Do not induce vomiting. Seek immediate medical attention.

STORAGE AND DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Follow proper procedures for the disposal of spray solutions. Rinse and dispose of empty containers according to approved local practices.

CONDITIONS OF SALE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Weather, crop conditions, timing and methods of applications, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions 10-17

