

# FBN Clethodim 240®

Use with FBN Clethodim/Companero Adjuvant

## EMULSIFIABLE CONCENTRATE POST- EMERGENCE HERBICIDE

GROUP	<b>1</b>	HERBICIDE
-------	----------	-----------

### CONTAINS CLETHODIM

POST-EMERGENCE HERBICIDE FOR CONTROL OF GRASSES IN CANOLA, FLAX (including low linolenic acid varieties), FIELD PEAS, LENTILS, DESI and KABULI CHICKPEAS, DRY BULB ONIONS, POTATOES, YELLOW MUSTARD, ORIENTAL (BROWN) MUSTARD (condiment and oilseed types), SOYBEANS, SEEDLING ALFALFA, SUNFLOWERS, DRY COMMON BEANS (*Phaseolus vulgaris* varieties only), FENUGREEK, CORIANDER, SPINACH, PRAIRIE CARNATION, Highbush Blueberry, SAFFLOWER, CRANBERRY, DILL, CARAWAY, BASIL, RED GARDEN (TABLE) BEET, PARSNIP, CARROT AND RADISH.

### COMMERCIAL (AGRICULTURAL)

**ACTIVE INGREDIENT: CLETHODIM .....240 g/L**

REGISTRATION NO. 33646 PEST CONTROL PRODUCTS ACT

**WARNING: EYE AND SKIN IRRITANT**

**KEEP OUT OF REACH OF CHILDREN**

**READ THE LABEL AND BOOKLET BEFORE USING**

**NET CONTENTS: 1 Litre - Bulk**

**Product Information:** 1-844-200-FARM (3276)

Farmer's Business Network Canada, Inc.  
PO Box 5607  
High River, Alberta  
T1V 1M7

In case of spills, poisoning or fire, telephone emergency response number CANUTEC 613-996-6666 or \*666 on a cellular phone. (24 hours a day).

## **PRECAUTIONS:**

KEEP OUT OF REACH OF CHILDREN. CAUSES EYE AND SKIN IRRITATION.

Avoid

contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Do not inhale fumes. Avoid breathing vapours or spray mist. After use, wash hands and other exposed skin. When using, do not eat, drink or smoke. Remove and launder contaminated clothing separately from household laundry before reuse. Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs.

Observe appropriate provincial buffer zones around bodies of water and wetland areas. Wear a long-sleeved shirt, long pants, rubber apron, chemical-resistant gloves, socks and rubber boots during mixing, loading, application, clean-up and repair. Gloves and rubber apron are not required during application. In addition, wear goggles/face shield during mixing/loading and when handling the concentrate.

Application is limited to agricultural crops only when there is low risk to drift to areas of human habitation or activity such as houses, cottages, schools and recreation areas, taking into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: [www.croplife.ca](http://www.croplife.ca)

## **ENVIRONMENTAL PRECAUTIONS:**

TOXIC to aquatic organisms and non-target terrestrial plants. Observe buffer zones specific under DIRECTIONS FOR USE.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains aromatic petroleum distillates that are toxic to aquatic organisms.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (for example, sandy soil) and/or the depth to the water table is shallow.

## **FIRST AID:**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give **any** liquid to the person. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Danger from lung aspiration of petroleum-based solvents must be weighed against toxicity when considering emptying stomach.

## **STORAGE AND DISPOSAL**

**STORAGE:** Store this product away from food or feed.

### **DISPOSAL:**

**RECYCLABLE CONTAINER DISPOSAL:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**RETURNABLE CONTAINER DISPOSAL:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**REFILLABLE CONTAINER DISPOSAL:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:** For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

SPECIMEN

**FBN Clethodim 240<sup>®</sup>**

Use with FBN Clethodim/Companero Adjuvant

# EMULSIFIABLE CONCENTRATE POST- EMERGENCE HERBICIDE

GROUP	1	HERBICIDE
-------	---	-----------

## CONTAINS CLETHODIM

POST-EMERGENCE HERBICIDE FOR CONTROL OF GRASSES IN CANOLA, FLAX (including low linolenic acid varieties), FIELD PEAS, LENTILS, DESI and KABULI CHICKPEAS, DRY BULB ONIONS, POTATOES, YELLOW MUSTARD, ORIENTAL (BROWN) MUSTARD (condiment and oilseed types), SOYBEANS, SEEDLING ALFALFA, SUNFLOWERS, DRY COMMON BEANS (*Phaseolus vulgaris* varieties only), FENUGREEK, CORIANDER, SPINACH, PRAIRIE CARNATION, HIGHBUSH BLUEBERRY, SAFFLOWER, CRANBERRY, DILL, CARAWAY, BASIL, RED GARDEN (TABLE) BEET, PARSNIP, CARROT AND RADISH.

## COMMERCIAL (AGRICULTURAL)

**ACTIVE INGREDIENT: CLETHODIM .....240g/L**

REGISTRATION NO. 33646 PEST CONTROL PRODUCTS ACT

**WARNING: EYE AND SKIN IRRITANT**

**KEEP OUT OF REACH OF CHILDREN**

**READ THE LABEL AND BOOKLET BEFORE USING**

**NET CONTENTS: 1 LITRE - BULK**

**Product Information:** 1-844-200-FARM (3276)

Farmer's Business Network Canada, Inc.  
PO Box 5607  
High River, Alberta  
T1V 1M7

In case of spills, poisoning or fire, telephone emergency response number 613-996-6666 or \*666 on a cellular phone (24 hours a day).

## GENERAL INFORMATION:

**FBN Clethodim 240** is a selective post-emergence herbicide for control of a broad range of grasses in canola, flax (including low linolenic acid varieties), field peas,

lentils, Desi and Kabuli chickpeas, dry bulb onions, potatoes, yellow mustard, oriental (brown) mustard (condiment and oilseed types), soybeans, seedling alfalfa, sunflowers, dry common beans (*Phaseolus vulgaris* varieties only), fenugreek, coriander, spinach, prairie carnation, highbush blueberry, safflower, cranberry, dill, caraway, basil, red garden(table) beet, parsnip, carrot and radish.

**FBN Clethodim 240 is to be used only with the FBN Clethodim / Companero Adjuvant.**

**FBN Clethodim 240** is a systemic herbicide which is translocated from the treated foliage to the growing points of the leaves, shoots and roots. Uptake into the plant is primarily through its leaves. Thorough coverage of the foliage is important for consistent grass control. Susceptible grasses that are actively growing will undergo a burn-back and colour change after treatment. Leaf foliage will first change from green to yellowish, then purplish and finally brown colour. The time required for complete control is normally 7 to 21 days following treatment, depending on growing conditions and crop competition.

### **PRECAUTIONS:**

**KEEP OUT OF REACH OF CHILDREN. CAUSES EYE AND SKIN IRRITATION.** Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Do not inhale fumes. Avoid breathing vapours or spray mist. After use, wash hands and other exposed skin. When using, do not eat, drink or smoke. Remove and launder contaminated clothing separately from household laundry before reuse. Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Observe appropriate provincial buffer zones around bodies of water and wetland areas. Wear a long-sleeved shirt, long pants, rubber apron, chemical-resistant gloves, socks and rubber boots during mixing, loading, application, clean-up and repair. Gloves and rubber apron are not required during application. In addition, wear goggles/face shield during mixing/loading and when handling the concentrate.

Application is limited to agricultural crops only when there is low risk to drift to areas of human habitation or activity such as houses, cottages, schools and recreation areas, taking into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

**DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.**

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: [www.croplife.ca](http://www.croplife.ca)

### **ENVIRONMENTAL PRECAUTIONS:**

**TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specific under DIRECTIONS FOR USE.

Toxic to certain beneficial insects. Minimize spray drift to reduce harmful effects on beneficial insects in habitats next to the application site such as hedgerows and woodland.

To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product contains aromatic petroleum distillates that are toxic to aquatic organisms.

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (for example, sandy soil) and/or the depth to the water table is shallow.

#### **FIRST AID:**

**If swallowed:** Call a poison control centre or doctor immediately for treatment advice. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do **not** give **any** liquid to the person. Do not give anything by mouth to an unconscious person. **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

**If in eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**If inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

**TOXICOLOGICAL INFORMATION:** Treat symptomatically. This product contains petroleum distillates. Vomiting may cause aspiration pneumonia. Danger from lung aspiration of petroleum-based solvents must be weighed against toxicity when considering emptying stomach.

#### **STORAGE AND DISPOSAL**

**STORAGE:** Store this product away from food or feed.

**DISPOSAL:**

**RECYCLABLE CONTAINER DISPOSAL:** Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site.

Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

- Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
- Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

**RETURNABLE CONTAINER DISPOSAL:** Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

**REFILLABLE CONTAINER DISPOSAL:** For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

**DISPOSAL OF UNUSED, UNWANTED PRODUCT:** For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

#### **DIRECTIONS FOR USE:**

As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

#### **TIME OF APPLICATION:**

Apply **FBN Clethodim 240** when the annual grasses and volunteer cereals are in the 2 to 6 leaf stage as specified under Application Rates of FBN Clethodim 240. Most effective control is achieved when application is made prior to tillering when annual grasses are small and actively growing.

For suppression or control of quackgrass, apply **FBN Clethodim 240** when the quackgrass is in the 2 to 6 leaf stage and 6 to 15 cm in height. Most effective results are achieved when application is made at the 3 to 5 leaf stage, when the canopy is uniform and actively growing.

**FBN Clethodim 240** will be less effective when plants are stressed by lack of moisture, excessive moisture, low-temperature and/or very low relative humidity.



Regrowth by tillering may occur if application is made under any of the above stress conditions. In wide row crops, where the canopy may be slow to close, cultivation may be necessary to control grasses that emerge after treatment.

Canola, flax (including low linolenic acid varieties), field peas, lentils, potatoes, yellow mustard, oriental (brown) mustard (condiment and oilseed types), soybeans, seedling alfalfa, sunflowers, dry common beans (*Phaseolus vulgaris* varieties only), highbush blueberry, coriander, caraway, basil, red garden (table) beet, parsnip, carrot and radish are tolerant to **FBN Clethodim 240** at all stages of growth. However, the pre-harvest interval outlined in the Crop Recommendations table must be followed to avoid excessive crop residues.

For Desi and Kabuli chickpeas, apply **FBN Clethodim 240** at a maximum of one application per season before the crop reaches the 9 node stage (18 cm height maximum) and when the annual grasses and volunteer cereals are in the 2 to 6 leaf stage.

For fenugreek and dill, apply FBN Clethodim 240 post-emergence, when fenugreek and dill are in the 3 - 5 leaf stage. Apply a maximum of one application may be made per season.

For coriander or prairie carnation, apply FBN Clethodim 240 post-emergence, when coriander or prairie carnation is in the 2 - 5 leaf stage. A maximum of one application may be made per year.

For safflower, apply FBN Clethodim 240 post-emergence, when safflower is in the 6-8 leaf stage at a maximum of one application per season.

Cranberry is tolerant to **FBN Clethodim 240** at all stages of growth, however do not apply FBN Clethodim 240 between the hook stage and full fruit set.

For dry bulb onions, apply FBN Clethodim 240 post-emergence, when crops are in the 1-4 leaf stage.

#### **APPLICATION RATES OF FBN Clethodim 240:**

Rates of use are given below to control various grass species. Refer to crop section for the maximum use rate on each crop.

<b>Grass Species</b>	<b>Leaf Stage</b>	<b>Application Rate of FBN Clethodim 240®</b>	<b>Application Rate of FBN Clethodim/ Companero Adjuvant.</b>
*Foxtail (green, yellow) Wild oats Volunteer cereals (wheat, barley, oats)	2-4	0.125 L/ha	0.5% v/v

Barnyard Grass Witchgrass Fall panicum Proso millet Volunteer Corn Volunteer canary grass	2-6	0.125 L/ha	0.5% v/v
Wild Oats Volunteer cereals (wheat, barley, oats) Foxtail (green, yellow) Persian darnel Crabgrass (smooth and large) Proso millet Witchgrass Fall panicum Barnyard grass Volunteer corn Volunteer canary grass Quackgrass suppression	2-6	0.19 L/ha	0.5% v/v
Quackgrass Control**	2-6	0.38 L/ha	1.0% v/v

\* **FBN Clethodim 240** applied at 0.125 L/ha for the control of weeds listed in this section of the table should only be applied under the following conditions:

- Good crop stand
- Early application (as in above table, prior to tillering)
- Do not tank-mix with other pesticides
- Do not apply to volunteer winter cereals
- Light to moderate weed infestation
- Adequate moisture and fertility
- Absence of stress, good growing conditions

If any one of the above conditions are not present at the time of application, use 0.19 L/ha rate of **FBN Clethodim 240**.

\*\*For adequate control of quackgrass use a minimum spray volume of 100 L/ha.

**CROP RECOMMENDATIONS:**

Crops	Maximum Application Rate of FBN Clethodim 240	Application Method <sup>7</sup>	Pre-Harvest Interval
Canola	0.38 L/ha	Ground and aerial	60 days

		application	
Flax (including low linolenic acid varieties)	0.38 L/ha	Ground and aerial application	60 days
Field Peas	0.38 L/ha	Ground and aerial application	75 days
Lentils	0.38 L/ha	Ground and aerial application	60 days
Desi and Kabuli Chickpeas	0.19 L/ha	Ground and aerial application	60 days
Potatoes	0.38 L/ha	Ground and aerial application	60 days
Yellow Mustard	0.38 L/ha	Ground and aerial application	60 days
Oriental (Brown) Mustard (condiment and oilseed types)	0.38 L/ha	Ground and aerial application	60 days
Soybeans	0.38 L/ha	Ground and aerial application	75 days
Seedling Alfalfa	0.38 L/ha	Ground application only	30 days
Sunflowers	0.38 L/ha	Ground and aerial application	72 days
Dry Common Beans <sup>1,2</sup> ( <i>Phaseolus vulgaris</i> varieties only)	0.19 L/ha	Ground and aerial application	60 days
Basil <sup>1,2,4,5</sup>	0.19 L/ha	Ground application only	30 days
Caraway <sup>1,5,6</sup>	0.38 L/ha	Ground application only	60 days
Coriander <sup>1,5,6</sup>	0.38 L/ha	Ground application only	60 days
Cranberry <sup>1,2,5</sup>	0.38 L/ha	Ground application only	30 days
Dill <sup>1,5</sup>	0.38 L/ha	Ground application only	40 days
Dry Bulb Onions <sup>1,5</sup>	0.38 L/ha	Ground application only	45 days
Fenugreek <sup>1,5,6</sup>	0.38 L/ha	Ground application only	30 days
Highbush Blueberry <sup>1,4,5</sup>	0.38 L/ha	Ground application only	14 days
Prairie	0.19 L/ha	Ground application only	N/A

Carnation <sup>1,5,6</sup>			
Red garden (table) beet, parsnip, carrot, radish <sup>3,5</sup>	0.38 L/ha	Ground application only	30 days
Safflower <sup>1,5</sup>	0.38L/ha	Ground application only	70 days
Spinach <sup>3,5</sup>	0.19L/ha	Ground application only	14 days

1. Apply a maximum of one application per year.
2. NOTE: Varieties of this crop may vary in their tolerance to herbicides, including to FBN Clethodim 240. Since not all crop varieties have been tested for tolerance to FBN Clethodim 240, first use of FBN Clethodim 240 should be limited to a small area of each variety to confirm tolerance prior to adoption as a general field practice. Additionally, consult your supplier for information on the tolerance of specific varieties to FBN Clethodim 240.
3. Apply a maximum of two applications per year. If repeat application is required, allow at least 14 days between first and second application.
4. Apply FBN Clethodim 240 as a broadcast spray directed to the ground.
5. Refer to section "MINOR USES" in this label for additional use instructions.
6. Do not harvest treated Prairie Carnations or greens of treated fenugreek as a vegetable or leaves of treated coriander (cilantro) and caraway for human consumption.
7. SPRAY VOLUME FOR AERIAL APPLICATION: Apply in 28 to 55 L/ha. Increase spray volumes up to 100 L/ha as grass or crop foliage becomes dense.

#### TANK-MIXES:

When applied as a tank-mix combination, read and observe all label directions, including rates, restrictions, and grazing limitations for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

#### FBN Clethodim 240 AND BUCTRIL® M:

**Flax (including low linolenic acid varieties):** FBN Clethodim 240 may be tank-mixed with BUCTRIL M for control of certain broadleaf and grassy weeds in flax (including low linolenic acid varieties) in one spray operation. Apply in 55 - 110 litres of water per hectare at a spray pressure of 275 kPa. DO NOT spray this tank-mix on plants under stress. DO NOT spray during periods of hot, humid weather. Follow all precautions, limitations and timing recommendations on the BUCTRIL M label. The most stringent label precautions for any individual pesticide product in a tank-mix should be used.

**Time of Application:** Flax (including low linolenic acid varieties) may be treated from 5 cm high to the early bud stage. For best results, apply when flax (including low linolenic acid varieties) is 5 cm to 10 cm high and weeds are in the seedling stage. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 - 0.38 L/ha tank-mixed with BUCTRIL M at 1.0 L/ha and the FBN Clethodim Adjuvant/**Companero Adjuvant** at 0.5% - 1.0% v/v.

**Weeds Controlled:** As listed for FBN Clethodim 240 used alone, plus certain seedling broadleaf weeds. Refer to the BUCTRIL M label for the broadleaf weeds controlled.

#### MIXING INSTRUCTIONS:

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- If tank-mixing FBN Clethodim 240 and BUCTRIL M for once-over grass and broadleaf weed control in flax (including low linolenic acid varieties), add the correct amount of BUCTRIL M. Continue to agitate.
- Add the correct amount of FBN Clethodim 240. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/ Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

*For disposal of cleaning solution, see section on DISPOSAL.*

#### FBN Clethodim 240 AND CURTAIL® M:

**Flax (including low linolenic acid varieties):** FBN Clethodim 240 may be tank-mixed with CURTAIL M for control of certain broadleaf and grass weeds in flax in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa.

Follow all precautions, limitations, restrictions and timing recommendations on the CURTAIL M label.

**Time of Application:** Flax may be treated from 5 to 15 cm in height. Apply when the grassy weeds are in the 2-6 leaf stage and the broadleaf weeds are actively

growing. Application to Canada thistle should be made when Canada thistle is between 10 cm and up to the pre-bud stage. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 L/ha tank-mixed with CURTAIL M at 1.5 to 2.0 L/ha and with the FBN Clethodim Adjuvant/ Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Weed claims listed on the FBN Clethodim 240 Herbicide label plus weed claims listed on Curtail M label. Refer to the CURTAIL M label for the broadleaf weeds controlled.

#### **Grazing Interval**

Do not cut treated crops for feed or graze until 60 days after application.

#### **MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- Add the required amount of CURTAIL M to the tank. Continue to agitate.
- Add the correct amount of FBN Clethodim 240. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/ Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
- If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

#### **FBN Clethodim 240 AND MCPA ESTER:**

**Flax:** FBN Clethodim 240 may be tank-mixed with MCPA ESTER for control of certain broadleaf and grassy weeds in flax in one spray operation. For ground applications, apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa. For aerial application apply this tank-mix in a minimum of 30 L/ha spray volume at a pressure of 235 kPa.

**NOTE: THIS TANK-MIX MAY NOT BE APPLIED TO LOW LINOLENIC ACID VARIETIES OF FLAX.**

**Time of Application:** Flax may be treated from 5 cm tall to just before the bud stage. Do not apply after the early bud stage. Apply when the grassy weeds are in the 2-6 leaf stage and broadleaf weeds are in the seedling stage and actively growing. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 - 0.38 L/ha tank-mixed with MCPA ESTER and the FBN Clethodim Adjuvant/ Companero Adjuvant at 0.5% - 1.0% v/v.

**Weeds Controlled:** As listed for FBN Clethodim 240 used alone, plus certain seedling broadleaf weeds. Refer to the MCPA ESTER label for the rates and broadleaf weeds controlled and specific information on flax.

#### MIXING INSTRUCTIONS:

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- Add the correct amount of MCPA ESTER. Continue to agitate.
- Add the correct amount of FBN Clethodim 240. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/ Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

#### FBN Clethodim 240 AND LONTREL® 360:

**Flax (including low linolenic acid varieties):** FBN Clethodim 240 may be tank-mixed with LONTREL 360 for control of certain broadleaf and grass weeds in flax in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa.

Follow all precautions, limitations and timing recommendations on the LONTREL 360 label.

**Time of Application:** Flax may be treated from 5 cm to 10 cm in height. Apply when the grassy weeds are in the 2-6 leaf stage and the broadleaf weeds are actively growing. Application to Canada thistle should be made when Canada thistle is in the rosette to pre-bud stage of growth. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 L/ha tank-mixed with LONTREL 360 at 0.56 to 0.83 L/ha and with the FBN Clethodim Adjuvant/ Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats, wild buckwheat.

Canada thistle	0.56 L LONTREL 360/ha	Season-long control
	0.83 L LONTREL 360/ha	Control extended into following year.

**MIXING INSTRUCTIONS:** For mixing instructions, refer to FBN Clethodim 240 and LONTREL 360 tank-mix on canola.

**Canola:** FBN Clethodim 240 may be tank-mixed with LONTREL 360 for control of certain broadleaf and grass weeds in canola in one spray operation. Apply in a minimum of 100 L/ha spray volume at a pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the LONTREL 360 label.

**Time of Application:** Canola may be treated from the 2 to 6 leaf stage. Apply when the grassy weeds are in the 2 to 6 leaf stage and the broadleaf weeds are actively growing. Application to Canada thistle should be made when Canada thistle is in the rosette to pre-bud stage of growth. If broadleaf weeds and grass weeds are not in recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 L/ha tank-mixed with LONTREL 360 at 0.42 to 0.83 L/ha and with the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats.

Wild Buckwheat	0.56 L LONTREL 360/ha	Season-long control
Canada thistle	0.42 L LONTREL 360/ha	Top growth control to 6 - 8 weeks
	0.56 L LONTREL 360/ha	Season-long control
	0.83 L LONTREL 360/ha	Control into following year.

**MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.



- Add the required amount of LONTREL 360 to the tank. Continue to agitate.
- Add the correct amount of FBN Clethodim 240. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
- If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

### **FBN Clethodim 240 AND PURSUIT®:**

**Field Peas:** FBN Clethodim 240 may be tank-mixed with PURSUIT for control of certain broadleaf and grassy weeds in field peas in one spray operation. Apply in a minimum of 100 litres of water per hectare at a spray pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the PURSUIT label.

**Time of Application:** Field peas may be treated up to the sixth (6<sup>th</sup>) trifoliate leaf stage. Apply when volunteer wheat, volunteer barley, wild oats and green foxtail are in the 2-6 leaf stage. Apply up to the 4-leaf stage for control of broadleaf weeds. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

**Application Rate:** FBN Clethodim 240 at 0.19 L/ha tank-mixed with PURSUIT at 210 mL/ha and the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Wild oats, green foxtail, volunteer barley, volunteer wheat plus the weeds as listed on the PURSUIT label.

### **MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- If tank-mixing FBN Clethodim 240 and PURSUIT for once over grass and broadleaf weed control in field peas, add the correct amount of PURSUIT. Continue to agitate.
- Add the required amount of FBN Clethodim 240 to the tank. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.

- Continue to agitate or run the by-pass system.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

**Imazethapyr Tolerant Canola (e.g. canola varieties with the SMART trait):** FBN Clethodim 240 may be tank-mixed with PURSUIT for control of certain broadleaf and grassy weeds in imazethapyr-tolerant canola varieties in one spray operation. Apply in a minimum of 100 litres of water per hectare at a spray pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the PURSUIT label.

**Time of Application:** Imazethapyr-tolerant canola varieties may be treated after the crop has developed one (1) fully expanded leaf. Apply when grassy weeds are in the 2 to 6-leaf stage. Apply up to the 4-leaf stage for control of broadleaf weeds.

**Application Rate:** FBN Clethodim 240 at 0.19L/ha tank-mixed with PURSUIT at 0.105 or 0.21 L/ha and the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** As listed for FBN Clethodim 240 plus the following additional weeds.

PURSUIT at 0.105 L/ha for control of chickweed, hempnettle, redroot pigweed\*, stinkweed, volunteer canola (not imazethapyr-tolerant), wild mustard and wild buckwheat\*, in the 1 to 4-leaf stage.

\*For the control of light infestations only. For heavy infestations, use the 0.21 L/ha rate of PURSUIT.

PURSUIT at 0.21 L/ha for control of weeds as listed on the PURSUIT label.

**MIXING INSTRUCTIONS:** For mixing instructions, refer to FBN Clethodim 240 and PURSUIT tank-mix on field peas.

#### **FBN Clethodim 240 AND MUSTER®:**

**Canola:** FBN Clethodim 240 may be tank-mixed with MUSTER for control of certain broadleaf and grassy weeds in canola in one spray operation. Apply in a minimum of 100 litres of water per hectare at a spray pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the MUSTER label.

**Time of Application:** Canola may be treated from the 2 leaf stage to the beginning of bolting. Apply when the grassy weeds are in the 2 to 6 leaf stage and when the

broadleaf weeds are in the cotyledon to 6 leaf stage. If broadleaf weeds and grassy weeds are not in the recommended treatment stage at the same time, then separate applications should be made.

**Application Rate: FBN Clethodim 240** at 0.19 L/ha tank-mixed with MUSTER at 20 or 30 g/ha and the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

### **Weeds Controlled:**

**Muster at 20 or 30 g/ha:** Wild oats, green foxtail, volunteer barley, volunteer oats, volunteer wheat, hempnettle, wild mustard, green smartweed, flixweed (spring seedlings), stinkweed\*, and redroot pigweed.

\* *Stinkweed suppression at the lower rate. Apply to stinkweed in the 1 to 4 leaf stage.*

### **MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- Add the required amount of MUSTER to the tank. Continue to agitate.
- Once MUSTER is in suspension, add the required amount of **FBN Clethodim 240**. Continue to agitate.
- Add the correct amount of FBN Clethodim Adjuvant/Companero Adjuvant along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system.
- On repeat tankloads, prepare a MUSTER/water slurry and add to required water volume in the spray tank prior to adding **FBN Clethodim 240** and FBN Clethodim Adjuvant/Companero Adjuvant.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight.
- Immediately after use, refer to MUSTER label for appropriate tank cleanout instructions.

### **FBN Clethodim 240 AND LIBERTY® 150 SN – PRAIRIE PROVINCES AND THE INTERIOR OF BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY**

**Canola:** For use on Glufosinate Ammonium tolerant canola varieties only (e.g. varieties or hybrids labelled LIBERTYLINK).

**FBN Clethodim 240** may be tank-mixed with LIBERTY 150 SN Herbicide for control of certain broadleaf and grass weeds in Glufosinate Ammonium tolerant canola varieties in one spray operation. For ground applications, apply in a minimum of 110

L/ha spray volume at a pressure of 275 kPa. For aerial applications apply this tank-mix in a minimum of 55 L/ha spray volume at a pressure of 300 kPa.

Follow all precautions, limitations and timing recommendations on the LIBERTY 150 SN Herbicide label.

**Time of Application:** Glufosinate Ammonium tolerant canola varieties may be treated from the cotyledon stage up to the early bolting stage. Apply when the wild oats, volunteer wheat, volunteer oats and green foxtail are in the 1 - 5 leaf stage. Refer to the LIBERTY 150 SN Herbicide label for application timing on all remaining grass and broadleaf weeds for 2.67 or 4.0 L/ha.

**Application Rate:** **FBN Clethodim 240** at 63 mL/ha tank-mixed with LIBERTY 150 SN Herbicide at 2.67 to 4.0 L/ha and with the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats plus all remaining grass weeds and broadleaf weeds listed on the LIBERTY 150 SN Herbicide label at the application rates of 2.67 and 4.0 L/ha.

#### **MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- Add the required amount of **FBN CLETHODIM ADJUVANT/COMPANERO ADJUVANT** to the tank. Continue to agitate until thoroughly mixed.
- STOP agitation. Add the correct amount of **LIBERTY 150 SN Herbicide** to the spray tank. Start agitation system.
- Add the correct amount of **FBN Clethodim 240** along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system and spray out immediately.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
- If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

**FBN Clethodim 240 AND LIBERTY® 200 SN – EASTERN CANADA AND BRITISH COLUMBIA (INCLUDING THE PEACE RIVER REGION) ONLY**

**Canola:** For use on Glufosinate Ammonium tolerant canola varieties only (e.g. varieties or hybrids labelled LIBERTYLINK).

**FBN Clethodim 240** may be tank-mixed with LIBERTY 200 SN Herbicide for control of certain broadleaf and grass weeds in Glufosinate Ammonium tolerant canola varieties in one spray operation. Apply in a minimum of 110 L/ha spray volume at a pressure of 275 kPa. Follow all precautions, limitations and timing recommendations on the LIBERTY 200 SN Herbicide label.

**Time of Application:** Glufosinate Ammonium tolerant canola varieties may be treated from the cotyledon stage up to the early bolting stage. Apply when the wild oats, volunteer wheat, volunteer oats and green foxtail are in the 1 – 5 leaf stage. Refer to the LIBERTY 200 SN Herbicide label for application timing on all remaining grass and broadleaf weeds for 2.0 or 2.5 L/ha.

**Application Rate:** **FBN Clethodim 240** at 63 mL/ha tank-mixed with LIBERTY 200 SN Herbicide at 2.0 to 2.5 L/ha and with the FBN Clethodim Adjuvant/Companero Adjuvant at 0.5% v/v.

**Weeds Controlled:** Wild oats, green foxtail, volunteer barley, volunteer wheat, volunteer oats plus all remaining grass weeds and broadleaf weeds listed on the LIBERTY 200 SN Herbicide label at the application rates of 2.0 and 2.5 L/ha.

#### **MIXING INSTRUCTIONS:**

- Thoroughly clean the sprayer by flushing the system with water containing detergent.
- Fill clean spray tank half full with clean water. Start agitation system.
- Add the required amount of **FBN CLETHODIM ADJUVANT/COMPANERO ADJUVANT** to the tank. Continue to agitate until thoroughly mixed.
- STOP agitation. Add the correct amount of **LIBERTY 200 SN Herbicide** to the spray tank. Start agitation system.
- Add the correct amount of **FBN Clethodim 240** along with the remaining amount of water necessary to fill the spray tank.
- Continue to agitate or run the by-pass system and spray out immediately.
- After any break in the spraying operation, agitate thoroughly before spraying again. Check inside the tank to ensure that sprayer agitation is sufficient to re-mix the spray materials. Do not allow the mixture to sit overnight. In the case of tank mixtures with broadleaf herbicides, settling will occur if agitation is not continuous.
- If an oil film starts to build up in the tank, drain it and clean tank with strong detergent solution.
- Immediately after use, thoroughly clean the sprayer by flushing the system with clean water containing detergent.

#### **MINOR USES:**

**NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS (BELOW):** The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Farmer's Business Network Canada, Inc. under the User Requested Minor Use Label Expansion program. For these uses, Farmer's Business Network Canada, Inc. has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

CROP	Weeds controlled	Application rate of FBN Clethodim 240	Application rate of FBN Clethodim Adjuvant/ Companero Adjuvant	REMARKS
<b>DRY BULB ONIONS</b>	Annual weeds listed on FBN Clethodim 240 label and suppression of annual bluegrass at the 2-6 leaf stage.	0.38 L/ha	0.5% v/v	<p><b>TIME OF APPLICATION:</b>            Apply <b>FBN Clethodim 240</b> post- emergent when the crop is in the 1- 4 leaf stage. Apply by ground application. One application per year.</p> <p>Observe a PHI of 45 days.</p> <p>Ground application only.</p>
<b>FENUGREEK</b>	Annual weeds listed on FBN Clethodim 240 label.	Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/ COMPANERO ADJUVANT weed/rate chart in this label.	Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/ COMPANERO ADJUVANT weed/rate chart in this label.	<p><b>TIME OF APPLICATION:</b>            Apply <b>FBN Clethodim 240</b> post- emergent when fenugreek is in the 3 – 5 leaf stage. Apply a maximum of one application per season, using ground equipment.</p> <p>Observe a PHI of 30 days.</p> <p>Do not graze or feed <b>FBN Clethodim 240</b>-treated fenugreek forage to livestock. Do not use the greens of treated fenugreek as a vegetable for human consumption.</p> <p>Ground application only.</p>
<b>CORIANDER, CARAWAY</b>	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 – 0.38 L/ha Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/ COMPANERO ADJUVANT weed/rate chart in this label.	0.5 – 1.0 % v/v Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/ COMPANERO ADJUVANT weed/rate chart in this label.	<p><b>TIME OF APPLICATION:</b>            Apply a maximum of one application per year, post-emergent using ground equipment. Apply in a minimum spray volume of 100 L/ha.</p> <p>Observe a PHI of 60 days.            Do not harvest leaves of treated coriander (cilantro) and caraway for human consumption.</p>

				Ground application only.
<b>SPINACH</b>	Annual weeds listed on FBN Clethodim 240 Label.	Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/COMPANERO ADJUVANT weed/rate chart in this label.	Refer to FBN Clethodim 240 + FBN CLETHODIM ADJUVANT/COMPANERO ADJUVANT weed/rate chart in this label.	<b>TIME OF APPLICATION:</b> Apply a maximum of two applications per year with 14 days between applications, post emergent by ground application when grassy weeds are in the 2-6 leaf stage.  Observe a PHI of 14 days.  Ground application only.

<b>CROP</b>	<b>Weeds controlled</b>	<b>Application rate of FBN Clethodim 240</b>	<b>Application rate of FBN Clethodim Adjuvant/Companero Adjuvant</b>	<b>REMARKS</b>
<b>PRAIRIE CARNATION</b>	Annual weeds listed on FBN Clethodim 240 label. Refer to weed/rate chart in this label.	0.19 L/ha	0.5% v/v	<b>TIME OF APPLICATION:</b> Apply <b>FBN Clethodim 240</b> post-emergence of weeds and crop. Apply a maximum of one application per year using ground equipment. Apply in a minimum spray volume of 110 L/ha. <b>Do not harvest treated prairie carnations for human consumption.</b>  Ground application only.
<b>HIGHBUSH BLUEBERRY</b>	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 – 0.38 L/ha	0.5 – 1.0 % v/v	<b>TIME OF APPLICATION:</b> Apply <b>FBN Clethodim 240</b> post-emergence of weeds and crop. Apply a maximum of one application per year, post-emergent by ground equipment. Apply as a broadcast spray directed to the ground. Apply in a minimum spray volume of 100 L/ha.  Observe a PHI of 14 days.  Ground application only.

<b>SAFFLOWER</b>	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 – 0.38 L/ha	0.5 – 1.0 % v/v	<p><b>TIME OF APPLICATION:</b> Apply FBN Clethodim 240 post-emergent when safflower is in the 6 – 8 leaf stage. Apply a maximum of one application per season, using ground equipment. Apply in a minimum spray volume of 100 litres of water per hectare.</p> <p>Observe a PHI of 70 days.</p> <p>Ground application only.</p>
<b>CRANBERRY</b>	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 – 0.38 L/ha	0.5 – 1.0 % v/v	<p><b>TIME OF APPLICATION:</b> Apply FBN Clethodim 240 post-emergence of weeds and crop. Do not apply between the hook stage and full fruit set. Apply a maximum of one application per year using ground equipment. Apply in a minimum spray volume of 110 litres of water per hectare.</p> <p>Observe a PHI of 30 days.</p> <p>Ground application only.</p>

<b>CROP</b>	<b>Weeds controlled</b>	<b>Application rate of FBN Clethodim 240</b>	<b>Application rate of FBN Clethodim Adjuvant/Companero Adjuvant</b>	<b>REMARKS</b>
<b>DILL</b>	Labelled grassy weeds. Refer to weed/rate chart in this label.	0.19 – 0.38 L/ha	0.5 – 1.0 % v/v	<p><b>TIME OF APPLICATION:</b> Apply FBN Clethodim 240 post-emergence when dill is in the 3-5 leaf stage. Apply a maximum of one application per year using ground equipment. Apply in a minimum spray volume of 100 L/ha.</p> <p>Do not harvest dillweed or dill seed within 40 days of application.</p> <p>Ground application only.</p>
<b>RED GARDEN (TABLE) BEET,</b>	Labelled grassy weeds. Refer to weed/rate chart	0.19 – 0.38 L/ha	0.5 – 1.0 % v/v	<p><b>TIME OF APPLICATION:</b> Apply FBN Clethodim 240 post-emergence of weeds and crop using ground</p>



<p><b>PARSNIP,          CARROT,          RADISH</b></p>	<p>in this label.</p>			<p>equipment. Apply a maximum of two applications per year. If repeat application is required, allow at least 14 days between first and second application. <b>Do not apply more than 0.38 L/ha (90 grams a.i./ha) per crop per season.</b> Apply in a minimum spray volume of 110 L/ha.</p> <p>Observe a PHI of 30 days.</p> <p>Ground application only.</p>
<p><b>BASIL</b></p>	<p>Labelled grassy weeds. Refer to weed/rate chart in this label.</p>	<p>0.19 L/ha</p>	<p>0.5% v/v</p>	<p><b>TIME OF APPLICATION:</b>                  Apply <b>FBN Clethodim 240</b> post- emergence of weeds and crop. Apply a maximum of one application per year using ground equipment. Apply as a broadcast spray directed to the ground. Apply in a minimum spray volume of 110 L/ha.</p> <p>Observe a PHI of 30 days.</p> <p>Ground application only.</p>

**APPLICATION INSTRUCTIONS:**

**GROUND APPLICATION:**

**Water volume and spray pressure:**

For ground application, use sprayers equipped with standard flat fan nozzles. For optimum spray coverage, use nozzles tilted 45° forward.

The use of flood jet or hollow cone nozzles is not recommended, because of uneven and inadequate spray coverage. Thorough and uniform spray coverage over the entire leaf area of the target weeds is necessary for optimum weed control.

Thoroughly clean all screens to prevent nozzle clogging, especially when water volumes of 55 to 110 L/ha are used.

For ground applications, use spray volumes of 55 to 225 L/ha and a minimum of 275 kPa pressure. For applications to dense weed infestations or to dense crop canopies, use a minimum of 110 litres of water per hectare at pressures of 275 to 425 kPa. In any situation, ensure the proper rate of FBN CLETHODIM ADJUVANT/COMPANERO

ADJUVANT is maintained as listed on the label under APPLICATION RATES OF **FBN Clethodim 240**.

**For dry common beans (*Phaseolus vulgaris* varieties only), a minimum spray volume of 110 L water/ha is recommended.**

#### **AERIAL APPLICATION:**

- Apply FBN Clethodim 240 alone or in tank mixes (ONLY with the recommended tank mix partners that are registered for aerial use) in no less than 28 L/ha spray volume at a pressure no less than 300 kPa.
- Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.
- Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended on this label.
- When applying **FBN Clethodim 240** by aircraft, uniform coverage is essential. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

#### **Aerial Use Precautions**

- Read and understand the entire label before opening this product. If you have questions, call Farmer's Business Network Canada, Inc. at 1-844-200-FARM (3276) or obtain technical advice from the distributor or your provincial agricultural representative.
- Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides.
- Do not apply to terrain where there is a potential for surface run-off to enter aquatic systems.
- Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed.
- Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift.
- Do not apply when wind speed is greater than 16 km/h at a flying height at the site of application.
- Do not apply with spray droplets smaller than ASAE medium classification.

#### **Aerial Operator Precautions:**

- Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.
- It is desirable that the pilots have communication capabilities at each treatment site at the time of application.

- The field crew and the mixer/loaders must wear a long-sleeved shirt, long pants, rubber apron, chemical-resistant gloves, goggles/face shield, socks and rubber boots. All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

**Product Specific Precautions:**

- Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-844-200-FARM (3276) or obtain technical advice from the distributor or your provincial agricultural representative.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply when wind speed is greater than 10 km/h at flying height at the site of application. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length **MUST NOT** exceed 65% of the wing- or rotorspan.

**Buffer zones:**

Spot treatments using hand-held equipment **DO NOT** require a buffer zone. The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands) and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands).

Method of application	Crop		Buffer Zones (meters) Required for the Protection of:		
			Freshwater Habitat of Depths		Terrestrial Habitat
			Less than 1m	Greater than 1m	
Field sprayer	All crops		1	1	1
Aerial	Canola, flax (including low linolenic acid varieties), field peas, lentils, potatoes, Oriental (brown) mustard (condiment and oilseed types), yellow mustard,	Fixed wing	5	1	40
		Rotary wing	5	1	35

	soybeans, sunflowers, Brassica carinata, dry bulb onions, garlic				
	Dry common beans, Desi and Kabuli chickpeas, pigeon peas	Fixed wing	1	1	20
		Rotary wing	1	1	20

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

#### ROTATIONAL CROP RESTRICTIONS

- A 30-day plant-back interval should be observed for all unlabelled crops.

#### RESTRICTIONS AND LIMITATIONS:

- Use only for crops listed on the label.
- Thorough preplant tillage operations are required to fields where sod or forage grass crops may have grown in the previous year.
- Take necessary precautions to avoid sprayer overlaps.
- Rainfall within one hour of application may reduce the effectiveness of the spray.
- Do not mix or apply **FBN Clethodim 240** with any other additive, pesticide or with fertilizer except as specifically recommended on the label.
- Allow 4 days between application of **FBN Clethodim 240** and any other chemical not recommended as a tank-mix combination on the label.
- Do not apply more than 0.38 L/ha (90 grams a.i./ha) per season.
- Do not use crops for grazing of livestock or green feed following application of **FBN Clethodim 240** until after the appropriate interval as specified in the crop recommendations table has been observed. Do not cut treated crops for forage until the appropriate interval as specified in the crop recommendations table has been observed.
- FBN Clethodim 240 may be applied by air for certain crops specified in this label.
- Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.
- Not for use in greenhouses.

#### HERBICIDE RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, FBN Clethodim 240 is a Group 1 herbicide. Any weed population may contain or develop plants naturally resistant to FBN Clethodim 240 and other Group 1 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of FBN Clethodim 240 or other Group 1 herbicides within a growing (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Farmer's Business Network Canada, Inc. or phone 1-844-200-FARM (3276).

**NOTICE TO USER:** This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

BUCTRIL® is a registered trademark of Bayer.

LONTREL® 360 and CURTAIL are registered trademarks of Dow AgroSciences LLC.

MUSTER® is a registered trademark of FMC Corporation.

LIBERTY® and PURSUIT® are registered trademarks of BASF.

COMPANERO™ is a trademark of Norac concepts Inc.