

Tribe 75 WDG

Herbicide

WETTABLE GRANULES

FOR SALE FOR USE IN THE PRAIRIE PROVINCES, PEACE RIVER REGION AND THE INTERIOR OF BRITISH COLUMBIA ONLY AS A PRE-SEEDING HERBICIDE ON FIELDS TO BE PLANTED TO WHEAT (SPRING AND DURUM) AND BARLEY, AND FOR WEED CONTROL IN SUMMERFALLOW
FOR SALE FOR USE IN EASTERN CANADA FOR WEED CONTROL IN LOWBUSH BLUEBERRIES IN EASTERN CANADA ONLY

COMMERCIAL

READ THE LABEL AND ENCLOSED BOOKLET BEFORE USING

ACTIVE INGREDIENT: TRIBENURON-METHYL 75%

REGISTRATION NO.: 34345 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.08 kg – 10 kg

WARNING - EYE IRRITANT- POTENTIAL SKIN SENSITIZER

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes eye irritation. DO NOT get in eyes
- POTENTIAL SKIN SENSITIZER
- DO NOT APPLY BY AIR
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.
- AVOID BREATHING SPRAY MIST OR VAPOURS. USE WITH ADEQUATE VENTILATION.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- WEAR A LONG-SLEEVED SHIRT, LONG PANTS, CHEMICAL-RESISTANT GLOVES, SOCKS AND SHOES DURING MIXING, LOADING, APPLICATION, CLEAN-UP AND REPAIR. IN ADDITION, WEAR PROTECTIVE EYEWEAR (GOGGLES OR FACE SHIELD) DURING MIXING AND LOADING. GLOVES ARE NOT REQUIRED DURING APPLICATION WITHIN A CLOSED CAB.
- WHEN USING, DO NOT EAT, DRINK OR SMOKE.
- WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE.
- STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE RE-USE
- WASH THOROUGHLY WITH SOAP AND WATER AFTER USE.
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS DURING THE RESTRICTED ENTRY INTERVAL (REI) OF 12 HOURS.
- APPLY ONLY WHEN THE POTENTIAL FOR DRIFT TO AREAS OF HUMAN HABITATION OR AREAS OF HUMAN ACTIVITY SUCH AS HOUSES, COTTAGES, SCHOOLS AND RECREATIONAL AREAS IS MINIMAL. TAKE INTO CONSIDERATION WIND SPEED, TEMPERATURE INVERSIONS, APPLICATION EQUIPMENT AND SPRAYER SETTINGS.

FIRST AID:

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

STORAGE:

Store this product away from food or feed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in any way that is inconsistent with the directions on the label.

<<DFU Booklet>>

Tribe 75 WDG

Herbicide

WETTABLE GRANULES

FOR SALE FOR USE IN THE PRAIRIE PROVINCES, PEACE RIVER REGION AND THE INTERIOR OF BRITISH COLUMBIA ONLY AS A PRE-SEEDING HERBICIDE ON FIELDS TO BE PLANTED TO WHEAT (SPRING AND DURUM) AND BARLEY, AND FOR WEED CONTROL IN SUMMERFALLOW
FOR SALE FOR USE IN EASTERN CANADA FOR WEED CONTROL IN LOWBUSH BLUEBERRIES IN EASTERN CANADA ONLY

COMMERCIAL

READ THE LABEL AND ENCLOSED BOOKLET BEFORE USING

ACTIVE INGREDIENT: TRIBENURON-METHYL 75%

REGISTRATION NO.: 34345 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 0.08 kg – 10 kg

WARNING - EYE IRRITANT- POTENTIAL SKIN SENSITIZER

Sharda Cropchem Limited
2nd Floor, Prime Business Park
Dashrathlal Joshi Road
Vile Parle (West)
Mumbai - 400056, India

Canadian Agent:
Sharda Cropchem Limited
63 Kingsview Blvd
Etobicoke, Ontario, CA
M9R1V1
1-844-810-5720
1-416-840-5639

PRECAUTIONS:

- **KEEP OUT OF REACH OF CHILDREN.**
- Causes eye irritation. DO NOT get in eyes
- POTENTIAL SKIN SENSITIZER
- DO NOT APPLY BY AIR
- MAY BE HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN.
- AVOID BREATHING SPRAY MIST OR VAPOURS. USE WITH ADEQUATE VENTILATION.
- DO NOT CONTAMINATE ANY BODY OF WATER.
- WEAR A LONG-SLEEVED SHIRT, LONG PANTS, CHEMICAL-RESISTANT GLOVES, SOCKS AND SHOES DURING MIXING, LOADING, APPLICATION, CLEAN-UP AND REPAIR. IN ADDITION, WEAR PROTECTIVE EYEWEAR (GOGGLES OR FACE SHIELD) DURING MIXING AND LOADING. GLOVES ARE NOT REQUIRED DURING APPLICATION WITHIN A CLOSED CAB
- WHEN USING, DO NOT EAT, DRINK OR SMOKE.
- WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING. REMOVE CONTAMINATED CLOTHING IMMEDIATELY AFTER USE.
- STORE AND WASH CONTAMINATED CLOTHING SEPARATELY FROM HOUSEHOLD LAUNDRY BEFORE RE-USE
- WASH THOROUGHLY WITH SOAP AND WATER AFTER USE.
- DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS DURING THE RESTRICTED ENTRY INTERVAL (REI) OF 12 HOURS.
- APPLY ONLY WHEN THE POTENTIAL FOR DRIFT TO AREAS OF HUMAN HABITATION OR AREAS OF HUMAN ACTIVITY SUCH AS HOUSES, COTTAGES, SCHOOLS AND RECREATIONAL AREAS IS MINIMAL. TAKE INTO CONSIDERATION WIND SPEED, TEMPERATURE INVERSIONS, APPLICATION EQUIPMENT AND SPRAYER SETTINGS.

FIRST AID:

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

IF ON SKIN OR CLOTHING:

Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

IF SWALLOWED:

Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

IF INHALED:

Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION: Treat symptomatically.

IMPORTANT INFORMATION - READ BEFORE USING

Injury to or loss of desirable trees, shrubs, flowers, vegetables, fruits or other vegetation may result from failure to observe the following: Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on crops.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

As this product is not registered for control of pests in aquatic systems, DO NOT use to control aquatic pests.

DO NOT APPLY BY AIR.

Do not apply in high winds.

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than spring wheat, durum or barley. Thoroughly clean all traces of tank mix from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Wash away any spray mixture from the outside of spray tank, nozzle or spray rig.

A MINIMUM RECROPPING INTERVAL OF 2 MONTHS SHOULD BE LEFT BETWEEN THE APPLICATION OF THIS PRODUCT AT THE 10 GRAM PER HECTARE RATE AND SEEDING OF THE NEXT CROP. After 2 months, the following crops can be seeded: canola, flax, lentils, alfalfa.

DO NOT APPLY TRIBE 75 WDG WITH A CARRIER OTHER THAN WATER OR ANY ADDITIVE OTHER THAN THOSE INDICATED ON THIS LABEL. Other carriers may accelerate breakdown of this product and reduce its effectiveness.

ENVIRONMENTAL PRECAUTIONS:

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative filter strip between the treated area and the edge of the water body.

TOXIC to aquatic organisms and non-target terrestrial plants. Observe spray buffer zones specified under **DIRECTIONS FOR USE**.

DIRECTIONS FOR USE

GENERAL INFORMATION:

Tribe 75 WDG provides control of certain broad-leaved weeds as a pre-seeding herbicide on fields to be planted to wheat (including durum) and barley.

Warm, moist growing conditions promote active weed growth and enhance the activity of Tribe 75 WDG by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress such as cold weather, drought stress or excessive heat may not be adequately controlled or suppressed, and regrowth may occur. Tribe 75 WDG may only be applied using ground spray equipment. For best results, ensure thorough spray coverage of target weeds.

Tribe 75 WDG rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application.

When tank-mixes are permitted, read and observe all label directions, including rates and restrictions for each product used in the tank-mix. Follow the more stringent label precautionary measures for mixing, loading and applying stated on both product labels.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

Spray Buffer Zones:

A spray buffer zone is NOT required for uses with hand-held application equipment permitted on this label.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Spray Buffer Zones (meters) Required for the Protection of:		
		Aquatic Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field Sprayer	summerfallow, fields to be planted to wheat (spring and durum), barley (only as a pre-seeding	0	0	3

	herbicide)			
	Wheat and barley	1	0	4
	Lowbush blueberries (Eastern Canada only)	1	0	10

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pest Management Regulatory Agency web site.

FOR WEED CONTROL IN SUMMERFALLOW

With any summerfallow treatment, allow at least 10 days to elapse between treatment and tillage. Only weeds that have emerged at the time of application will be controlled.

APPLICATION TIMING:

Tribe 75 WDG + 2,4-D Ester Tank Mix

Apply Tribe 75 WDG at 10 grams per hectare with 2,4-D ester at 420 grams per hectare (0.6 L 2,4-D Ester LV-700).

Apply the tank mix in the spring after emergence, up to the early flowering stage of the target weeds.

WEEDS CONTROLLED:

Flixweed and Stinkweed (fall rosettes and spring seedlings)

CONTROL OF BROADLEAF WEEDS AND GRASSES IN FALLOW AND PRE-SEED TO SPRING WHEAT (INCLUDING DURUM) AND BARLEY.

With any summerfallow treatment, allow 10 days to elapse between treatment and tillage. Only weeds emerged at time of application will be controlled.

Fields treated with Tribe 75 WDG can be seeded to wheat (spring or durum) or barley a minimum of 24 hours after application.

Apply Tribe 75 WDG at 10 grams per hectare, tank mixed with one of the following companion herbicides and adjuvant (where required) in a total spray volume of 55-110 L/ha.

Companion Herbicide	Application Rate	Application Stage	Weeds Controlled	Weeds Suppressed*
Roundup Transorb or Roundup Weathermax	1.25 L/ha 0.83 L/ha	Up to 8 cm	Canada fleabane, Common ragweed,	

			Narrow- leaved hawk's beard	
		Up to 15 cm	Dandelion, Downy brome, Flixweed, Giant foxtail, Green foxtail, Hemp nettle, Kochia, Lady's thumb, Lamb's-quarters, Persian darnel, Redroot pigweed, Russian thistle, Stinkweed, Volunteer barley, Volunteer canola (including glyphosate-tolerant varieties), Volunteer flax, Volunteer wheat, Wild mustard, Wild oats	
		Up to 3-leaf	Wild buckwheat, Cow cockle	
		Rosette		Canada thistle
Roundup Transorb, plus Agral 90	0.75 L/ha 0.35% v/v	Any stage	Volunteer canola (including glyphosate tolerant varieties)	
		Seedlings and rosettes (nonflowering)		Narrow-leaved hawk's beard
		Up to 10 cm	Kochia, Lamb's-quarters	Redroot pigweed, Russian thistle
		Up to 8 cm	Green foxtail, Lady's thumb, Stinkweed, Volunteer barley, Volunteer wheat, Wild mustard, Wild oats (1-3-leaf), Wild buckwheat (1-3 leaf)	
Roundup Transorb, plus Agral 90	1.0 L/ha 0.35% v/v	Rosettes (nonflowering)	Dandelion (top growth)	Canada thistle (top growth)

*Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

Fields treated with Tribe 75 WDG + Roundup herbicides can be seeded with wheat (spring or durum) or barley a minimum of 24 hours after application.

LOWBUSH BLUEBERRY – Eastern Canada Only

For bunchberry control, apply Tribe 75 WDG at 40 g/ha with 0.2% v/v AGRAL 90* (2 L per 1000 L of water) in the spring of the sprout year**. Apply in 150 to 250 L water per hectare.

For best results, application should be made when the majority of the emerged bunchberry plant leaves have unfolded to form a 45 degree angle, but no later than when the first white blossoms are visible on the most advanced plants. Tribe 75 WDG + AGRAL* 90 should be applied before blueberry sprout regrowth exceeds 2 cm in height. Applications made at later stages of blueberry development or applications made in spring burnt fields are not recommended due to potential crop injury and yield reductions. Blueberry plant stunting can result following Tribe 75 WDG + AGRAL* 90 application, but they recover and fruit bud numbers and yields should not be affected. Recommended fertilizer applications following Tribe 75 WDG + AGRAL* 90 applications can be beneficial.

Control will not be achieved if AGRAL 90 is not applied with Tribe 75 WDG.

**Make only one application per crop cycle (2 or 3 years) in the sprout year.

Tribe 75 WDG is taken up mainly through actively growing plant tissue, thus no specific soil restrictions apply.

NOTE TO USER: READ THE FOLLOWING BEFORE USING THIS PRODUCT FOR THE INDICATED SPECIAL USE APPLICATIONS:

The DIRECTIONS FOR USE for the uses described in this section of the label were developed by persons other than Sharda Cropchem Limited under the User Requested Minor Use Label Expansion program. For these uses, Sharda Cropchem Limited has not fully assessed performance (efficacy) and/or crop tolerance (phytotoxicity) under all environmental conditions or for all crop varieties when used in accordance with the label. The user should test the product on a small area first, under local conditions and using standard practices, to confirm the product is suitable for widespread application.

LOWBUSH BLUEBERRY: Bunchberry control – fall application For bunchberry control, apply Tribe 75 WDG at 40 g/ha with 0.2% v/v AGRAL 90* (2 L per 1000 L of water) 1-4 weeks after completion of harvest. Apply in 150 to 250 L water per hectare. Do not apply more than once in a crop cycle.

LOWBUSH BLUEBERRY: To control yellow loosestrife, speckled alder, wild rose, and bracken fern in lowbush blueberries, mix 0.25 grams Tribe 75 WDG per litre of water plus AGRAL* 90 at a rate of 0.2% v/v (2 L per 1000 L spray solution).

Apply Tribe 75 WDG as a spot treatment. Application should be made in mid summer of the prune or vegetative year of the crop to fully expanded foliage of bracken fern and yellow loosestrife. Wild rose and speckled alder may also be controlled by early fall applications. Apply enough spray solution to thoroughly wet the weed foliage. Stunting and reduction of fruit buds may occur if the crop is sprayed directly.

MIXING INSTRUCTIONS:

1. Add one-third of the necessary volume of water to the spray tank.
2. With the agitator running, add the Tribe 75 WDG.
3. When Tribe 75 WDG is in suspension, add the required amount of the appropriate tank mix partners.
4. Specific Tank Mix Instructions:

Tribe 75 WDG + 2,4-D Ester: Once Tribe 75 WDG is completely in suspension, add the required amount of 2,4-D Ester while maintaining agitation. Do not add a surfactant.

Tribe 75 WDG + Roundup Transorb or Roundup Weathermax: Once Tribe 75 WDG is completely in suspension, add the required amount of Roundup Transorb or Roundup Weathermax as indicated in the APPLICATION RATES AND TIMING tables. Once components are in suspension, if required, add AGRAL 90 at 3.5 L per 1000 L of spray solution (0.35% v/v).

1. Add the rest of the water. If an antifoam agent is required, add this last.
2. Use a minimum spray volume of 55 L/ha.
3. For repeat tank loads, reduce the material remaining in the tank to 10% of the original volume or less before proceeding with step 1. If this is not possible, pre-slurry Tribe 75 WDG in a small amount (5-10 L) of water before adding to the tank.

Continuous agitation is required to keep Tribe 75 WDG in suspension. If the spray preparation is left standing without agitation, thoroughly agitate to re-suspend the mixture before spraying. Use spray preparation within 24 hours or product degradation may occur.

MINIMUM RECROPPING INTERVAL AND CROP ROTATION GUIDELINES:

A minimum recrop interval of 2 months should be left between the application of this product at the 10 g/ha rate and seeding of the next crop. The following crops can be seeded two months after application of Tribe 75 WDG + 2,4-D Ester: canola, flax, lentils, and alfalfa.

EQUIPMENT-SPRAY VOLUMES:

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep the spray materials in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Reduce drift with high water volume, low pressure, coarse sprays and drop nozzles.

SPRAYER CLEANUP:

It is difficult to remove all traces of 2,4-D Ester from a sprayer. A sprayer used to apply this herbicide should not be used to spray crops sensitive to this herbicide. The following clean-out procedure will remove traces of Tribe 75 WDG from the sprayer.

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes to remove all visible residues.
2. Fill the tank with clean water then add 1 litre household AMMONIA (containing a minimum of 3% ammonia) per 100 litres of water. Fill boom and hoses with solution and allow sprayer to sit for 15 minutes. Drain.
3. Repeat step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of ammonia, rinse the tank, hoses and booms thoroughly with clean water.
5. Dispose of tank rinseate according to Provincial directions.

CAUTION: Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour which can cause eye, nose, throat and lung irritation. Do not clean equipment in an enclosed area.

RESISTANCE MANAGEMENT RECOMMENDATIONS:

For resistance management, Tribe 75 WDG is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Tribe 75 WDG and other Group 2 herbicides. The resistance biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible rotate the use of Tribe 75 WDG or other Group 2 herbicides within a growing season (sequence) or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from different groups when such use is permitted. To delay resistance, the less resistance-prone partner should control the target weed(s) as effectively as the more resistance-prone partner.
- Herbicide use should be based on an integrated weed management program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical control methods), cultural (for example, higher crop seeding rates; precision fertilizer application method and timing to favour the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Monitor weed populations after herbicide application for signs of resistance development (for example, only one weed species on the herbicide label not controlled). If resistance is suspected, prevent weed seed production in the affected area if possible by an alternative herbicide from a different group. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- Have suspected resistant weed seeds tested by a qualified laboratory to confirm resistance and identify alternative herbicide options.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact your local Sharda Cropchem Limited representative at 1-844-810-5720 for further information.

USE PRECAUTIONS:

Licensed wheat and barley varieties have shown acceptable tolerance to Tribe 75 WDG. Consult Sharda Cropchem Limited prior to use on unlicensed varieties for crop tolerance information.

Do not apply to irrigated land where tail water will be used to irrigate other crop land.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

As this product is not registered for control of pests in aquatic systems, DO NOT use to control aquatic pests.

Use a maximum of 10 grams of Tribe 75 WDG per hectare per year in summerfallow and on fields to be planted to wheat and barley.

Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.

Do not apply to fields to be planted to spring wheat, durum wheat or barley under sown with legumes or grasses as injury to the forage may result.

Tribe 75 WDG applied to fields to be planted to spring wheat, durum wheat or barley that is stressed by severe conditions such as drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury, grade or yield loss.

Do not remove soil from treated fields for use on lawns, in gardens or for backfill.

Under certain conditions such as prolonged cool weather temperature or wide fluctuations in day/night temperatures just prior to or soon after treatment, temporary yellowing and/or crop stunting may occur.

OVERSPRAY OR DRIFT TO IMPORTANT WILDLIFE HABITATS SUCH AS SHELTERBELTS, WETLANDS, SLOUGHS OR DRY SLOUGH BORDERS, OR WOODLOTS, SHOULD BE AVOIDED. LEAVE A 15 METER SPRAY BUFFER ZONE BETWEEN THE LAST SPRAY SWATH AND THE EDGE OF ANY OF THESE HABITATS.

Control of weeds growing in wheel tracks may be reduced Tribe 75 WDG is applied under dry, dusty conditions.

STORAGE:

Store this product away from food or feed.

DISPOSAL:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER:

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in any way that is inconsistent with the directions on the label.