

155(S)082404A11M1

GAUCHO[®] 480

FLOWABLE

ACTIVE INGREDIENTS:

Imidacloprid:1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine.....40.7%

OTHER INGREDIENTS:.....59.3%

TOTAL.....100.0%

Contains 4 lbs. active per gallon (480 grams per liter) @ 20°C

EPA Reg. No. 7501- 155

EPA Est. 7501-ID-1

NET CONTENTS: 2.5 GALLONS

KEEP OUT OF REACH OF CHILDREN

CAUTION

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

SEE ADDITIONAL PRECAUTIONARY STATEMENTS

A quality product manufactured by:



Gustafson LLC
1400 Preston Rd., Suite 400
Plano, TX 75093
Questions? (972) 985-8877

KEEP OUT OF REACH OF CHILDREN**CAUTION****PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND
DOMESTIC ANIMALS**

Harmful if swallowed, absorbed through skin or inhaled. Avoid breathing vapor or spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

PERSONAL PROTECTIVE EQUIPMENT (PPE): Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for Category C on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear: Long-sleeved shirt and long pants, socks and shoes and chemical-resistant gloves (such as nitrile, butyl, neoprene, barrier laminate, polyvinyl chloride or Viton). Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS: Users should wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Users should remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change clothing.

FIRST AID

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.

**IN CASE OF AN EMERGENCY ENDANGERING LIFE OR PROPERTY INVOLVING THIS
PRODUCT CALL:**

CHEMTREC: (800) 424-9300 (Toll free – US)

CHEMTREC: (703) 527-3887 (Domestic and International)

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds and aquatic invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Cover or incorporate spilled treated seeds.

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

For use in commercial seed treaters only, with the exception of application to canola, cotton (delinted seed), field corn, sorghum, wheat and barley, which may be made either by commercial seed treatment or as an end-use seed treatment on agricultural establishments at, or immediately before, planting. This product is to be used in liquid or slurry treaters.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Mix thoroughly before use or use entire container at one time. All tank mixes should be pre-tested to determine physical compatibility between formulations. Observe all cautions and limitations on labeling of all products used in mixtures.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the seed is treated with the product and the treated seed is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls, waterproof gloves, shoes plus socks, protective eyewear.

Stored Seed Protection:

Applied at labeled rates equal to or above 1.0 fl. oz. per hundredweight, GAUCHO 480 Flowable will provide protection to seed against injury from the following insects: Indian Meal Moth (*plodia interpunctella*), Rice Weevil (*Sitophilus oryzae*), Red Flour Beetle (*Tribolium castaneum*), and Lesser Grain Borer (*Rhizopertha dominica*). It is recommended that seed with existing populations of stored grain pests be fumigated prior to treating and bagging seed.

For Early Season Protection Against Certain Sucking Insects:

GAUCHO 480 Flowable will aid in the protection of seeds and seedlings against injury by certain early season insects.

Wheat, Barley, Triticale:

To provide early season protection of seedlings against injury by aphids (including Bird cherry-oat, English grain, Greenbug, and Russian wheat aphid), Hessian fly and Wireworms, apply 1.0 to 3.0 fluid ounces per hundredweight of seed prior to planting as a slurry treatment. Low use rate for "wireworm-only" protection: GAUCHO 480 applied at the 0.16 - 0.32 fl. oz. per hundredweight of seed offers suppression of wireworm activity on seed and young seedlings. To provide early season protection from grasshopper damage, GAUCHO 480 should be applied to seed at 1.5 to 3.0 fluid ounces per hundredweight. To reduce early season damage caused by grasshopper, GAUCHO treated seed may be planted as a 50 to 60 foot border around the edges of the field. Consult your local university extension entomologist for details regarding grasshopper control in your area. For maximum effectiveness, seed should be treated uniformly. Ensure thorough coverage. Use the higher rate to provide increased length of protection and from heavy insect pressure and from wireworms, and to reduce potential spread of Barley yellow dwarf virus due to aphid vectors. Do not graze or feed livestock on treated areas for 45 days after planting.

Sorghum:

To provide early season protection of seedlings against injury by Aphids (including Corn leaf, English grain, Greenbug, and Yellow sugar cane aphid), Chinch bugs, Fire ants and Wireworms, apply 8 fluid ounces per hundredweight of seed prior to planting as a slurry treatment. Ensure thorough coverage. Do not graze or feed livestock on treated areas for 45 days after planting.

Sugar Beets:

To provide early season protection of seedlings against injury by Whitefly, Aphids, Leafhoppers (including to reduce potential for spread of Curly top and Yellow mosaic virus due to aphid and leafhopper vectors), Root

aphid, Thrips and Wireworms, apply as a commercial seed treatment at 3.0 to 6.3 fluid ounces of GAUCHO 480 Flowable in or on a unit of pelleted sugar beet seed with a weight ratio of 2:1 pelleting mixture to raw seed (seed count 100,000 seed - approximately 1 kilogram by weight). Apply in a film coat directly to raw seed (100,000 seed or approximately 1 kilogram by weight) at a rate of 3.0 fluid ounces per unit of seed. If rates exceed 3.0 fluid ounces per unit, seed must be pelleted.

Cotton (Delinted Seed Only):

To provide protection of seedlings against injury by early season thrips and aphids and where specific application rate is desired on an individual seed basis apply 0.375 mg. a.i. per seed (do not apply more than 16.0 fluid ounces per hundredweight of seed) prior to planting as a slurry treatment so as to ensure thorough coverage. Otherwise, apply at 16.0 fluid ounces per hundredweight of seed. Regardless of the type of application (seed treatment, soil or foliar) do not apply more than a total of 0.5 lb. of imidacloprid per acre per cropping cycle.

Sweet Corn:

(Not registered for use in California)

For the protection of sweet corn plants from the insect pests listed below. Apply as a seed treatment at the recommended rates listed.

PEST	GAUCHO 480 FL. OZ. / CWT. OF SEED
Flea beetle Early season corn leaf aphid Seed corn maggot Wireworm	8
Imported fire ant Early season corn leaf aphid Seed corn maggot Wireworm	4-8
Imported fire ant Seed corn maggot (seed protection) Wireworm (seed protection)	2-4
Wireworm (seed protection)	1-2

The final slurry rate should be adjusted to apply a rate of 16 - 20 fl. oz. of dilute solution per hundredweight of seed with commercial application equipment.

Asparagus Bean, Broad Bean (Succulent Or Dry), Chinese Longbean, Field Bean, Guar Bean, Jackbean, Kidney Bean, Lima Bean (Succulent Or Dry), Moth Bean (Succulent Or Dry), Mung Bean, Navy Bean, Pinto Bean, Rice Bean, Runner Bean, Snap Bean, Sword Bean, Wax Bean, Yardlong Bean, Blackeyed Pea (Succulent Or Dry), Chickpea, Cowpea (Succulent Or Dry), Crowder Pea, Dwarf Pea, Edible-Pod Pea, English Pea, Field Pea, Garden Pea, Green Pea, Pigeon Pea (Succulent Or Dry), Snow Pea, Southern Pea (Succulent Or Dry), Sugar Snap Pea, Lentil: *

To provide early season protection of seedlings against injury by wireworm, bean leaf beetle, imported fire ant, and aphid, apply as a commercial seed treatment at 2 - 4 fl. oz. per hundredweight of seed prior to planting.

* Seed-and-pod vegetable seed treated in California must be destined for planting in states other than California and is not to be planted in California.

ROTATIONAL CROPS:

Treated areas may be replanted with any crop specified on an imidacloprid label, or any crop for which a tolerance exists for the active ingredient, as soon as practical following the last application. The following plant-back intervals should be observed for listed crops:

ROTATIONAL PLANT-BACK INTERVALS*				
IMMEDIATE PLANT-BACK				30-DAY PLANT-BACK
Artichoke	Cranberry	Mustard Seed	Strawberry	Cereals, including: buckwheat millet oats rice rye Safflower Soybean
Barley	Cucurbits	Okra	Sugarbeet	
Brassica (cole) leafy veg.	Eggplant	Pepinos	Tomatillo	
Canola	Ground cherry	Pepper	Tomato	
Cilantro	Leaf petiole veg.	Popcorn	Triticale	
Corn, Field	Leafy veg.	Potato	Veg. Root and Tuber	
Corn, Sweet	Legume veg. (succulent or dried, except soybean)	Rapeseed	Watercress	
Cotton		Sorghum	Wheat	

- * Cover crops for soil building or erosion control may be planted any time, but do not graze or harvest for food or feed.
- * For all other crops not listed on an Imidacloprid label, or for crops for which no tolerance for the active ingredient has been established, a 12-month plant-back interval should be observed.

NOTIFICATION of the crop rotational restriction must be conveyed to the grower by appropriate seed tag labeling or bag printing on all seed units.

NOTE: The purchaser of this product is responsible for ensuring that all seed treated with this product are adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.1001. Federal regulations have established official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

Treated seed must not be used for or mixed with food or animal feed or processed for oil. Seed commercially treated with GAUCHO 480 Flowable must be labeled in accordance with all applicable requirements of the Federal Seed Act.

Labels for commercially treated seed should include the following addition to the Environmental Hazards statements:

Exposed treated seed may be hazardous to birds. Dispose of all excess treated seed and seed packaging by burial away from bodies of water. Cover or incorporate spilled treated seeds.

USE RESTRICTION:

Rape greens grown and harvested from GAUCHO 480 treated seed must not be used for human and feed consumption. Rapeseed grown and harvested from GAUCHO 480 treated seed is only for industrial uses and can not be used for edible oil or any other human / feed consumption.

STORAGE AND DISPOSAL

STORAGE: Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures.

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal, or by cleaning of equipment. Do not reuse empty container. Open dumping is prohibited.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Contact Gustafson or your chemical supplier for container return instructions. If container is not returned in accordance with instructions, triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTICE – READ CAREFULLY

CONDITIONS OF SALE AND LIMITED WARRANTY STATEMENT

Treatment of highly mechanically damaged seed, or seed of known low vigor and poor quality, may result in reduced germination and/or reduction of seed and seedling vigor. Treat and conduct germination tests on a small portion of seed before committing the total seed lot to a selected chemical treatment. Due to seed quality conditions beyond the control of Gustafson LLC, no claims are made to guarantee germination of carry-over seed.

As its sole express warranty, Gustafson LLC warrants that this product conforms to the chemical description on the label and is reasonably fit for purposes stated on the label only when used in accordance with directions and instructions specified on the label, subject to the inherent risks set forth above. ONLY AS PERMITTED BY APPLICABLE LAW, IN THE EVENT OF A BREACH OF THIS LIMITED WARRANTY, GUSTAFSON LLC SHALL NOT BE LIABLE FOR CONSEQUENTIAL DAMAGES. SUBJECT TO APPLICABLE LAW, GUSTAFSON LLC NEITHER MAKES NOR AUTHORIZES ANY OF ITS AGENTS OR REPRESENTATIVES TO MAKE ANY WARRANTY OF FITNESS OR MERCHANTABILITY, GUARANTY OR REPRESENTATION, EXPRESS OR IMPLIED, CONCERNING THIS MATERIAL. Buyer assumes the responsibility to handle, use, and store this product in accordance with safety instructions and use directions contained on the label.

Buyer/User purchases this product subject to the foregoing Conditions of Sale and Limited Warranty which, subject to applicable law, may be varied only by a written agreement signed by a duly authorized representative of Gustafson LLC. If these terms are not acceptable, return all products to the place of purchase, unopened for a full refund.

Gustafson (logo) is a registered trademark of Gustafson LLC, Plano, Texas
GAUCHO is a registered trademark of Bayer Ag, Germany

RTU is a registered trademark of Gustafson LLC, Plano, Texas
VITAVAX is a registered trademark of Crompton Corporation
ALLEGIANCE is a registered trademark of Gustafson LLC, Plano, Texas