

Keylate[®] CoMo Classic



SPECIMEN

3-0-0 GUARANTEED ANALYSIS

Total Nitrogen (N)	3.0%
3.0% Water Soluble Nitrogen	
Cobalt (Co)	2.0%
2.0% Chelated Cobalt	
Molybdenum (Mo)	3.0%

Derived from cobalt EAHP and sodium molybdate.

F2399

Z-KEYCOMO

Rev: 21C10

Density: 10.3 lb/gal (1.23 kg/L) @ 68°F (20°C)

NET CONTENT

- 2.50 gal (9.5 L)
- 2.64 gal (10.0 L)
- 5.00 gal (18.9 L)
- 5.28 gal (20.0 L)
- 55.00 gal (208.2 L)
- 275.00 gal (1041.0 L)
- BULK

NET WEIGHT

- 25.8 lb (11.7 kg)
- 27.2 lb (12.4 kg)
- 51.5 lb (23.4 kg)
- 54.4 lb (24.7 kg)
- 566.5 lb (257.3 kg)
- 2832.5 lb (1286.7 kg)

CAUTION

KEEP OUT OF REACH OF CHILDREN; MAY BE HARMFUL IF SWALLOWED, INHALED OR IN CONTACT WITH SKIN; MAY CAUSE EYE IRRITATION. MAY CAUSE DAMAGE THROUGH PROLONGED EXPOSURE.

PRECAUTIONS:

Do not breathe dust/fume/gas/mist/vapors/spray. Wash hands thoroughly after handling. Do not eat, drink or smoke when using this product. Use only outdoors or in well-ventilated area. Avoid release to environment. Wear protective gloves/protective clothing/eye protection/face protection. Keep container tightly closed. Keep cool and protected from sunlight.

If medical advice is needed, have product container or label at hand. For more details regarding the safe use/handling, storage, and disposal of this product as well as additional hazards and first aid measures, please obtain the current Safety Data Sheet at www.stollerusa.com.

STORAGE AND DISPOSAL

We recommend storing pails a maximum of three pails high. Do not stack pallets. Store container tightly closed in a cool/well-ventilated dry place protected from sunlight above 40°F. Do not allow container to freeze. Store locked up. DISPOSE of contents/container at a permitted facility or as advised by your local regulatory authority. Do not contaminate water, food or feed by storage or disposal. Do not apply directly to areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment wash water. Do not reuse empty container.

CAUTION: The application of fertilizing materials containing molybdenum may result in forage crops containing levels of molybdenum which are toxic to ruminant animals.

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.aapfco.org/metals.html>

Always read and follow label instructions.

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 800-539-5283 or 713-461-1493 | © 2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.

GENERAL INFORMATION

KEYLATE COMO CLASSIC is a convenient plant available source of the essential nutrients cobalt and molybdenum. **KEYLATE COMO CLASSIC** is formulated to be applied as a seed treatment, applied along with liquid fertilizer solutions, impregnated onto dry fertilizer, or applied with pesticide spray solutions and is non-phytotoxic when used as directed.

DIRECTIONS FOR USE

It may be a violation of federal law to use this product inconsistent with its labeling.

KEYLATE COMO CLASSIC can be applied as a seed treatment, in-furrow, transplant water or drench, as a side-dress or foliar application, and through flood, drip sprinkler, overhead, pivot and micro irrigation systems. Apply the recommended amount in a volume sufficient to provide uniform coverage. **KEYLATE COMO CLASSIC** is safe to apply as a seed treatment with inoculant at planting.

RECOMMENDED USES

FOR SEED TREATMENT APPLICATION:

Apply 1-4 oz/cwt of seed. **KEYLATE COMO CLASSIC** is safe to apply with seed treatment inoculants at recommended rates. *Do not use treated seed for food, feed, or oil purposes. Label treated seed in accordance with the requirements of the Federal Seed Act and applicable state and local laws. Use an approved dye to distinguish treated seed and prevent inadvertent use for food, feed or oil purposes.

FOR IN-FURROW APPLICATION:

Apply in-furrow or banded 2x2 alongside the furrow, at 4-8 ounces either with or without starter fertilizer solutions.

FOR FERTILIZER IMPREGNATION:

Apply 2-6 ounces per 100 pounds of fertilizer. Incorporate into liquid fertilizer solutions or impregnation onto dry fertilizer are both acceptable methods of application.

FOR SOIL APPLICATION:

Apply 4 -16 ounces per acre via fertigation through all fertigation methods, drip, flood, pivot, or sprinkler irrigation. Repeat every 7-21 days as needed.

FOR FOLIAR APPLICATION:

Apply 4 to 16 oz per acre for field and vegetable crops, and for fruit, and nut crops, apply 8 to 16 oz. per acre, in a volume sufficient to provide uniform coverage. Repeat every 7-21 days as needed.

Use **KEYLATE COMO CLASSIC** at the above-recommended rates on the following crops:

FIELD CROPS: Alfalfa, bean crops, canola, clover, corn, cotton, flax, forage grasses, peanut, potato, rape (canola), rice, sorghum, soybean, sugar beet, sugarcane, sunflower, tobacco, and wheat, barley, oat, or other cereal grain.

VEGETABLE CROPS: Artichoke, asparagus, broccoli, cauliflower, cabbage, kale, and other cruciferous vegetables; leaf lettuce, head lettuce, dandelion, radicchio, and other leafy vegetables; cucumber, melon, pumpkin, squash, and other cucurbits, onion, garlic, and other bulb vegetables; tomato, pepper, eggplant and other fruiting vegetables; strawberry; carrot, sweet potato, radish, beet and other root and tuber vegetables.

FRUIT AND NUT CROPS: Apple, avocado, blueberry and other bush berries, blackberry, raspberry and other cane berries, sweet orange, lemon, mandarin and other citrus, date palm, fig, grapes, hops, nectarines, oil palm, olive, persimmon, cherry, peach, plums, prune, apricot and other stone fruit, almond, pecan, pistachio, walnut and other tree nuts, pear, pomegranate, and subtropical fruit.

MIXING INSTRUCTIONS: Use the following mixing order: (1) Water (2) **KEYLATE COMO CLASSIC** (3) Pesticide/Fertilizer. **KEYLATE COMO CLASSIC** will disperse in water with little agitation. **KEYLATE COMO CLASSIC** is compatible with most fertilizers and pesticides. Always conduct a jar test when using new or untried combinations.

Notice: The information contained herein is true and accurate to the best of our knowledge and belief, but is presented without guarantee since field conditions and use are beyond our control. Neither the manufacturer nor the seller makes any warranty, expressed or implied concerning the use of this product other than indicated on the label. Buyer assumes all risk of use of this material when such use is contrary to label instruction. Read and follow the label directions carefully.

CONDITIONS OF SALE: 1. Seller warrants that this product consists of the ingredients specified and is reasonably fit for the purpose stated on this label when used in accordance with directions under normal conditions of use. No one, other than an officer of Seller, is authorized to make any warranty, guarantee of direction concerning this product. 2. Because of time, place, rate of application and other conditions of use are beyond seller's control, Seller's liability from handling, storage and use of this product is limited to replacement of product or refund of purchase price.

Always read and follow label instructions.

Manufactured and Guaranteed by Stoller Enterprises, Inc.

9090 Katy Freeway, Suite 400 Houston, Texas 77024 | www.stollerusa.com | 800-539-5283 or 713-461-1493 | © 2021 Stoller USA Inc. All rights reserved. Printed in the U.S.A.