

3-10-1 GUARANTEED ANALYSIS

Total Nitrogen (N)	3.0%
3.0%	
Available Phosphate (P2O5)	10.0%
Soluble Potash (K,O)	1.0%

Derived from anhydrous ammonia, phosphoric acid and potassium hydroxide



WARNING. Causes eye installon. May be harmful in contact with skin. Harmful if swallowed. Avoid breating dust if limit pags mist vipous spray. Avoid relates to the environment. Wear protective probability dust if limit pags mist vipous spray. Avoid relates to the environment. Wear protective water for 515 p.20 minutes. Remove contact leness, if present and continue reining eyes. IF ON SORV. Wash with plenty of water for 155 p.20 minutes. Remove contact leness, if present and continue reining eyes. IF ON SORV. Wash with plenty of water for 155 p.20 minutes. If medical advice is needed, have product container or label available. Now not of reached forbidition. Read label plenty in the contribution of the product of the product of the contribution.

	Net Contents 5 gallons (18.9 liters), 50.5 lbs
	10.11 lbs. per gallon (1212 grams/liter) @ 68° l

Net Contents 265 gallons (1003.1 liters), 2,679 lbs 10.11 lbs. per gallon (1212 grams/liter) @ 68° F

DIRECTIONS FOR USE

Kwik Start LS® may be used as a pop-up, in-furrow at 4 to 8 gallons per acre. Kwik Start LS® can be applied to most vegetable and courbit crops, field crops, row crops, deciduous fruit and nut frees, grapes, berries, olives, avocados and citrus. Mixing with water for field storage is generally not recommend ed. Products blended with water should be used immediately and not stored for prolonged periods. Kwik Start LS® is non-phytotoxic when used as directed and is compatible with most commonly used pesticides and fertilizers.

SOIL APPLICATION RATES

Kwik Start LS® may be applied pre-plant, as a pop-up in furrow, as a starter, side dress, fertigated, top dressed, broadcast or banded to the soil. To apply alone, may be diluted with water to ensure uniform distribution.

ROW CRÓPS, FIELD CROPS & VEGETABLE CROPS: Kwik Start LS® may be used as a pop-up, in-furrow at 4 to 8 gallons per acre.

ROW CROPS, FIELD CROPS & VEGETABLES: Apply 5 to 30 gallons per acre in a surface or sub-surface band. Repeat applications are beneficial FRUIT CROPS, ORCHARDS & VINEYARDS: Inject 5 to 30 gallons per acre 3

to 5 inches deep at the drip line, or surface band and water into the soil. Repeat applications are beneficial.

TRANSPLANT WATER - FIELD & ROW CROPS, VEGETABLE & FRUIT CROPS, ORCHARDS & VINEYARDS: Apply 5 to 8 gallons per acer in a minimum of 100 gallons of solution, not to exceed a 3% Kwik Start LS® to water

FERTIGATION APPLICATION RATES

FIELD & ROW CROPS, ORCHARDS & VINEYARDS, VEGETABLE & FRUIT CROPS: Use 5 to 20 gallons per acre per irrigation. Continue irrigation for a minimum of 1 hour after application to maximize fertilizer distribution. Repeat applications are beneficial.

APPLICATION THROUGH DRIP IRRIGATION - inject after irrigation filter.

FOLIAR APPLICATION RATES

solution concentration.

FIELD & ROW CROPS, VEGETABLE & FRUIT CROPS, ORCHARDS & VINE-YARDS: Apply 1/2 to 2 gallons per acre in enough water for thorough coverage. The maximum recommended concentration of Actagro Kwik Start LS® to water is 4.5.

IS 1.3. VEGETABLES AND LEAFY CROPS: Apply 1/2 to 2 gallons per acre in enough water for thorough coverage. The maximum recommended concentration of Actagro Kwik Start LS® to water is 1:5. Repeat applications are beneficial.

GENERAL INFORMATION

Kwik Start® LS will be influenced by various soil and environmental factors and should be applied under conditions of optimum plant growth. Optimum conditions are defined as adequate humidity, moderate temperatures and adequate soil moisture. Optimum conditions for most crops generally occur during morning and late evening hours.

CAUTION: If any other chemicals are to be used in the spray mixture, be sure to check compatibility. Kwik Start® LS is not compatible with inorganic salts of heavy metals. Do not mix Kwik Start® LS with conventional calcium based fartificate.

CAUTION: Foliar application of Kwik Start® LS close to harvest may leave a brown residue on the harvestable portion of the crop. Rainfall, sprinkler irrigation or high humidity should alleviate this condition.

WARRANTY: Since weather, crops, soil and other conditions may vary, neither Actagor®, LLC, nor the Seller make a warranty of any kind, expressed or implied, concerning the use of this product. The user assumes all risk of use and handling whether or not in accordance with directions or suggestions.

Information regarding the contents and levels of metals in this product is available on the internet at http://www.aapfco.org/metals.htm

Not all Actagro® products are compatible with one another or when mixed with other fertilizers and/or pesticides on the market. A small scale compatibility test is recommended prior to mixing.

MIXING INSTRUCTIONS

- 1. Partially fill tank with water and start agitation.
- 2. Add pesticides or other nutrients.
- 3. Add Kwik Start® LS to circulating tank mix.
- 4. Fill tank and agitate thoroughly.

