

MAX-IN™ Sulphur 13%

By WinField United

0-0-19

GUARANTEED MINIMUM ANALYSIS

Soluble Potash (K₂O).....19.0%
Sulphur (S).....13.0%

STORE ABOVE 4°C

KEEP OUT OF REACH OF CHILDREN



WARNING

Causes serious eye irritation.
Causes skin irritation.

PREVENTION: Wash hands thoroughly after handling. Wear protective gloves/eye protection/face protection.

FIRST AID: IF IN EYES: Rinse cautiously with water for 15 minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical attention. **IF ON SKIN:** Wash with plenty of water. If skin irritation occurs: Get medical attention. Take off contaminated clothing and wash before reuse.

STORAGE: This product can be corrosive to tanks, pumps, meters and fittings. Store in stainless steel or polyethylene tanks with stainless steel or polyethylene fittings.

DISPOSAL: Dispose of contents/container in accordance with Federal, provincial and territorial regulations.

For Chemical Spill, Leak, Fire, Exposure or Accident Call, CHEMTREC 1-800-424-9300

Manufactured for: Winfield Solutions, LLC, P.O. Box 64589, St. Paul, MN 55164 ~ USA

Imported by: United Suppliers Canada Inc., 101-302 Wellman Lane, Saskatoon, Saskatchewan S7T 0J1 ~ CAN

Telephone: 1-888-975-4769

Net Volume: / Volume net :

Lot No.: _____

2/01183

GENERAL INFORMATION: **MAX-IN SULPHUR** contains technology which assists in absorption and movement of nutrients in the plant. **MAX-IN SULPHUR** is not to be used as a replacement for any surfactant in recommendations tank-mixed with herbicides. The use of ammonium sulfate is always recommended when tank-mixed with glyphosate based products and other herbicides.

Check compatibility with orthophosphates, high potash suspensions and chemical mixtures.

MAX-IN SULPHUR can be used in conjunction with glyphosate products. A jar test for compatibility should be conducted prior to large scale mixing. When using **MAX-IN SULPHUR** with pesticides read and follow the pesticide label instructions. Do not mix **MAX-IN SULPHUR** where prohibited by the pesticide label. Mixing **MAX-IN SULPHUR** with certain pesticides or adjuvants, especially oil based products, can increase the risk of phytotoxicity to sensitive crops in conditions that may increase the activity of the spray solution. Do not mix **MAX-IN SULPHUR** with pesticides unless potential phytotoxicity can be tolerated. If uncertain, spray a small test area with the spray mixture before making large scale applications.

MIXING INSTRUCTIONS:

1. Add water and agitate.
2. Add an ammonium sulfate source or water conditioner.
3. Add **MAX-IN SULPHUR**.
4. Add Pesticides.

NOTE: If wettable powders or water dispersible granules (WDG) are used, slurry them in water prior to addition.

MIXING INSTRUCTIONS WHEN MIXED WITH GLYPHOSATE:

1. Fill tank ½ full with water and agitate.
2. Add ammonium sulfate source or water conditioner.
3. Add **MAX-IN SULPHUR**, continue agitation.
4. Fill tank to ¾ full and continue agitation.
5. Add glyphosate and continue agitation.
6. Fill tank to full.

Do not mix **MAX-IN SULPHUR** and glyphosate together in the cone inductor. If using a cone induction system, rinse it out thoroughly with water between the addition of **MAX-IN SULPHUR** and glyphosate. At the end of the day, rinse out spray booms.

DIRECTIONS FOR USE: This product must only be used for corrections of deficiencies as determined by foliar and/or soil testing. For specific recommendations for crops in your area, consult an agricultural representative or a professional agricultural consultant. **CAUTION:** Plant and leaf injury may occur on some crops when certain weather and growing conditions are present. Do not apply to foliage of any crop when temperatures are above 32°C.

A number of variables influence the rate of application necessary to prevent or correct a specific deficiency. For best results, follow good agronomic practices and tissue test.

Foliar Applications:

Crop	Rate (L/ha)	Application Timing:
Alfalfa	5	Apply after each cutting when sufficient re-growth has occurred.
Cereal Crops		Apply between tillering and early joint stages. Later application may be needed; tissue samples should be used to determine later usages.
Corn		Apply when corn is in the V4 to V7 stage. Later application may be needed; tissue samples should be used to determine later usages.
Other Crops	Apply according to directions of your regional Winfield Plant Nutrition Specialist or local agricultural professional	
<ul style="list-style-type: none"> • Use in a minimum of 100 L/ha of water. 		

WARRANTY DISCLAIMER

Winfield Solutions, LLC warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, NOR IS ANY REPRESENTATIVE OF SELLER AUTHORIZED TO MAKE ANY SUCH WARRANTY OR MODIFY THESE TERMS. This warranty does not extend to the storage, handling or use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such storage, handling or use. Seller shall not be responsible for incidental or consequential damages, if any, resulting from a breach of warranty.