Caution: This product contains BORON. Determine plant sensitivity prior to use. Crops sensitive to this elements may be injured.



MICRO MN + B

GUARANTEED ANALYSIS

Boron (B)	2.00%
Manganese (Mn)	
10.00% Chelated Manganese (Mn)	
Derived From: Carboxylic Acid, Manganese EDTA, Boric Acid.	
Chelating agent is EDTA (ethylenediaminetetraacetic acid).	
Chloride (CI) maximum.	0.12%



Suspected of damage to fertility or to the unborn child. Wash face, hands, and any exposed skin after handling. Do not eat, drink, or smoke when using this product. If swallowed: Call a Poison Center if you feel unwell. Rinse mouth.

GENERAL INFORMATION

MICRO MN + B is formulated to be compatible with Glyphosate and other selective and non-selective herbicides. Research conducted with tank mixes of the chelating agents used in MICRO MN + B helped to maintain the efficacy of the herbicides. This product gives best results when crops have adequate supplies of water and major nutrients, and are not under stress for other reasons. Conditions which are responsible for one particular deficiency can also induce deficiencies of other trace elements.

MICRO MN + B is particularly suitable for foliar application and for general soil applications. Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops. Plants need only very small amounts of micronutrients for maximum growth.

KEEP OUT OF REACH OF CHILDREN

MANUFACTURED BY
MILLER CHEMICAL & FERTILIZER, LLC
P.O. Box 333
Hanover, PA 17331
1-800-233-2040

Net Weight 15 lb. (6.795 kg)

CROPS SUSCEPTIBLE TO NUTRITIONAL MICRONUTRIENT DEFICIENCY

MICRO MN + B is recommended for crops such as alfalfa, almonds, avocados, apples, broccoli, cabbage, canola, cauliflower, carrots, celery, citrus, corn, cotton, grapes, lettuce, milo, melons, nectarines, rice, pears, peaches, pecans, peppers, plums, prunes, potatoes, peanuts, sorghum, soybeans, sugar beets, sweet corn, sugar cane, strawberries, tomatoes, turnips, walnuts, watermelons, wheat and most other crops.

FIRST AID

EYE CONTACT: Immediately flush eyes with plenty of water for at least 15 minutes, while holding eyelids apart to ensure flushing of entire surface. Call a physician. SKIN CONTACT: Immediately flush skin with plenty of water for at least 15 minutes, while removing contaminated clothing and shoes. Thoroughly clean clothing and shoes before reuse. Call a physician.

INHALATION: Remove to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. If breathing is difficult give oxygen. Call a physician. **INGESTION:** If swallowed, DO NOT induce vomiting. Rinse mouth with water. Dilute stomach contents by drinking water. If vomiting occurs spontaneously, keep head below hips to prevent breathing vomit into lungs. NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. Call a physician immediately!

MIXING

MICRO MN + B is soluble in water. When mixing; do NOT pre-mix a small amount of water before adding to the spray tank. MICRO MN + B should be added slowly to the majority of the spray water volume while it is being agitated. Agitation should continue for a minute or two after the addition of MICRO MN + B. If pesticides are to be used in the spray mix, add them after the MICRO MN + B has dissolved.

RECOMMENDED RATES

Soil: Apply 1 lb. per acre of MICRO MN + B in combination with starter or pop-up fertilizers. MICRO MN + B can also be applied at 1 lb. per acre broadcast with pre-plant or pre-emergence herbicide application.

Foliar: Apply 1 lb. per acre of MICRO MN + B in a minimum of 20 gallons of water per acre. Applications made in lower volumes of water, especially if the crop is under stress, may cause injury. Therefore, use no more than ½ lb of MICRO MN + B per 5 gallons of water used per acre.

Aerial application: Apply at the same use rate of 1/4 lb. per 5 gallons of water used per acre. **DO NOT** exceed 1 lb. per acre per application even if exceeding 20 gallons of water per acre.

CAUTION: Do not apply more than 1 lb. per acre of **MICRO MN + B** per foliar application as crop injury may occur. Multiple applications should be made at no less than 10 to 14 day intervals to reduce the potential for crop injury. Wash hands with soap and water before meals and after work. Store in the original container tightly closed, in a dry place. Store away from children, pets, livestock and foodstuffs.

Use this product in accordance with good agronomic practices, which include utilizing proven spray equipment set for proper coverage. Do not make applications when temperatures are too hot, as crop damage may occur. Applications should be made at temperature levels and when other environmental conditions in your area are such that your experience indicates the application will be compatible and will accomplish the desired result.

Notice of Warranty: Miller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Miller. In no case shall Miller be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the buyer. MILLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Information concerning the raw materials composing this product can be obtained by writing to Miller Chemical & Fertilizer, LLC, P.O. Box 333, Hanover, PA 17331. Please refer to the batch number found on this container.