

NIMAXXA™

Suspension Concentrate Seed Treatment Use

A suspension concentrate biological nematicide for seed treatment use for protection against labeled nematodes.

Active Ingredients:

	By Wt.
<i>Bacillus subtilis</i> strain CH4000*	3.33%
<i>Bacillus paralicheniformis</i> strain CH0273*	3.33%
<i>Bacillus paralicheniformis</i> strain CH2970*	3.33%
Other Ingredients:	<u>90.01%</u>
Total:	100.00%

*Contains a minimum of 1.67×10^{10} colony forming units (CFU) of each active ingredient per mL of product.

EPA Reg. No. 2375-5-70506

EPA Est. No. 2375-BRA-1

KEEP OUT OF REACH OF CHILDREN

See other panels for additional precautionary information.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY: 1-866-673-6671.

FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire, exposure or accident) CALL CHEMTREC: 1-800-424-9300 or +1-703-527-3887.

Net Contents: 2.5 Gallons

Not for sale or use after:

Batch/Lot code:

Sold By: **UPL NA Inc.**
PO Box 12219, Research Triangle Park
NC 27709 U.S.A. • 1-800-438-6071
BR LBL-061 10/24



PRECAUTIONARY STATEMENTS

Personal Protective Equipment (PPE)

Applicators and other handlers must wear:

- protective eyewear
- long-sleeved shirt
- long pants
- shoes plus socks
- waterproof gloves

Mixers/loaders and applicators must wear a minimum of a NIOSH-approved particulate filtering facepiece respirator with any N, R, or P filter; OR a NIOSH-approved elastomeric particulate respirator with any N, R, or P filter; OR a NIOSH-approved powered air-purifying respirator with an HE filter. Repeated exposures to high concentrations of microbial proteins can cause allergic sensitization.

User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

Engineering Controls

When handlers use closed systems, or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607 (d) and (e)], the handler PPE requirements may be reduced or modified as specified in the WPS.

IMPORTANT: When reduced PPE is worn because a closed system is being used, handlers must be provided all PPE specified above for "applicators and other handlers" and have such PPE immediately available for use in an emergency, such as a spill or equipment break-down.

User Safety Recommendations

Users should:

- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove clothing/PPE immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

For container sizes <5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

For container sizes ≥5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. Do not discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), restricted-entry interval, and notification to workers. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.

Exception: If the product is soil-injected or soil incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water) is:

- Protective eyewear
- Coveralls
- Waterproof gloves
- Shoes plus socks

Non-Agricultural Use Requirements

The requirements in this box apply to uses that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Keep unprotected persons out of treatment area until seeds have dried or been packaged.

Product Information

NIMAXXA™ is a bionematicide seed treatment suspension concentrate formulation that provides suppression and control of labeled nematodes in a variety of horticultural and broadacre crops for any and all seed practices including farm, greenhouse, nursery, or any other seed development and application processes. Seed must be sound and well-cured before treatment.

Seed Treatment Equipment

NIMAXXA™ may be used both for commercial and for on-farm application. NIMAXXA™ can be applied with mechanical, slurry, or mist-type seed treating equipment, as long as the equipment can be calibrated to accurately and uniformly apply the product to seed without undue mechanical damage to the seed. Uniform application to seed is important for all seed treatment products.

Application Instructions

Seed must be sound and well-cured before treatment. Refer to the **COMPATIBILITY AND TANK MIXING INSTRUCTIONS** below and follow the crop-specific application rates below.

COMPATIBILITY AND TANK MIXING INSTRUCTIONS

COMPATIBILITY

NIMAXXA™ is physically compatible with many commonly used fungicides, herbicides, insecticides, and/or nutrient products. However, the compatibility of NIMAXXA™ with all potential tank-mix partners has not been fully investigated. If tank mixing with other products is desirable, conduct a jar test with the volumes and rates typically used in labeled applications. Using a small container of water, add the proportionate amounts of the products in the following order: wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 15 minutes. Look for signs of separation, globules, sludge, flakes, or other precipitates. Physical compatibility is indicated if the combination remains mixed or can be remixed readily.

TANK MIXING INSTRUCTIONS

When tank-mixing NIMAXXA™ with other registered pesticides and/or other products, always read and follow all use directions, restrictions, and precautions of both NIMAXXA™ and the tank-mix partner(s). It is the pesticide user's responsibility to ensure that all tank-mix products are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture. Use of the resulting tank mix must be in accordance with the more restrictive label limitations and precautions. DO NOT exceed label dosage rates.

All applications of NIMAXXA™ involve the following general tank mixing procedures. Begin with clean supply, mixing and application equipment. Add one-half of the required amount of non-chlorinated water to the supply/mixing tank and start agitation. Dissolve the needed amount of NIMAXXA™ into a small volume of water to make a pre-mix slurry. Add needed amounts of products to the mix tank in the following order: water conditioners, water soluble packages (wait for them to completely dissolve), wettable powders and water-dispersible granular products, followed by liquids, including, liquid slurries (add NIMAXXA™ pre-mix slurry here), flowables, suspension concentrates, emulsifiable concentrates, with surfactants and/or adjuvants added last. After adding the required quantity of NIMAXXA™ and other products to the mix tank, complete filling with water to the required final volume and maintain agitation throughout

application. DO NOT allow tank mixture to remain in the tank overnight. DO NOT allow tank mixture to sit for long periods during the day without agitation. If non-chlorinated water is not available, DO NOT let NIMAXXA™ remain in chlorinated water for long periods of time as it may have detrimental effect on the bacterial endospores.

The appropriate volume of slurry for seed treatment depends on crop, temperature, humidity, type of equipment, and other factors and must be adjusted per normal treating practices for the circumstances. Contact your local supplier or distributor representative for additional specifications for slurry volumes and questions on any tank mixes or restrictions.

Recalibrate equipment to compensate for the required slurry rate to ensure all products are applied at the correct application rate.

NOTE: This product does not contain dye and is not covered by an appropriate tolerance, tolerance exemption, or other clearance under the Federal Food, Drug and Cosmetic Act. To comply with 40 CFR 153.155, therefore, all seed treated commercially with this product must be colored with an EPA-approved dye or colorant of a suitable color to prevent accidental use as food for man or feed for animals.

SEED BAG LABEL REQUIREMENTS

The Federal Seed Act requires that bags containing seeds treated with this product shall be labeled with the following information:

- This seed has been treated with *Bacillus subtilis* strain CH4000 and *Bacillus paralicheniformis* strains CH0273 and *Bacillus paralicheniformis* strain CH2970.
- Do not use for feed, food or oil purposes. Store away from feed and food stuffs.

User is responsible for ensuring that the seed bag meets all requirements under the Federal Seed Act.

Do not use NIMAXXA™ in combination with other seed treatment products unless compatibility has been verified. Read and carefully follow all label directions of each product used in the treating mixture. When using combinations of products, the most restrictive of label limitations and precautions must be followed.

Application Rates, Crops, and Pests

Application Rates

NIMAXXA™ may be applied as a seed treatment following the product rates described below according to the specific crop directions. Use the lower application rate for light infestations and the higher labeled rate for heavy infestations.

Specific Crop Directions

Crop Rotation Restrictions

There are no crop rotation restrictions. Any crop can be rotated at any time after application of NIMAXXA™.

Crop Group 1 -	Root and Tuber Vegetables Group: Artichoke
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 5 -	Brassica (Cole) Leafy Vegetables Group: Broccoli, Cabbage, Cauliflower, Brussels Sprouts, Collards, Kale, Mustard Greens, Kohlrabi, and other brassica crops
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 15 -	Cereal Grains Group: (Exclude Corn) Barley, Buckwheat, Millet, Oat, Pearl Millet, Proso Millet, Rice**, Rye, Sorghum, Teosinte, Triticale, Wheat, and other cereal grain crops
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

**Do not apply to flooded fields.

Crop Group 15 -	Cereal Grains Group: Corn (Field corn, Popcorn, Sweet corn, and Corn Grown for Seed)
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 20 -	Oilseed Group: Cottonseed
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Rotylenchulus reniformis*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 9 -	Cucurbit Vegetables Group: Chayote (fruit), Chinese wax gourd (Chinese preserving melon), Citron melon, Cucumber, Cantaloupe, Edible Gourd (such as Chinese Okra, Cucuzza, Hechima, and Hyotan), Gherkin, Melon, <i>Momordica</i> spp. (such as Balsam apple, Balsam pear, Bitter melon, and Cucumber), Muskmelon, Pumpkin, Summer and Winter Squash (such as Acorn, Butternut, Calabaza, Crookneck, Hubbard, Scallop, Spaghetti, and Zucchini), Squash, Watermelon, Chinese Watermelon, and other cucurbit crops
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 8 -	Fruiting Vegetables Group: African eggplant, Bell pepper, Bush tomato, Cocona, Currant tomato, Eggplant, Garden huckleberry, Goji berry, Groundberry, Martynia, Naranjilla, Nonbell pepper, Okra, Pea eggplant, Pepino, Roselle, Scarlet eggplant, Sunberry, Tomatillo, and Tomato
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i> <i>Globodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 4 -	Leafy Vegetables (Except <i>Brassica</i> Vegetables) Group: Lettuce, Celery, Spinach, Parsley, Radicchio, and other leafy vegetables crops including those grown for seed production
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 6 -	Legume Vegetables (Succulent or Dried): (Exclude Lentils and Soybeans) Beans, Snap beans, Garbanzo beans, Lima beans, Peas, Chick peas, Dried bean and pea (<i>Lupinus</i> spp., <i>Phaseolus</i> spp., <i>Vigna</i> spp., and <i>Pisum</i> spp.), and other legume vegetable crops including those grown for seed production
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Lentils
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 18 -	Nongrass Animal Feeds (Forage, Fodder, Straw, and Hay) Group: Alfalfa, Clover, and Forages
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i> <i>Globodera*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 20 -	Oilseed Group: (Excludes Cottonseed, Peanut, and Soybeans) Canola, Castor, Flax, Palm, Olive, Rapeseed, Safflower, Sesame, Sunflower, and other oil seed crops
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop -	Peanut
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Belonolaimus*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 1 -	Root and Tuber Vegetables Group: Potato
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i> <i>Globodera*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 1 -	Root and Tuber Vegetables Group: (Exclude Potato, Sugar Beet, and Artichoke) Arracacha, Arrowroot, Bean and True Yam, Beets, Bitter and Sweet Cassava, Black and Spanish Salsify, Carrot, Celeriac (Celery Root), Chayote (Root), Chicory, Chufa, Dasheen (Taro), Edible Burdock, Edible Cana, Ginger, Ginseng, Green Beet, Horseradish, Leren, Oriental Radish, Parsnip, Radish, Rutabaga, Salsify (Oyster plant), Skirret, Sweet Potato, Tanier (Cocoyam), Turmeric, Turnip, Turnip-rooted Chervil, and Turnip-rooted Parsley, and other root and tuber vegetable crops
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i> <i>Globodera*</i>	0.5 ml/kg to 5 ml/kg or 0.77 fl oz/100 lbs to 7.67 fl oz/100 lbs

*Not registered for use in California.

Crop Group 6 -	Legume Vegetables (Succulent or Dried): Soybeans
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i> <i>Rotylenchulus*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

Crop Group 1 -	Root and Tuber Vegetables Group: Sugarbeet
Target Nematode	Application Rate: Product/Seed Wt:
<i>Meloidogyne*</i> <i>Pratylenchus*</i> <i>Heterodera*</i>	0.25 ml/kg to 3 ml/kg or 0.38 fl oz/100 lbs to 4.60 fl oz/100 lbs

*Not registered for use in California.

USE RESTRICTIONS

Do not use treated seed for food, feed or oil purposes. For at-planting seed treatments, treat only those seeds needed for immediate use, minimizing the interval between treatment and planting. Care must be exercised in the handling of treated seed. Augers used for handling treated seed must not be used to move seed for feed, food or oil processing. Do not re-use bags from treated seed to handle food or feed products.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place under low humidity. Avoid direct sunlight and avoid excess heat. Keep out of reach of children and animals. Store in original container only. Carefully open container. After partial use, replace lid of container and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and Spills) at 1-800-424-9300. To confine spill, dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. If a leaky container must be contained within another, mark the outer container to identify the contents.

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Plastic Containers (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume, and recap. Shake for 30 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

Nonrefillable Plastic Containers (Containers greater than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 60 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

IMPORTANT INFORMATION READ BEFORE USING PRODUCT

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

NOTICE: Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.** UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

UPL, the UPL logo and NIMAXXA are trademarks of a UPL Corporation Limited Group Company.

©2024 UPL Corporation Limited Group Company

2911455-A (1124)

ESL092724-12001-111924

NIMAXXA™

Suspension Concentrate Seed Treatment Use

A suspension concentrate biological nematocide for seed treatment use for protection against labeled nematodes.

Active Ingredients:	By Wt.
<i>Bacillus subtilis</i> strain CH4000*	3.33%
<i>Bacillus paralicheniformis</i> strain CH0273*	3.33%
<i>Bacillus paralicheniformis</i> strain CH2970*	3.33%
Other Ingredients:	90.01%
Total:	100.00%

*Contains a minimum of 1.67×10^{10} colony forming units (CFU) of each active ingredient per mL of product.

EPA Reg. No. 2375-5-70506
EPA Est. No. 2375-BRA-1

KEEP OUT OF REACH OF CHILDREN

See other panels for additional precautionary information.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**FOR 24-HOUR MEDICAL EMERGENCY ASSISTANCE
CALL ROCKY MOUNTAIN POISON AND DRUG SAFETY:**
1-866-673-6671.

**FOR 24-HOUR CHEMICAL EMERGENCY (Spill, leaks, fire,
exposure or accident) CALL CHEMTREC:** 1-800-424-9300
or +1-703-527-3887.

PRECAUTIONARY STATEMENTS

Environmental Hazards

For container sizes <5 gallons

For terrestrial uses: Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read entire label before using this product.

Net Contents: 2.5 Gallons

Sold By: **UPL NA Inc.** • PO Box 12219, Research Triangle Park
NC 27709 U.S.A. • 1-800-438-6071

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage

Store in a cool, dry place under low humidity. Avoid direct sunlight and avoid excess heat. Keep out of reach of children and animals. Store in original container only. Carefully open container. After partial use, replace lid of container and close tightly. Do not put concentrate or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by storage or disposal. In case of spill, avoid contact, isolate area, and keep out animals and unprotected persons. Confine spills. Call CHEMTREC (Transportation and Spills) at 1-800-424-9300. To confine spill, dike surrounding area or absorb with sand, cat litter or commercial clay. Place damaged package in a holding container. If a leaky container must be contained within another, mark the outer container to identify the contents.

Pesticide Disposal

To avoid wastes, use all material in this container by application according to label directions. If wastes cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

Container Handling

Nonrefillable Plastic Containers (Containers equal to or less than 5 gallons): Nonrefillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank, and drain for 10 seconds after the flow begins to drip. Add water – at least 2% of the container volume, and up to 1/3 of the volume of water needed to make the proper slurry composition with a maximum of 1/4 of the container volume, and recap. Shake for 30 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. If used in application equipment, adjust the slurry volume application rate to account for any added rinsate water. Then offer container for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill or by incineration. If burned, stay out of smoke.

