

NIXELA™

0-0-5

GUARANTEED ANALYSIS:

Soluble Potash (K ₂ O)	5.00 %
Calcium (Ca)	2.64 %
Magnesium (Mg)	0.84 %
Boron (B)	0.24 %

Derived from: Potassium Hydroxide, Calcium Hydroxide, Magnesium Oxide, Boric Acid

NIXELA™ is a pending trademark of AECI Ltd.

WARNING – Product contains Boron. The use of the fertilizing material on any crops other than those recommended may result in serious injury to the crop(s).

KEEP OUT OF REACH OF CHILDREN



WARNING

Causes serious eye irritation. Wear protective gloves. Wear eye or face protection. Use only outdoors or in a well-ventilated area. Avoid breathing vapor. Wash hands thoroughly after handling.

FIRST AID:

If in eye: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If on skin: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice. Have a person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

DIRECTIONS FOR USE: USE ONLY AS INDICATED - INCLUDED BUT NOT LIMITED TO THE CROPS LISTED BELOW

BEFORE USING THIS PRODUCT READ THE LABEL CAREFULLY

NIXELA should not be added with any sulfur or phosphite based products. **Do not apply less than 25 fluid ounces per acre of NIXELA for any given crops.**

APPLICATION TABLE:

CROP	APPLICATION RATE	APPLICATION METHOD
APPLES, PEARS	25 to 40 fluid oz/100 gallons of water to a max of 40 fluid oz/acre	Apply 3-4 applications from 50% flowering and repeat with 21-day intervals. For use on Pears: Do not combine NIXELA with other foliar feeds or adjuvants.
CEREALS (BARLEY, OATS, WHEAT, TEFF)	25 to 30 fluid oz/100 gallons of water to a max of 30 fluid oz/acre	Apply at 4 and 8 weeks after emergence, or prior to expected heat waves or cold spells.

CITRUS	25 to 40 fluid oz/100 gallons of water to a max of 40 fluid oz/acre	Apply 3 to 4 applications from 100% petal drop and repeat with 28-day intervals.
CORN	25 to 30 fluid oz/100 gallons of water to a max of 30 fluid oz/acre	Apply at 4 and 8 weeks after emergence, or prior to expected heat waves or cold spells.
CUCURBITS (PUMPKINS, WATERMELONS, MELONS AND CUCUMBERS)	48 to 65 fluid oz/100 gallons of water to a max of 65 fluid oz/acre	Apply at flowering and repeat with 21-day intervals.
HEMP	25 to 40 fluid oz/100 gallons of water to a max of 40 fluid oz/acre	Apply 2 applications after flowering with a 10 to 14-day interval period.
LEGUMES (SOYBEAN, ALFALFA, PEANUT)	25 to 30 fluid oz/100 gallons of water to a max of 30 fluid oz/acre	Apply at 4 and 8 weeks after emergence, or prior to expected heat waves or cold spells.
LETTUCE, CAULIFLOWER, CABBAGE, CHINESE CABBAGE	48 to 65 fluid oz/100 gallons of water to a max of 65 fluid oz/acre	Begin applications prior to onset of head formation and repeat at 14 to 21-day intervals.
PEACHES, NECTARINES, PLUMS, APRICOTS, ALMONDS, OLIVES	25 to 40 fluid oz/100 gallons of water to a max of 40 fluid oz/acre	Apply 3 to 4 applications from 75% petal drop and repeat with 21 to 28-day intervals.
POTATOES	48 to 65 fluid oz/100 gallons of water to a max of 65 fluid oz/acre	Apply at tuber initiation and repeat every 21 to 28 days.
TOMATOES, PEPPERS, BEANS, PEAS, ONIONS	48 to 65 fluid oz/100 gallons of water to a max of 65 fluid oz/acre	Begin applications prior to fruit set and repeat at 14 to 21-day intervals.
TOBACCO	48 to 65 fluid oz/100 gallons of water to a max of 65 fluid oz/acre	Apply 2 to 3 applications bi-weekly, starting at 12 inches height.
VINES	25 to 30 fluid oz/100 gallons of water to a max of 30 fluid oz/acre	Apply after flowering every three weeks, and apply before expected heat waves or cold spells.

IMPORTANT: BEFORE APPLYING NIXELA, CONSULT A QUALIFIED AGRONOMIST OR AGRICULTURAL ADVISER ON THE USE THEREOF.

Information regarding the contents and levels of metals in this product is available on the internet at

<http://www.aapfco.org/metals.html>

Net Content: 2.5 Gallons (9.46 liters)

Density: 10.52 lbs/US gal @ 68° F

Net Weight: 26.3 lbs (11.92 kg)

GUARANTEED BY:

Nulandis (A division of AECI® Ltd)
AECI CHEM PARK, BLOCK NO 2 - 2ND FLOOR
200 BERGRIVIER DRIVE, CHLOORKOP
KEMPTON PARK, 1619
SOUTH AFRICA
T: +27 11 823 8000
www.aeciworld.com