

NoMate® Duo

CM/LRX

NoMate® Duo CM/LRX is a mating disruption product for control of codling moth, obliquebanded, pandemis, fruittree, threelined, and european leafrollers.

Active Ingredients:

(E,E)-8,10-Dodecadien-1-ol	6.35%
Z-11-Tetradecen-yl Acetate	6.11%
Other Ingredients.....	87.54%
Total	100.00%

*330 mg active ingredient per dispenser.

CAUTION

KEEP OUT OF REACH OF CHILDREN

FIRST AID

IF ON SKIN OR CLOTHING

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

IF IN EYES

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses if present after the first 5 minutes then continue rinsing.
- Call a poison control center or doctor for treatment advice.

IF INHALED

- Move person to fresh air.
- If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call a poison control center or doctor for further treatment advice.

IF SWALLOWED

- Call the poison control center or doctor immediately for treatment advice.
- Have the person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to do so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact the National Pesticide Information Center (NPIC) at 1-800-858-7378 for emergency treatment.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

Caution: Harmful if absorbed through skin, inhaled, or swallowed. Causes moderate eye irritation. Avoid contact with skin, eyes, and clothing. Avoid breathing vapor. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash before reuse. Wear the appropriate Personal Protective Equipment (PPE).

SCENTRY
BIOLOGICALS INC.

610 Central Avenue • Billings, Montana 59102
(406) 248-5856 • 800-735-5323
www.scentry.com

Lot No.

ENVIRONMENTAL HAZARDS

For terrestrial uses: Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not open pouches until ready to use.

INSTRUCTIONS FOR USE

NoMate® Duo CM/LRX is intended to prevent damage to pome fruit, stone fruit and tree nut crops caused by Codling moth (*Cydia pomonella*), Obliquebanded leafroller (*Choristoneura rosaceana*), Pandemis leafroller (*Pandemis pyrusana*), Fruittree leafroller (*Archips argyrospilus*), Threelined leafroller (*Pandemis limitata*), and European leafroller (*Archips rosanus*). The product functions by disrupting the mating communication between the sexes of the adult moths.

RATE OF APPLICATION

The rate of application is 200-400 dispensers per acre (500-1000 per hectare). Use the spirals in 1-3 applications per year as needed. Do not exceed 150 grams of active ingredient (454 Dispensers) per acre per year.

METHOD OF APPLICATION

NoMate Duo CM/LRX is applied directly by wrapping the spiral around the stems or branches. Open the bag just prior to application. Apply product to the upper third of the canopy. For maximum effectiveness, it may be necessary to increase the density of the dispenser placement around the orchard perimeter.

TIMING OF APPLICATION

Monitor using Scentry Pheromone lures and Traps to determine the time of application. Make the first application within 2 to 4 days after the first male moth is captured in a trap (biofix). Prediction Degree Day Models can provide assistance in determining application timing. Model information and economic thresholds may vary according to geographical regions, consult local extension offices and crop advisors in the area for more information.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed through storage or disposal. Pesticide Storage: Store in a cool place until used. Prolonged storage should be at or less than 40°F (4°C). Pesticide Disposal: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility. Container Disposal: Nonrefillable container. Do not reuse or refill this container. Dispose of empty bag in the trash.

WARRANTY STATEMENT

Scentry Biologicals warrants that this product conforms to the description of this label and is reasonably fit for the purposes stated on this label when used in accordance with the directions on this label under normal conditions of use, to the extent consistent with applicable law. If this product is defective, buyer's exclusive remedy, as permitted by law, shall be the replacement of the product, or if replacement is impracticable refund of the purchase price.

SCENTRY BIOLOGICALS MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

NoMate, Scentry are trademarks of:
Scentry Biologicals Inc.
Scentry Biologicals Inc. 406-248-5856

EPA REG. No. 36638-47
EPA EST. No. 36638-MT-01
Net contents: 300 Spirals