



Parka®

POWERED BY **SureSeal™**
BIOFILM TECHNOLOGY

NET CONTENTS:

2.5 gal

(9.5 Liters)

CROP CUTICLE SUPPLEMENT

**Read the label carefully before using the product.
Use reserved for farmers and professional applicators.**

**IN ORDER TO AVOID RISKS FOR PEOPLE AND THE ENVIRONMENT,
CAREFULLY FOLLOW THE INSTRUCTIONS FOR USE.**

Manufactured By:

cultiva®

Cultiva LLC
4780 West Harmon Ave., Suite 6
Las Vegas, NV 89103
Telephone (888) 638-1955

Parka is a trademark of Cultiva.

U.S. Patent #8,752,328

CROP CUTICLE SUPPLEMENT

Parka® Crop Cuticle Supplement is a patented blend of phospholipids designed to supplement the cuticle of growing fruit and foliage. Supplementing the cuticle of the fruit reduces micro-fractures to minimize cracking from rain, improve finish, reduce post harvest disorders as well as provide sunburn protection. Parka® applications provide support to the cuticle of the foliage and reduce stress to improve fruit quality and yield. Parka is exempt from tolerance and there is no limit on the number of applications per season. *It all goes back to the cuticle.™*

Crop	Rate per Acre (Hectare)
Cherries	1 gal (9.5 L)
Other Stone Fruit	1 gal (9.5 L)
All Berry Crops	2 – 4 qts (4.75 – 9.5 L)
Grapes	0.5 – 1 gal (4.75 – 9.5 L)
Pome Fruit	1 gal (9.5 L)
Nuts	1 gal (9.5 L)
Field Crops	2 – 4 qts (4.75 – 9.5 L)
Citrus, Avocado, Pomegranate	2 – 4 qts (4.75 – 9.5 L)

FIRST AID	
If in Eyes:	Rinse with slowly and gently running water for 15-20 minutes. If wearing contacts remove after five (5) minutes of rinsing and then continue rinsing. If irritation persists get medical attention.
If Swallowed:	Give large quantities of water. DO NOT induce vomiting. Never give anything by mouth to an unconscious person. Get medical attention.
If on Skin:	Remove contaminated clothing. Thoroughly wash exposed skin area(s) with soap and water. If irritation occurs get medical attention. Do not reuse contaminated clothing until washed and cleaned.
If Inhaled:	Remove individual from area. Give symptomatic and supportive care if required. Get medical attention.

Crop	USE GUIDELINES
Cherries	Apply in a 2 application program with the first application at 100% shuck fall and the second application at straw color. If tank mixed with GA, ensure this application occurs 72 hours prior to a rain event.
Other Stone Fruit	Apply in a 3 application program with the first application at fruit set. Reapply at 21–30 day intervals.
All Berry Crops	Apply in a 3-4 application program with the first application when 1st fruit is 5-7 mm in size. Reapply at 14–21 day intervals. Rate: 2-4 qts per acre starting at fruit set of first bloom.
Grapes	Apply 3-4 applications with the first application starting at 10 mm (pea size), a second application at pre-bunch closure, followed by a 3rd application at veraison. A 4th application should be made 10-14 days prior to harvest if increased post harvest quality is desired.
Pome Fruit	Apply in a 3 application program with the first application at fruit set. Reapply at 21–30 day intervals.
Nuts	Apply in a 3 application program with the first application 10 days after nut set. Reapply at 21–30 day intervals.
Field Crops	Onion: Apply when tops fall and bulb is exposed, every 14 days. Tomato, Peppers: Apply at first fruit set; reapply every 14-21 days. Melons: Apply at first fruit set; reapply every 21 days. Leafy Greens: Apply 10-14 days post transplant, reapply every 14-21 days.
Citrus, Avocado, Pomegranate	Apply at fruit set. Reapply every 21-30 days.

For best results, complete coverage on the crop is required. Apply in the morning with low temperatures for optimal drying time. Parka may be applied outside the above recommendations prior to extreme rain or heat events. Additionally, Parka may be applied immediately pre-harvest for improved shelf life.

COMPATIBILITY: PARKA® is compatible with most other crop protection products, provided application coincides with the conditions on each label. Do not tank mix, or overlap Parka® applications within 10 days of applications of or ethyl mercaptan (Captan). If using micronized or dusting sulfur, do not apply Parka® within 3 days of a sulfur application. If using Lime Sulfur wait 7 days before applying Parka®. Do not tank mix with EC based materials. Do not tank mix with any material containing oil. Do not mix with surfactants, stickers or pinolene based materials. Parka® should be the last product added to the tank. For best results, finished spray solution pH should be between 5 and 7.

STORAGE AND HANDLING

Storage: Protect from freezing. Store in a cool, dry, shaded area away from children and animals. Store above 50F.

Spills: Surround and dike area of spill. Prevent runoff into streams, other waterways, or sewer systems. Soak up spill and place in containers. Wash contaminated area with soap and water.

Disposal: Dispose of material and rinsate as directed by local state, or Federal authorities. Keep out of streams and other waterways.

WARRANTY AND DISCLAIMER STATEMENT

To the extent consistent with applicable law, seller makes no warranty, express or implied, of merchantability, fitness or otherwise concerning the use of this product other than as indicated on the label. To the extent consistent with applicable law, user assumes all risks of use, storage or handling not in strict accordance with accompanying directions.

CONDITIONS: Use Directions must be followed carefully. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of factors including, but not limited to: weather conditions; presence of other materials; or the manner of use or application, all of which are beyond the control of the manufacturer or seller. The buyer shall assume all such risks.



Manufacturer: Cultiva LLC
4780 West Harmon Ave., Suite 6
Las Vegas, NV 89103
Telephone (888) 638-1955



www.cultiva.com

Emergency Contact/Number: (888) 638-1955 (US only); (702) 987-6203 (outside US)
Parka® is a trademark of Cultiva.