

[LABEL]

GROUP	2	HERBICIDE
-------	---	-----------

**Precede™**

HERBICIDE

DRY FLOWABLE

For Sale for Use on Wheat (spring and durum), Barley, and Summerfallow in the Prairie Provinces and the Peace River Region of British Columbia

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.: 29483 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: Tribenuron-methyl 75%  
Contains the allergens sulfites and milk

WARNING – EYE IRRITANT  
POTENTIAL SKIN SENSITIZER

NET CONTENTS: 160 grams – 13.6 Kg

FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104  
1-833-362-7722

PRECAUTIONS: WARNING - EYE IRRITANT.  
POTENTIAL SKIN SENSITIZER.  
DO NOT APPLY BY AIR.  
KEEP OUT OF REACH OF CHILDREN.  
MAY BE HARMFUL IF SWALLOWED, INHALED OR  
ABSORBED THROUGH SKIN.  
AVOID BREATHING SPRAY MIST. USE WITH ADEQUATE  
VENTILATION.  
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.  
WEAR SUITABLE CHEMICAL RESISTANT GLOVES AND  
GOGGLES OR FACE SHIELD DURING MIXING AND  
LOADING.  
MAY IRRITATE SKIN AND EYES. WHEN USING, DO NOT  
EAT, DRINK OR SMOKE.  
WASH THOROUGHLY WITH SOAP AND WATER AFTER  
HANDLING. REMOVE CONTAMINATED CLOTHING  
IMMEDIATELY AFTER USE.  
DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS FOR  
12 HOURS FOLLOWING APPLICATION.  
**TOXIC** TO AQUATIC ORGANISMS AND NON-TARGET TERRESTRIAL  
PLANTS. OBSERVE BUFFER ZONES SPECIFIED UNDER **DIRECTIONS FOR  
USE.**

## **FIRST AID**

In case of poisoning, contact a physician or poison control centre.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INGESTED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call 1-833-362-7722 (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

### **STORAGE AND DISPOSAL**

**STORAGE:** Store product in original container in a secure, dry area, away from fertilizer, seed, food or feed. Not for use or storage in or around the home. Keep container tightly closed.

**DISPOSAL:**

**FOR REFILLABLE CONTAINERS:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

### **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the PEST CONTROL PRODUCTS ACT to use this product in a way that is inconsistent with the directions on the label.

<sup>®/TM</sup> Precede is a Registered trademarks/trademarks of FMC Corporation or an affiliate.

[BOOKLET]

**GROUP 2 HERBICIDE**

# Precede™

HERBICIDE

DRY FLOWABLE

For Sale for Use on Wheat (spring and durum), Barley, and Summerfallow in the Prairie Provinces and the Peace River Region of British Columbia

COMMERCIAL

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

REGISTRATION NO.: 29483 PEST CONTROL PRODUCTS ACT

ACTIVE INGREDIENT: Tribenuron-methyl 75%

WARNING – EYE IRRITANT  
POTENTIAL SKIN SENSITIZER  
Contains the allergen sulfites and milk

FMC Corporation  
2929 Walnut Street  
Philadelphia, PA 19104  
1-833-362-7722

**PRECAUTIONS:** WARNING - EYE IRRITANT.  
POTENTIAL SKIN SENSITIZER.  
DO NOT APPLY BY AIR.  
KEEP OUT OF REACH OF CHILDREN.  
MAY BE HARMFUL IF SWALLOWED, INHALED OR  
ABSORBED THROUGH SKIN. AVOID BREATHING SPRAY  
MIST. USE WITH ADEQUATE VENTILATION.  
AVOID CONTACT WITH SKIN, EYES AND CLOTHING.  
WEAR SUITABLE CHEMICAL RESISTANT GLOVES AND  
GOGGLES OR FACE SHIELD DURING MIXING AND  
LOADING.  
MAY IRRITATE SKIN AND EYES. WHEN USING, DO NOT  
EAT, DRINK OR SMOKE.  
WASH THOROUGHLY WITH SOAP AND WATER AFTER  
HANDLING. REMOVE CONTAMINATED CLOTHING  
IMMEDIATELY AFTER USE.  
DO NOT ENTER OR ALLOW WORKER ENTRY INTO TREATED AREAS FOR  
12 HOURS FOLLOWING APPLICATION.

**FIRST AID:**

In case of poisoning, contact a physician or poison control centre  
IMMEDIATELY

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

**IF INGESTED:** Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

For medical emergencies call **1-800-331-3148** (24 hours).

**TOXICOLOGICAL INFORMATION:** Treat symptomatically.

#### IMPORTANT INFORMATION - READ BEFORE USING

Injury to or loss of desirable trees, shrubs, flowers, vegetables, fruits or other vegetation may result from failure to observe the following: Do not apply, drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants. Do not contaminate any body of water, including irrigation water that may be used on crops.

DO NOT APPLY BY AIR.

Do not apply in high winds or on hot days (over 27 degrees C).

Carefully observe sprayer clean-up instructions, as spray tank residue may damage crops other than spring wheat, durum wheat, barley or summerfallow. Thoroughly clean all traces of tank mix from application equipment immediately after use. Flush tank, pump, hoses, and boom with several changes of water after removing nozzle tips and screens (clean these parts separately). Wash away any spray mixture from the outside of spray tank, nozzle or spray rig.

A MINIMUM RECROPPING INTERVAL OF 2 MONTHS SHOULD BE LEFT BETWEEN THE APPLICATION OF THIS PRODUCT AND SEEDING OF THE NEXT CROP. After 2 months, the following crops can be seeded: canola, flax, lentils, and alfalfa.

DO NOT APPLY PRECEDE™ WITH A CARRIER OTHER THAN WATER OR ANY ADDITIVE OTHER THAN THOSE INDICATED ON THIS LABEL. Other carriers may accelerate breakdown of this product and reduce its effectiveness.

#### ENVIRONMENTAL HAZARDS:

· **TOXIC** to aquatic organisms and non-target terrestrial plants. Observe buffer zones specified under **DIRECTIONS FOR USE**

#### GENERAL INFORMATION

Precede™ provides control of certain broad-leaved weeds in spring planted wheat (including durum), barley and summerfallow.

Warm, moist growing conditions promote active weed growth and enhance the activity of Precede™ by allowing maximum foliar uptake and contact activity. Weeds hardened off by environmental stress such as cold weather, drought stress or excessive heat may not be adequately controlled or suppressed, and regrowth may occur. Precede™ may only be applied using ground spray equipment. For best results, ensure thorough spray coverage of target weeds.

Precede™ rapidly stops growth of susceptible weeds. However, typical symptoms (discolouration) of dying weeds may not be noticeable for 1 to 3 weeks after application, depending on growing conditions and weed susceptibility. Degree of control and duration of effect depend on weed sensitivity, weed size, spray coverage and growing conditions. Activity of the herbicide mixture may be delayed by cold, dry conditions after application.

## DIRECTIONS FOR USE:

**Field sprayer application:** **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

### Buffer Zones:

Use of the following spray methods or equipment **DO NOT** require a buffer zone: hand-held or backpack sprayer, spot treatment, inter-row hooded sprayer, soil drench and soil incorporation.

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of Application	Crop	Buffer Zones (metres) Required for the Protection of:		
		Aquatic Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field sprayer*	Summerfallow, fields to be planted to wheat (spring and durum), barley (only as pre-seeding herbicide)	0	0	3
	Wheat and barley	1	0	4

\* For field sprayer application, buffer zones can be reduced with the use of drift-reducing spray shields.

When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled buffer zone can be reduced by 30%.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) [fine medium or coarse] classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

## **FOR WEED CONTROL IN SPRING WHEAT (INCLUDING DURUM) AND BARLEY**

Precede™ + 2,4-D ESTER TANK MIX:

Apply Precede™ at 10 grams per hectare with 2,4-D Ester at 420 g/ha acid equivalent (0.6 L/ha 2,4-D Ester LV-700). Agral<sup>†</sup> 90 is not required in this tank mix.

### **WEEDS CONTROLLED**

Annual sunflower  
Ball mustard  
Canada thistle (top growth control)  
Cow cockle  
Flixweed (fall rosettes and spring seedlings)  
Hare's-ear mustard  
Indian mustard  
Kochia (2-10 leaf)  
Lamb's-quarters  
Narrow-leaved hawk's-beard (fall rosettes and spring seedlings)  
Prickly lettuce  
Redroot pigweed  
Russian pigweed  
Russian thistle  
Shepherd's-purse (fall rosettes and spring seedlings)  
Stinkweed (fall rosettes and spring seedlings)



Sweetclover  
 Thyme-leaved spurge  
 Tumble mustard  
 Wild mustard  
 Wild radish  
 Wormseed mustard  
 WEEDS SUPPRESSED\*

Wild buckwheat

\* Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

#### WHEN TO APPLY

Apply the tank mix from the full 3-leaf stage to just before the flag leaf (shot blade) stage of wheat (including durum) and barley, prior to head emergence. Do not apply after head has emerged.

For best results apply to young, emerged actively growing weeds that are less than 10 cm tall or across, and before the crop canopy closes. Weeds that emerge after treatment will not be controlled.

For wild buckwheat, apply to ACTIVELY GROWING wild buckwheat in the 1- to 3-leaf stage. Under dry conditions, suppression may be reduced. Plants larger than the 3-leaf stage may regrow after treatment.

Precede™ + ASSERT† + 2,4-D ESTER TANK MIX:

For control of WILD OATS and control or suppression of broadleaf weeds in spring (including durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
Precede™	10 g/ha
ASSERT†	1.3 to 1.6 L/ha
2,4-D Ester LV-700	0.6 L/ha

Use the low rate of ASSERT when the majority of wild oats are in the 1 to 3-leaf stage and the higher rate if wild oats are in the 4-leaf stage. Apply when the crop is in the full 3-leaf stage.

Precede™ + 2,4-D ESTER + PUMA† 120 SUPER TANK MIX:

For control of GREEN FOXTAIL (WILD MILLET) and control or suppression of broadleaf weeds in spring (including durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
Precede™	10 g/ha
PUMA† 120 SUPER	385 mL/ha
2,4-D Ester LV-700	0.6 L/ha

Apply this tank mix from the full 3-leaf stage to the 6-leaf + 3 tiller stage of wheat (including durum) and the full 3-leaf stage to the 5-leaf + 2 tiller stage of barley.

For green foxtail control, apply at the 1-6 leaf (up to emergence of the 3<sup>rd</sup> tiller) stage.

### **FOR WEED CONTROL IN SPRING WHEAT (EXCLUDING DURUM) AND SPRING BARLEY**

Precede™ + 2,4-D ESTER + BANVEL† or BANVEL† II TANK MIX:

For control of sulfonylurea resistant kochia and control or suppression of broadleaf weeds in spring (excluding durum) wheat and barley, apply the tank mix as follows:

TANK MIX PARTNER	RATE
Precede™	10 g/ha
BANVEL† or BANVEL† II	110 mL/ha
2,4-D Ester LV-700 (or acid equivalent)	0.6 L/ha (or 420 g/ha acid equivalent)

Apply the tank mix from the full 3-leaf stage to the 6-leaf + 3 tiller stage of wheat (including durum) and barley and to kochia plants that are 2.5 – 10 cm in height. Expect superior results when kochia is small.

## **FOR CONTROL OF FLIXWEED AND STINKWEED (FALL ROSETTES AND SPRING SEEDLINGS) IN SUMMERFALLOW**

Precede™ + 2,4-D ESTER TANK MIX:

Apply Precede™ at 10 g/ha with 2,4-D Ester at 420 g/ha (0.6 L 2,4-D Ester LV-700). AGRAL<sup>†</sup> 90 is not required in this tank mix.

Apply the tank mix in the spring after emergence, up to the early flowering stage of the target weeds. Only weeds that have emerged at the time of application will be controlled.

## **CONTROL OF BROADLEAF WEEDS AND GRASSES IN FALLOW AND PRE-SEED TO SPRING WHEAT (INCLUDING DURUM) AND BARLEY.**

Fields treated with Precede™ can be seeded to spring wheat (including durum) or barley a minimum of 24 hours after application.

Apply Precede™ at 10 grams per hectare, tank mixed with one of the following companion herbicides and adjuvant (where required) in a total spray volume of 55-110 L/ha.

Consult the labels of the glyphosate products to be used as tankmix partners with Precede™ for specific instructions of "Directions for Use" and restrictions.

*With summerfallow treatment, allow at least 10 days to elapse before tillage. Only weeds emerged at time of application will be controlled.*

Companion Herbicide	Application Rate (grams acid equivalent per hectare)	Application Stage	Weeds Controlled	Weeds Suppressed *
Glyphosate (present as potassium salt, isopropylamine salt, ammonium salt or trimethylsulfonium salt)	450 g ae/ha	Up to 8 cm	Canada fleabane, Common ragweed, Narrow-leaved hawk's beard	
		Up to 15 cm	Dandelion, Downy brome, Flixweed, Giant foxtail, Green foxtail, Hemp nettle, Kochia, Lady's thumb, Lamb's-quarters, Persian dandelion, Redroot pigweed, Russian thistle, Stinkweed, Volunteer barley, Volunteer canola (including glyphosate-tolerant varieties), Volunteer flax, Volunteer wheat, Wild mustard, Wild oats	
		Up to 3-leaf Rosette	Wild buckwheat, Cow cockle	
Glyphosate (present as potassium salt, isopropylamine salt, ammonium	270 g ae/ha	Any stage Seedlings and rosettes (non-flowering)	Volunteer canola (including glyphosate tolerant varieties)	Narrow-leaved hawk's beard

salt or trimethylsulfonium salt) plus  Agral <sup>†</sup> 90	0.35% v/v	Up to 8 cm	Green foxtail, Lady's thumb, Stinkweed, Volunteer barley, Volunteer wheat, Wild mustard, Wild oats (1-3- leaf) Wild buckwheat (1-3 leaf)	
		Up to 10 cm	Kochia, Lamb's-quarters,	Redroot pigweed, Russian thistle
Glyphosate (present as potassium salt, isopropylamine salt, ammonium salt or trimethylsulfonium salt) plus  Agral <sup>†</sup> 90	360 g ae/ha  0.35 % v/v	Rosettes (non-flowering)	Dandelion (top growth)	Canada thistle (top growth)

\*Weed suppression is a visual reduction in weed competition (reduced population or vigour) as compared to an untreated area. Degree of suppression will vary with size of weed and environmental conditions prior to and following treatment.

## MIXING INSTRUCTIONS

1. Add one-third of the necessary volume of water to the spray tank.
2. With the agitator running, add the required amount of Precede™.
3. When Precede™ is in suspension, add the required amount of the appropriate tank mix partners.
4. Specific Tank Mix Instructions: Precede™ alone: add AGRAL<sup>†</sup> 90 at a rate of 2 L/1000 L spray solution (0.2% v/v).

Precede™ + 2,4-D Ester: once Precede™ DF 75% HERBICIDE is completely in suspension, add the required amount of 2,4-D Ester while maintaining agitation. Do not add a surfactant.

Precede™ + 2,4-D Ester + ASSERT<sup>†</sup>: once Precede™ DF 75% HERBICIDE is completely in suspension, add the required amount of Spray Water pH Adjuster and agitate until dissolved. Add the required amount of ASSERT<sup>†</sup> while maintaining agitation. Add 2,4-D Ester. Addition of a surfactant is not required.

Precede™ + 2,4-D Ester + BANVEL<sup>†</sup> or BANVEL<sup>†</sup> II: add 2,4-D Ester last. A surfactant is not required with this mixture.

Precede™ + 2,4-D Ester + PUMA<sup>†120</sup> SUPER: add the required amount of PUMA<sup>120</sup> SUPER last. A surfactant is not required with this mixture.

Precede™ + Glyphosate (present as potassium salt, isopropylamine salt, ammonium salt or trimethylsulfonium salt): once Precede™ is completely in suspension, add the required amount of glyphosate (present as potassium salt, isopropylamine salt, ammonium salt or trimethylsulfonium salt) while maintaining agitation. . Once components are completely mixed, if required, add Agral 90 at 3.5L per 1000 L of spray solution (0.35% v/v)

5. Add the rest of the water. If an antifoam agent is required, add this last.
6. Use a minimum spray volume of 55 L/ha on spring wheat (including durum), barley and summerfallow.
7. For repeat tankloads, reduce the material remaining in the tank to 10% of the original volume or less before proceeding with step 1. If this is not

possible, pre-slurry Precede™ in a small amount (5-10 L) of water before adding to the tank.

Continuous agitation is required to keep Precede™ in suspension. If the spray preparation is left standing without agitation, thoroughly agitate to resuspend the mixture before spraying. Use spray preparation within 24 hours or product degradation may occur.

## **MINIMUM RECROPPING INTERVAL AND CROP ROTATION GUIDELINES**

A minimum recrop interval of 2 months should be left between the application of this product and seeding of the next crop. The following crops can be seeded two months after application of Precede™ + 2,4-D Ester: canola, flax, lentils, and alfalfa.

## **EQUIPMENT-SPRAY VOLUMES**

Apply the spray mixture uniformly with properly calibrated ground equipment only. Ensure thorough coverage and a uniform spray pattern. Flat fan nozzles are recommended (minimum 55 litres spray volume per hectare). Use 50 mesh filter screens or larger (metal or nylon).

Continuous agitation is required to keep the spray materials in suspension. Apply uniformly and avoid overlapping. Shut off spray boom while starting, turning, slowing or stopping to prevent crop injury from over application.

Reduce drift with high water volume, low pressure, coarse sprays and drop nozzles. Do not apply with more than 275 kPa spray pressure.

DO NOT APPLY BY AIR.

## **SPRAYER CLEANUP**

It is difficult to remove all traces of 2,4-D Ester and MCPA Amine from a sprayer. A sprayer used to apply these herbicides should not be used to spray crops sensitive to these herbicides. The following clean-out procedure will remove traces of Precede™ from the sprayer.

1. Drain tank; then flush tank, boom and hoses with clean water for a minimum of ten minutes to remove all visible residues.
2. Fill the tank with clean water then add 1 litre household AMMONIA (containing a minimum of 3% ammonia) per 100 litres of water. Fill

boom and hoses with solution and allow sprayer to sit for 15 minutes. Drain.

3. Repeat step 2.
4. Nozzles and screens should be removed and cleaned separately. To remove traces of ammonia, rinse the tank, hoses and booms thoroughly with clean water.
5. Dispose of tank rinseate according to Provincial directions.

**CAUTION:** Do not use ammonia with chlorine bleach. Using ammonia with chlorine bleach will release a gas with a musty chlorine odour that can cause eye, nose, throat, and lung irritation. Do not clean equipment in an enclosed area.

## **RESISTANCE-MANAGEMENT RECOMMENDATIONS**

For resistance management, Precede™ is a Group 2 herbicide. Any weed population may contain or develop plants naturally resistant to Precede™ and other Group 2 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- ◆ Where possible rotate the used of Precede™ or other Group 2 herbicides with different herbicide groups that control the same weeds in a field.
- ◆ Use tank mixtures with herbicides from a different group when such use is permitted
- ◆ Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- ◆ Monitor treated weed populations for resistance development.
- ◆ Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- ◆ Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- ◆ For further information or to report suspected resistance, contact your local FMC representative or the FMC hotline at 1-833-362-7722 for further information.



## **USE PRECAUTIONS:**

Licensed wheat and barley varieties have shown acceptable tolerance to Precede™. Consult FMC Corporation prior to use on unlicensed varieties for crop tolerance information.

Do not apply to irrigated land where tail water will be used to irrigate other cropland.

Do not contaminate irrigation water.

Do not use on highly variable soils that have large gravelly or sandy areas, eroded knolls or calcium deposits.

If rain occurs soon after application, control may be reduced. At least 4-6 hours of dry weather is needed to allow the tank mixture to be absorbed by weed foliage. Environmental conditions that slow the drying of the spray mixture on the foliage such as high relative humidity, cool air temperatures or cloud cover, may increase the time required.

Use a maximum of 10 grams of Precede™ per hectare per year on spring planted wheat, barley, and summerfallow.

Precede™ applied to summerfallow, spring wheat, durum wheat or barley that is stressed by severe conditions such as drought, low fertility, saline soils, waterlogged soils (soils at or near field capacity), disease or insect damage may result in crop injury. Drought, disease or insect damage following application may also result in crop injury, grade or yield loss.

Do not apply to spring wheat, durum wheat or barley undersown with legumes or grasses as injury to the forage may result.

Do not remove soil from treated fields for use on lawns, in gardens or for backfill.

Under certain conditions such as prolonged cool weather temperature or wide fluctuations in day/night temperatures just prior to or soon after treatment, temporary yellowing and/or crop stunting may occur.

**OVERSPRAY OR DRIFT TO IMPORTANT WILDLIFE HABITATS SUCH AS SHELTERBELTS, WETLANDS, SLOUGHS OR DRY SLOUGH BORDERS, OR WOODLOTS, SHOULD BE AVOIDED. LEAVE A 15-METER BUFFER ZONE BETWEEN THE LAST SPRAY SWATH AND THE EDGE OF ANY OF THESE HABITATS.**

Control of weeds growing in wheel tracks may be reduced if Precede™ is applied under dry, dusty conditions.

## **STORAGE AND DISPOSAL**

### **STORAGE:**

Store products in original containers only, away from fertilizer, seeds, food or feed.

Not for use or storage in or around the home. Keep container closed.

### **DISPOSAL:**

#### **FOR REFILLABLE CONTAINERS:**

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

## **NOTICE TO USER**

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the Pest Control Products Act to use this product in a way that is inconsistent with the directions on the label.

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact FMC Corporation at 1-833-362-7722 for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by FMC Corporation.

®/™ Precede is a Registered trademarks/trademarks of FMC Corporation or an affiliate.

† All other products mentioned are trademarks of their respective companies.