

 **Dow AgroSciences**
Restore™ A Herbicide

GROUP	4	HERBICIDE
--------------	----------	------------------

This component must only be used in Restore Herbicide

COMMERCIAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

GUARANTEE: Aminopyralid, present as triisopropanolamine salt 240 g a.e./L
Solution

REGISTRATION NO. 28551 PEST CONTROL PRODUCTS ACT

NET CONTENTS: 3.0 L – 450 L

Dow AgroSciences Canada Inc.
2400, 215 – 2nd Street S.W.
Calgary, Alberta
T2P 1M4
1-800-667-3852

®™Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN

PERSONAL PROTECTIVE EQUIPMENT

Applicators must wear coveralls and chemical resistant gloves.

Mixers and loaders handling concentrated product, as an extra precaution, should wear coveralls, chemical resistant gloves, goggles, and rubber boots.

Worker reentry – Do not enter or allow worker entry to treated area for 12 hours following application.

If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at www.croplife.ca.

SPRAY DRIFT PRECAUTIONS

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

OPERATOR USE PRECAUTIONS

- Wash hands before eating, drinking, using tobacco or using the washroom.
- If herbicide penetrates clothing, remove immediately; then wash thoroughly and put on clean clothing.
- Remove personal protective equipment immediately after handling this product. As soon as possible, wash thoroughly and change into clean clothing. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash personal protective equipment separate from household laundry.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient.

ENVIRONMENTAL HAZARDS

TOXIC to terrestrial and aquatic plants. Observe terrestrial and aquatic buffer zones specified under DIRECTIONS FOR USE.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL

Recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Restore Herbicide



Dow AgroSciences

Restore™ Herbicide

GROUP	4	HERBICIDE
-------	---	-----------

FOR CONTROL OF BROADLEAF WEEDS IN RANGELAND AND PASTURE AREAS OF CANADA

COMMERCIAL

READ THE LABEL AND BOOKLET BEFORE USING
KEEP OUT OF REACH OF CHILDREN

This co-package contains:

Restore A Herbicide

REGISTRATION NO. 28551 PEST CONTROL PRODUCTS ACT

GUARANTEE: Aminopyralid, present as triisopropanolamine salt 240 g a.e./L
solution

Restore B Herbicide

REGISTRATION NO. 28552 PEST CONTROL PRODUCTS ACT

GUARANTEE: 2,4-D, present as dimethylamine salt 564 g a.e./L
solution

WARNING POISON

NET CONTENTS: Restore A Herbicide 3.0 L – 450 L
 Restore B Herbicide 7.65 L – 450 L

Dow AgroSciences Canada Inc.

2400, 215 – 2nd Street S.W.

Calgary, Alberta

T2P 1M4

1-800-667-3852

®™Trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow

PRECAUTIONS

KEEP OUT OF REACH OF CHILDREN.

Avoid contact with skin, eyes and clothing. Wash concentrate from skin or eyes immediately. Do not inhale fumes. Avoid breathing spray mist. Avoid spray drift. Store the container tightly closed and away from seeds, feeds, fertilizer, plants and foodstuffs. Avoid contamination of ponds, streams, rivers and other water sources.

PROTECTIVE CLOTHING AND EQUIPMENT

All Application Equipment

- When mixing/loading, wear coveralls over a long-sleeved shirt, long pants, chemical-resistant gloves, socks, shoes, protective eye wear (face shield or safety glasses) and rubber boots. Rinse gloves before removal.
- When applying or during clean-up and repair, wear coveralls over a long sleeved shirt, long pants, chemical resistant gloves, socks and shoes. Rinse gloves before removal.
- When handling more than 470 L of Restore B Herbicide per day workers must also use a closed system.

Application Using Aerial Equipment

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities.
- No human flaggers are permitted.

Application Using Groundboom Equipment

- Applicators must wear coveralls over a long-sleeved shirt and long pants. Chemical-resistant gloves must also be worn during clean-up and repair activities

Worker reentry - Do not enter or allow worker entry to treated area for 12 hours following application.

Refer to Directions for Use section for any additional requirements.

OPERATOR USE PRECAUTIONS

- Wear freshly laundered clothing and clean protective equipment daily.
- Rinse gloves before removal.
- Wash hands before eating, drinking, using tobacco or using the toilet.
- If herbicide penetrates clothing, remove immediately, then wash thoroughly and put on clean clothing. Throw away clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate.
- Remove clothing and personal protective equipment immediately after handling this product. As soon as possible, wash hands and exposed skin thoroughly with soap and water and change into clean clothing. Follow manufacturer's instructions for cleaning personal protective clothing and equipment. If no such instructions for washables are provided, use detergent and hot water. Keep and wash clothing and personal protective equipment separate from household laundry.
- After work, remove all clothing and shower using soap and water.

TRANSFER SYSTEM

For containers larger than 20L: Use a transfer system that avoids open pouring when transferring the liquid concentrate from such containers into the spray tank.

SPRAY DRIFT PRECAUTIONS

Apply only when the potential for drift to areas of human habitation or areas of human activity such as houses, cottages, schools and recreational areas is minimal. Take into consideration wind speed, wind direction, temperature, application equipment and sprayer settings.

FIRST AID

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

TOXICOLOGICAL INFORMATION

No specific antidote. Employ supportive care. Treatment should be based on judgment of the physician in response to reactions of the patient. 2,4-D may cause severe irritation to the eyes. Overexposure to 2,4-D may cause coughing, burning, dizziness or temporary loss of muscle coordination. Other possible effects of overexposure include fatigue, muscle weakness or nausea. Treat symptomatically.

ENVIRONMENTAL HAZARDS

- Toxic to small wild mammals, birds, aquatic organisms and non-target terrestrial plants.
- Observe buffer zones specified under DIRECTIONS FOR USE.
- This product will harm other broadleaved plants in the vicinity of the treatment area. If applying this product using a handheld sprayer, do not directly spray or allow the spray to drift onto ornamentals or gardens.
- Do not spray exposed roots of trees and ornamentals.

LEACHING

The use of this chemical may result in contamination of groundwater particularly in areas where soils are permeable (e.g. sandy soil) and/or the depth to the water table is shallow.

RUNOFF

- To reduce runoff from treated areas into aquatic habitats avoid application to areas with moderate to steep slope, compacted soil, or clay.
- Avoid application of this product when heavy rain is forecast.
- Contamination of aquatic areas as a result of runoff may be reduced by including a strip of untreated vegetation between the treated area and the edge of the water body.

STORAGE

To prevent contamination store this product away from food or feed.

DISPOSAL

Recyclable containers:

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

3. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
4. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial requirements.

Returnable containers:

Do not reuse this container for any purpose. For disposal, this empty container may be returned to the point of purchase (distributor/dealer).

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency. Contact the manufacturer and the provincial regulatory agency in case of a spill, and for clean-up of spills.

DIRECTIONS FOR USE

READ ALL DIRECTIONS CAREFULLY BEFORE APPLYING. FAILURE TO FOLLOW LABEL INSTRUCTIONS MAY RESULT IN ERRATIC WEED CONTROL OR CROP DAMAGE.

GENERAL USE PRECAUTIONS

Avoid spray drift to any desirable vegetation as damage may occur. Avoid contamination of ponds, streams, rivers and other water sources. Do not spray during periods of high winds when spray is likely to drift. Coarse sprays are less likely to drift. Do not spray during periods of high temperatures when desirable vegetation may be under stress as damage to the desirable vegetation may occur. Applications should be avoided under conditions of drought or other environmental stresses.

For best results, apply Restore Herbicide to young weeds that are actively growing at time of application.

Restore Herbicide is highly active against broadleaf plants. This product is recommended for use on areas where loss of broadleaf forage plants, including legumes, can be tolerated.

Do not apply more than 0.50 L per hectare of Restore A Herbicide per annual growing season to rangeland and pasture.

Only one application per growing season is permitted.

Restore Herbicide cannot be applied on domestic or commercial turf grass.

- As this product is not registered for the control of pests in aquatic systems, DO NOT use to control aquatic pests.
- DO NOT contaminate irrigation/drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. Boom height must be 60 cm or less above the crop or ground.

Aerial application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. DO NOT apply when wind speed is greater than 16 km/h at flying height at the site of application. DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) coarse classification. To reduce drift caused by turbulent wingtip vortices, the nozzle distribution along the spray boom length MUST NOT exceed 65% of the wing- or rotorspan. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Spray Equipment Precautions

Clean all spray equipment thoroughly before use. Select nozzle tips to apply the recommended volume per hectare. Higher water volumes will reduce the risk of crop injury. Calibrate the sprayer and adjust the boom height to ensure uniform coverage.

Nozzles must be in good condition and spraying the same volume. Flat fan nozzle tips are recommended.

Equipment Clean-Up

Thoroughly clean the herbicide sprayer before applying any other chemical on crops that may be damaged (see GENERAL USE PRECAUTIONS). Wash the outside of the sprayer and drain the tank completely. Remove and clean filters, screens and nozzle tips separately. Fill the sprayer tank with clean water. With the agitator running flush out the lines and boom, then drain. Fill the sprayer with clean water and for each 100L add 1 L household ammonia. Operate the pump and agitator for 15 minutes. If possible, let the solution remain in the tank and hoses overnight; recirculate and flush out the lines and boom, then drain. Rinse out twice with clean water, recirculating and draining each time. DO NOT contaminate irrigation or drinking water when cleaning the sprayer.

Tank Mixtures

In some cases, tank mixing a pest control product with another pest control product or a fertilizer can result in biological effects that could include, but are not limited to: reduced pest efficacy or increased host crop injury. The user should contact Dow AgroSciences Canada Inc at 1-800-667-3852 or www.dowagro.ca for information before mixing any pesticide or fertilizer that is not specifically recommended on this label. The user assumes the risk of losses that result from the use of tank mixes that do not appear on this label or that are not specifically recommended by Dow AgroSciences Canada Inc.

APPLICATION DIRECTIONS

Mixing

Shake this product well before using. Half fill the spray tank with clean water. Add the required amount of product and agitate thoroughly. Fill the tank with clean water and agitate well before use.

NOTE: If spray solution has been left standing, agitate thoroughly before use.

Application Rates

Herbicide Component	Rate	Weeds Controlled
Restore A Herbicide	0.29 L/ha	Control of annual sow thistle, bluebur, burdock (<4 leaf), cocklebur, common plantain, flixweed, goat's beard, prickly lettuce, ragweeds, stinging nettle, sweet clover, curled dock (<4 leaf), hawkweed, peppergrass, season long control of Canada thistle, spotted knapweed, scentless chamomile, Canada goldenrod, and top growth control of blue lettuce, bull thistle, buttercup, gumweed, hoary cress, perennial sowthistle
Restore B Herbicide	1.5 L/ha	

Herbicide Component	Rate	Weeds Controlled
Restore A Herbicide	0.38 L/ha	all weeds listed above plus season long control of absinth wormwood, dandelion
Restore B Herbicide	1.9 L/ha	

Herbicide Component	Rate	Weeds Controlled
Restore A Herbicide	0.50 L/ha	all weeds listed above plus season long control of western snowberry, common tansy
Restore B Herbicide	2.6 L/ha	

Maximum 1 application per year.

Preharvest/Grazing Intervals

- There is no restriction on livestock (except lactating dairy animals) grazing in treated areas.
- Do not permit lactating dairy animals to graze fields within 7 days after application.
- Allow 3 days of grazing on an untreated pasture (or feed untreated hay) before transferring livestock to areas where sensitive broadleaf crops may be grown. Withdraw meat animals from treated fields at least 3 days before slaughter. Do not harvest forage or cut for hay within 30 days after application.

MIXING METHODS**Mixing with Water**

To prepare the spray, add half the desired amount of water in the spray tank. Then with agitation, add the recommended amount of Restore A Herbicide and then Restore B Herbicide. Finally, with continued agitation, add the rest of the water.

APPLICATION METHODS**Broadcast Treatment (Ground and Aerial Applications)**

Applications of Restore Herbicide may be made once grasses are well-established (have developed a good secondary root system and show good vigor).

(1) Ground Application

Using ground equipment, apply Restore Herbicide as a broadcast treatment to control listed broadleaf weeds and woody plants. Apply Restore Herbicide at the recommended rates in a minimum spray volume of 100 L/ha.

(2) Aerial Application

Use Restore Herbicide as a broadcast treatment by air to control listed broadleaf weeds and woody plants. Apply Restore Herbicide at the recommended rates in a minimum spray volume of 19 L/ha by air. Refer to the tank-mix partner label for additional instructions and precautions.

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the "National Aerial Pesticide Application Manual, developed by the Federal/Provincial/Territorial Committee on Pest Management and Pesticides".

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other non-target areas. Specified buffer zones should be observed. Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto the aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application. Do not use human flaggers.

The field crew and mixer/loaders must wear coveralls over a long-sleeved shirt, long pants, chemical resistant gloves and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating, drinking, using tobacco or the toilet. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-800-667-3852 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

- Avoid spray drift at the application site. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. Users are responsible for considering all these factors when making decisions.

SPRAYER CLEAN-OUT INSTRUCTIONS

Do not use spray equipment used to apply Restore Herbicide for other applications to land planted to, or to be planted to susceptible crops or desirable sensitive plants, unless it has been determined that all residue of this herbicide has been removed by thorough cleaning of equipment. Do not contaminate water through cleaning of equipment or disposal of wastes.

Equipment used to apply Restore Herbicide should be thoroughly cleaned before reusing to apply any other chemicals.

1. Rinse and flush application equipment thoroughly after use. Dispose of rinse water in non-cropland area away from water supplies.
2. Rinse a second time, adding 1 litre of household ammonia or tank cleaning agent for every 1000 L of water. Circulate the solution through the entire system so that all internal surfaces are contacted (15 to 20 minutes). Let the solution stand for several hours, preferably overnight.
3. Flush the solution out the spray tank through the boom.
4. Rinse the system twice with clean water, recirculating and draining each time.
5. Nozzles and screens should be removed and cleaned separately.

BUFFER ZONES TO PROTECT SENSITIVE HABITATS

Seasonal water bodies require buffer zones if there is water in them during application. Water bodies which do not fill on an annual basis need not be buffered.

Buffer Zones – Sensitive Habitats

The buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands), and estuarine/marine habitats.

Buffer Zone Table

Method of Application	Crop		Buffer Zones (metres) Required for the Protection of Terrestrial and Aquatic Habitat:			
Field sprayer	Pastures, rangelands		10			
	Aerial	Pastures, rangelands	Spray Quality	ASAE Coarse (VMD = 385.22 µm)	ASAE Coarse- Very Coarse (VMD = 439.39 µm)	ASAE Very Coarse (VMD = 477.94 µm)
Fixed wing			175	125	100	80
Rotary wing			150	100	90	70

NOTE: Applicators may recalculate a site-specific buffer zone by combining information on current weather conditions and spray configuration for the following applications: all airblast applications, and for field and aerial applications which specify the following droplet size category wording on the product label: 'DO NOT apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) [Fine or Medium or Coarse] classification.' To access the Buffer Zone Calculator, please visit the Pest Management Regulatory Agency web site.

When a tank mixture is used, consult the labels of the tank-mix partners and observe the largest (most restrictive) buffer zone of the products involved in the tank mixture.

RESISTANCE MANAGEMENT RECOMMENDATIONS

For resistance management, Restore Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Restore Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual

chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Restore Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical), cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Dow AgroSciences Canada Inc. at 1-800-667-3852 or at www.dowagro.ca.

NOTICE TO USER: This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.