



# MOCCASIN® II PLUS

## HERBICIDE

**Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

For weed control in beans, peas, and lentils; corn; cotton; grasses grown for seed; horseradish; peanuts; potatoes; pumpkin; rhubarb; safflowers; sweet, grain, or forage sorghum; soybeans; soybeans, immature seed; sugarcane; sugar beets; sunflowers; tomatoes.

<b>Active Ingredient:</b>	<b>BY WT.</b>
S-metolachlor (CAS No. 87392-12-9) .....	82.4%
<b>Other Ingredients:</b> .....	17.6%
<b>Total:</b> .....	100.0%

MOCCASIN® II PLUS Herbicide contains 7.64 lbs of active ingredient per gallon. MOCCASIN II PLUS Herbicide is formulated as an emulsifiable concentrate (EC).

EPA Reg. No. 70506-333

### KEEP OUT OF REACH OF CHILDREN WARNING - AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
<b>If in eyes</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a Poison Control Center or doctor for treatment advice.</li> </ul>
<b>If swallowed</b>	<ul style="list-style-type: none"> <li>• Call a Poison Control Center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to by a Poison Control Center or doctor.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If inhaled</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a Poison Control Center or doctor for further treatment advice.</li> </ul>
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. For medical treatment, call Rocky Mountain Poison and Drug Safety at 1-866-673-6671.</p>	
<p><b>FOR CHEMICAL EMERGENCY: Spill, leak, fire, exposure, or accident, call CHEMTREC 1-800-424-9300.</b></p>	

See inside for additional Precautionary Statements and complete Directions for Use.



NET CONTENTS: \_\_\_\_\_ GALLONS



## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### WARNING

Causes substantial but temporary eye injury. Harmful if absorbed through skin. Harmful if swallowed. Do not get in eyes or on clothing. Wear protective eyewear (goggles, face shield, or safety glasses).

#### Personal Protective Equipment (PPE)

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

#### Engineering Control Statements

Mixers and loaders supporting aerial applications are required to use closed systems. The closed system must be used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d)]. When using the closed system, the mixers' and loaders' PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.607(d-f)], the handler PPE requirements may be reduced or modified as specified in the WPS.

#### User Safety Recommendations

**Users should:**

- Wash hands thoroughly with soap and water immediately after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash water or rinsate.

Reporting Ecological Incidents: To report ecological incidents, including mortality, injury, or harm to plants and animals, call UPL NA Inc. at 1-800-438-6071.

#### Groundwater Advisory

S-metolachlor is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

#### Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several weeks or months after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of s-metolachlor from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is expected to occur within 48 hours.

#### Non-target Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of non-target organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of non-target organisms by following label directions intended to minimize spray drift.

#### Mixing/Loading Instructions

When using this product take care to prevent back-siphoning into wells, spills, or improper disposal of excess pesticide, spray mixtures, or rinsates.

Check-valves or antisiphoning devices must be used on all mixing and/or irrigation equipment.

This product may not be mixed or loaded within 50 ft of perennial or intermittent streams and rivers, natural or impounded lakes, and reservoirs. This product may not be mixed/loaded or used within 50 ft of all wells, including abandoned wells, drainage wells, and sink holes. Operations that involve mixing, loading, rinsing, or washing of this product into or from pesticide handling or application equipment or containers within 50 ft of any well are prohibited, unless conducted on an impervious pad constructed to withstand the weight of the heaviest load that may be positioned on or moved across the pad. Such a pad shall be designed and maintained to contain any product spills or equipment leaks, container or equipment rinse or wash water, and rain water that may fall on the pad. Surface water shall not be allowed to either flow over or from the pad, which means the pad must be self-contained. The pad shall be sloped to facilitate material removal. An unroofed pad shall be of sufficient capacity to contain at a minimum 110% of the capacity of the largest pesticide container or application equipment on the pad. A pad that is covered by a roof of sufficient size to completely exclude precipitation from contact with the pad shall have a minimum containment capacity of 100% of the capacity of the largest pesticide container or application equipment on the pad. Containment capacities as described above shall be maintained at all times. The above-specified minimum containment capacities do not apply to vehicles when delivering pesticide shipments to the mixing/loading site.

**NOTICE:** Read the entire **Directions for Use** and **Conditions of Sale and Limitation of Warranty and Liability** before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Endangered Species Protection Requirements:

It is a Federal offense to use any pesticide in a manner that results in an unauthorized "take" (e.g., kill or otherwise harm) of an endangered species and certain threatened species, under the Endangered Species Act section 9. When using this product, you must follow the measures contained in the Endangered Species Protection Bulletin for the area in which you are applying the product. You must obtain a Bulletin no earlier than six months before using this product. To obtain Bulletins, consult <http://www.epa.gov/espp/>, call 1-844-447-3813, or email [ESPP@epa.gov](mailto:ESPP@epa.gov). You must use the Bulletin valid for the month in which you will apply the product.

Use MOCCASIN II PLUS Herbicide only in accordance with directions on this label or in separately published EPA accepted supplemental labeling directions for this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 24 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

### FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, as crop injury may result.

**Sale, use and distribution of this product in Nassau and Suffolk Counties in the State of New York is prohibited.**

## PRODUCT INFORMATION

Observe all precautions, restrictions and limitations on the labels of each product used in tank mixtures. Tank mixtures are permitted only in those states where the tank mix partner is registered. Refer to and follow the label for each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

MOCCASIN II PLUS Herbicide is a selective herbicide for use as a preplant surface-applied, preplant incorporated, or preemergence treatment in water or fluid fertilizer for control of most annual grasses and certain broadleaf weeds in labeled crops.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Precautions

- Dry weather following preemergence application of MOCCASIN II PLUS Herbicide or a tank mixture may reduce effectiveness. Cultivate if weeds develop.
- If MOCCASIN II PLUS Herbicide is incorporated, any supplemental tillage before planting must not exceed the depth of incorporation.
- Certain states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for additional information. It is a violation of this label to deviate from state use regulations.
- Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor, or consistent control at a level below that generally considered acceptable for commercial weed control.

- Injury may occur following the use of MOCCASIN II PLUS Herbicide under abnormally high soil moisture conditions during early development of the crop.

### Restrictions

- **DO NOT** apply under conditions which favor runoff or wind erosion of soil containing this product to nontarget areas.
- To prevent off-site movement due to runoff or wind erosion:
  - **DO NOT** treat powdery dry or light sand soils when conditions are favorable for wind erosion. Under these conditions, the soil surface should first be settled by rainfall or irrigation.
  - **DO NOT** apply to impervious substrates, such as paved or highly compacted surfaces.
  - **DO NOT** use tailwater from the first flood or furrow irrigation of treated fields to treat nontarget crops, unless at least 1/2 inch of rainfall has occurred between application and the first irrigation.

## SPRAY DRIFT MANAGEMENT

### MANDATORY SPRAY DRIFT MANAGEMENT

#### Ground Boom Applications

- User must only apply with the release height recommended by the manufacturer, but no more than 4 ft above the ground or crop canopy.
- Applicators are required to select the nozzles and pressure that deliver medium or coarser droplets (ASABE S572).
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

#### Aerial Applications:

- Do not release spray at a height greater than 10 ft above the ground or vegetative canopy, unless a greater application height is necessary for pilot safety.
- Applicators are required to select the nozzle and pressure that deliver medium or coarser droplets (ASABE S641).
- If the wind speed is 10 miles per hour or less, applicators must use 1/2 swath displacement upwind at the downwind edge of the field. When the wind speed is between 11 - 15 miles per hour, applicators must use 3/4 swath displacement upwind at the downwind edge of the field.
- Do not apply when wind speeds exceed 15 mph at the application site. If the wind speed is greater than 10 mph, the boom length must be 65% or less of the wingspan for fixed wing aircraft and 75% or less of the rotor diameter for helicopters. Otherwise, the boom length must be 75% or less of the wingspan for fixed-wing aircraft and 90% or less of the rotor diameter for helicopters.
- Do not apply during temperature inversions.

#### Boomless Ground Applications:

- Applicators are required to select the nozzle and pressure that deliver a medium or coarser droplet size (ASABE S572.3) for all applications.
- Do not apply when wind speeds exceed 15 mph at the application site.
- Do not apply during temperature inversions.

## SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT. BE AWARE OF NEARBY NON-TARGET SITES AND ENVIRONMENTAL CONDITIONS.

#### Boomless Ground Applications:

- Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

## IMPORTANCE OF DROPLET SIZE

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

### Controlling Droplet Size – Ground Boom

- Volume – Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure – Use the lowest spray pressure recommended for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle – Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

### Controlling Droplet Size – Aircraft

- Adjust Nozzles – Follow nozzle manufacturers' recommendations for setting up nozzles. Generally, to reduce fine droplets, nozzles should be oriented parallel with the airflow in flight.

## BOOM HEIGHT – Ground Boom

For ground equipment, the boom should remain level with the crop and have minimal bounce.

## RELEASE HEIGHT – Aircraft

Higher release heights increase the potential for spray drift.

## SHIELDED SPRAYERS

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

## TEMPERATURE AND HUMIDITY

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

## TEMPERATURE INVERSIONS

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

## WIND

Drift potential generally increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.**

Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

## Herbicide Resistance Management

For resistance management, MOCCASIN II PLUS Herbicide is a Group 15 herbicide. Any weed population may contain or develop plants naturally resistant to MOCCASIN II PLUS Herbicide and other Group 15 herbicides. Weed species with acquired resistance to Group 15 may eventually dominate the weed population if Group 15 herbicides are used repeatedly in the same field. Appropriate resistance-management strategies should be followed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of MOCCASIN II PLUS Herbicide or other Group 15 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field. Whenever possible incorporate multiple weed control practices including mechanical cultivation, biological management practices, and crop rotation.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.

- Adopt an integrated weed-management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation, and that considers tillage (or other mechanical control methods), cultural (e.g. higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties) and other management practices.
- Scout fields before application to identify the weed species present and their growth stage to determine if the intended application will be effective. Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of non-controlled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method including hoeing or tillage. Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields, and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product and switch to another management strategy or herbicide with a different mode of action (MOA), if available. Treat weed escapes with an herbicide with a different MOA or use non-chemical methods to remove escapes. To the extent possible do not allow weed escapes to produce seeds, roots, or tubers.
- Contact your local sales representative, crop advisor, or extension agent to find out if suspected resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed. Contact UPL NA at 1-800-438-6071.

## SOIL TEXTURES AND HERBICIDE RATES

Where rates are based on coarse-, medium-, or fine-textured soils, it is understood that soil textural classes are generally categorized as follows:

COARSE	MEDIUM	FINE
Sand	Loam	Sandy clay loam
Loamy sand	Silt loam	Silty clay loam
Sandy loam	Silt	Clay loam
		Sandy clay
		Silty clay
		Clay

Within rate ranges in the rate tables and elsewhere on this label, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter.

**Note:** MOCCASIN II PLUS Herbicide may be applied preemergence alone, or in combination with tank mix partners specified on this label, following preplant incorporated herbicides when used according to their label instructions, provided that such use is not prohibited on the respective labels.

Thoroughly clean sprayer or other application device before using. Dispose of cleaning solution in a responsible manner. Do not use a sprayer or applicator contaminated with any other materials, or crop damage or clogging of the application device may result.

## MOCCASIN II PLUS HERBICIDE APPLIED ALONE

### Weeds Controlled

MOCCASIN II PLUS Herbicide is taken up by the shoots and/or roots of emerging weeds. This uptake results in the inhibition of shoot and root tissue growth soon after weed germination. Because MOCCASIN II PLUS Herbicide will not control emerged weeds, apply before weed emergence.

If MOCCASIN II PLUS Herbicide is incorporated, do not exceed a 2- to 3-inch depth. Any tillage after the MOCCASIN II PLUS Herbicide incorporation and before planting should not exceed 2 - 3 inches.

Dry weather following application of MOCCASIN II PLUS Herbicide may reduce weed control. Cultivate if weeds develop.

Where reference is made to weeds partially controlled, partial control can either mean erratic control from good to poor or consistent control at a level below that generally considered acceptable for commercial weed control. Control of these weeds can be erratic, due partially to variable weather conditions. The following procedures may improve the control of weeds listed as partially controlled in **Table 1**:

1. Thoroughly till soil to destroy germinating and emerged weeds.
2. Plant crop into moist soil immediately after tillage. If MOCCASIN II PLUS Herbicide is to be used preemergence, apply at planting or immediately after planting.
3. If available, sprinkler irrigate within 2 days after application. Apply 1/2 - 1 inch of water. Use lower water volume (1/2 inch) on *coarse-textured soils* and higher volume (1 inch) on *fine-textured soils*. Also, refer to the section on **Center Pivot Irrigation Application** for this method of applying MOCCASIN II PLUS Herbicide.
4. If irrigation is not possible and rain does not occur within 2 days after planting and application, weed control may be decreased. Under these conditions, a uniform, shallow cultivation is recommended as soon as weeds emerge.

**Table 1**

**Weeds Controlled or Partially Controlled by MOCCASIN II PLUS Herbicide Applied Prior to Weed Emergence**

Common Name	Scientific Name	Control (C) or Partial Control (PC)
<b>GRASSES</b>		
Barnyardgrass	<i>Echinochloa crus-galli</i>	C
Crabgrass, large	<i>Digitaria ischaemum</i>	C
Crabgrass, smooth	<i>Digitaria sanguinalis</i>	C
Crowfootgrass	<i>Dactyloctenium aegyptium</i>	C
Cupgrass, Prairie	<i>Eriochloa contracta</i>	C
Cupgrass, Southwestern	<i>Eriochloa acuminata</i>	C
Cupgrass, woolly	<i>Eriochloa villosa</i>	PC <sup>1</sup>
Foxtail, bristly	<i>Setaria verticillata</i>	C
Foxtail, giant	<i>Setaria faberi</i>	C
Foxtail, green	<i>Setaria viridis</i>	C
Foxtail, millet	<i>Setaria italica</i>	C
Foxtail, yellow	<i>Setaria pumila</i>	C
Goosegrass	<i>Eleusine indica</i>	C
Johnsongrass (seedling)	<i>Sorghum halepense</i>	PC
Millet, wild-proso	<i>Panicum miliaceum</i>	PC <sup>1</sup>
Panicum, fall	<i>Panicum dichotomiflorum</i>	C
Panicum, Texas	<i>Panicum texanum</i>	PC
Rice, red	<i>Oryza punctata</i>	C
Sandbur, field	<i>Cenchrus spinifex</i>	PC
Ryegrass, Italian	<i>Lolium multiflorum</i>	C
Sandbur, Southern	<i>Cenchrus spinifex</i>	PC
Shattercane	<i>Sorghum bicolor</i>	PC
Signalgrass, broadleaf	<i>Urochloa platyphylla</i>	C
Sorghum (volunteer)	<i>Sorghum bicolor</i>	PC
Witchgrass	<i>Panicum capillare</i>	C

(continued) 5

**Table 1 (continued)**

**Weeds Controlled or Partially Controlled by MOCCASIN II PLUS Herbicide Applied Prior to Weed Emergence**

Common Name	Scientific Name	Control (C) or Partial Control (PC)
<b>BROADLEAF WEEDS</b>		
Amaranth, Palmer	<i>Amaranthus palmeri</i>	C
Amaranth, Powell	<i>Amaranthus powellii</i>	C
Beggarweed, Florida	<i>Desmodium tortuosum</i>	PC
Carpetweed	<i>Mollugo verticillata</i>	C
Eclipta	<i>Eclipta prostrata</i>	PC
Galinsoga, hairy	<i>Galinsoga quadriradiata</i>	C
Galinsoga, smallflower	<i>Galinsoga parviflora</i>	C
Nightshade, Eastern black	<i>Solanum ptychanthum</i>	C
Nightshade, hairy	<i>Solanum physalifolium</i>	PC
Pigweed, prostrate	<i>Amaranthus blitoides</i>	C
Pigweed, redroot	<i>Amaranthus retroflexus</i>	C
Pigweed, smooth	<i>Amaranthus hybridus</i>	C
Pigweed, tumble	<i>Amaranthus albus</i>	C
Purslane, common	<i>Portulaca oleracea</i>	PC
Pusley, Florida	<i>Richardia scabra</i>	C
Spiderwort, tropical	<i>Commelina benghalensis</i>	C
Waterhemp, common	<i>Amaranthus rudis</i>	C
Waterhemp, tall	<i>Amaranthus tuberculatus</i>	C
<b>SEDGES</b>		
Nutsedge, yellow	<i>Cyperus esculentus</i>	C

<sup>1</sup> Refer to the **CORN** section of this label for additional information.

**ROTATIONAL CROPS SECTION**

**Replanted Crop Directions**

This section covers replant crops that may be planted following a lost crop that has had an application of MOCCASIN II PLUS Herbicide.

If a crop treated with MOCCASIN II PLUS Herbicide is lost, any crop on this label, or on a supplemental MOCCASIN II PLUS Herbicide label, may be replanted immediately provided that the rate of MOCCASIN II PLUS Herbicide applied to the previous crop was not greater than the labeled rate for the crop to be replanted. If the first application was banded and the replant crop is planted in the center of the untreated bands, a second banded treatment may be applied at the rate for the use pattern for the replant crop, provided the application does not overlap the first application band.

**Rotational Crop Directions**

Any food or feed crops not listed below or in the **Additional Rotational Crop Options** table must NOT be planted within 12 months of the last application of this product.

Crop(s)	Replant Interval
Tobacco, Buckwheat, Rice	The spring following the last treatment
Barley, Oats, Rye, Wheat	4.5 months
Alfalfa	4.0 months
Clover	9.0 months

In order to make a replant application of this product, the previous crop must have had applied 2.0 pounds active ingredient, or less, of *S-metolachlor* (2 pints or less) per acre, and the previous crop must not have had a lay-by or other post-emergence application of this product.

**Note:** If rotating to a crop listed in one of the crop groupings below within 60 days of last application, injury to the new crop may result. Also, if the amount of this product applied to the previous crop was more than listed below, the crop must not be planted until the following spring.

### Additional Rotational Crop Options

Note that not all crops within each group are specifically listed. The plant-back interval applies to all the crops in the specific EPA crop group or subgroup listed.

Amount of MOCCASIN II PLUS Herbicide Applied Previously (Pints per Acre)	Crops that may be Planted 60-days after the Last Application of MOCCASIN II PLUS Herbicide
≤ 1.33 pts/A	<p><b>Crop Subgroup 1B: Root Vegetables (except sugar beet, except carrot)</b> Beet, garden; burdock, edible; celeriac; chervil, turnip-rooted; chicory; ginseng; horseradish; parsnip; radish; radish, oriental; rutabaga; salsify; skirret; turnip</p> <p><b>Crop Subgroup 3-07B: Onion, green</b> Chive leek, onion, Beltsville bunching; onion, fresh; onion, green; onion, Welsh; Shallot</p> <p><b>Crop Subgroup 4-16B: Brassica, leafy greens</b> Bok choy; broccoli, Chinese; cabbage, Chinese (napa); collards; kale; greens, mustard; greens, turnip</p> <p><b>Crop Group 9: Vegetable, cucurbit</b> Cantaloupe; citron melon; cucumber; gourd; muskmelon; pumpkin; squash, summer; squash, winter; watermelon</p> <p>Carrot; leaf lettuce; sesame; strawberry; Swiss chard</p>
≤ 1.67 pts/A	<p><b>Crop Group 8-10: Fruiting Vegetables (except Tabasco pepper)</b> Eggplant; ground cherry (<i>Physalis</i> spp.); okra, pepino; peppers, bell, chili, cooking, pimento &amp; sweet; tomatillo; tomato</p>
≤ 2.0 pts/A	<p><b>Crop Subgroup 1C: Tuberous and Corm Vegetables</b> Arracacha; arrowroot; artichoke (Jerusalem, Chinese); Cassava (bitter, sweet); chayote (root); chufa; dasheen (taro); ginger; potato; potato, sweet; taniar; turmeric; yam bean; yam, true</p> <p><b>Crop Group 3-07A: Onion, Bulb</b> Garlic, bulb, garlic, great-headed; onion, dry bulb; shallot</p> <p><b>Crop Subgroup 22A: Stalk and Stem Vegetable (except Kohlrabi)</b> Agave; asparagus; celtuce; fennel, Florence; fern, edible; kale, sea</p> <p><b>Crop Subgroup 22B: Leaf Petiole Vegetable</b> Cardoon; celery, Chinese; celery; rhubarb</p> <p><b>Crop Subgroup 5-16: Brassica Head and Stem Vegetable</b> Broccoli; Brussel sprouts; cabbage; cabbage, Chinese; cauliflower</p>

**NOTE:** A second application of *S-metolachlor* to the crops in the above table must NOT be made within 60 days of the original application.

## APPLICATION PROCEDURES

### APPLICATION TIMING

MOCCASIN II PLUS Herbicide alone or in some tank mixtures with other labeled herbicides may be applied for weed control in certain crops at various times. Refer to the given crop section of the label to determine if application timings listed below are indicated.

Application Method	Instructions
<b>Preplant Surface-Applied</b>	Refer to individual crop section on this label to determine if early preplant surface application is indicated. For minimum-tillage or no-tillage systems only, MOCCASIN II PLUS Herbicide alone and some MOCCASIN II PLUS Herbicide tank mixtures may be applied up to 45 days before planting certain crops. For treatments made 30 - 45 days before planting, use only split applications, with 2/3 the labeled broadcast rate for the crop and soil texture applied initially and the remaining 1/3 at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (for example, products containing glyphosate or paraquat). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.
<b>Preplant Incorporated</b>	Apply MOCCASIN II PLUS Herbicide to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate MOCCASIN II PLUS Herbicide after bed formation, unless specified otherwise.
<b>Preemergence</b>	Apply MOCCASIN II PLUS Herbicide during planting (behind the planter) or after planting, but before weeds or crops emerge.
<b>Postemergence</b>	MOCCASIN II PLUS Herbicide will not control emerged weeds so apply only to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Refer to the individual crop section of this label if a postemergence application is indicated.

## SPECIAL APPLICATION PROCEDURES

### CA Only (Beans, Peas, and Lentils; Corn; Safflowers): Preplant Incorporated:

Broadcast MOCCASIN II PLUS Herbicide alone or with tank mix partners listed on this label to the soil and thoroughly incorporate with a disk or similar implement set to till 4 - 6 inches deep. For more thorough incorporation, till the soil in 2 different directions (cross-till). Crops may be planted on flat surface or on beds. Use caution when forming the beds that only soil from the MOCCASIN II PLUS Herbicide treated zone is used (i.e., do not bring untreated soil to soil surface). If the application is made to preformed beds, incorporate MOCCASIN II PLUS Herbicide with a tillage implement set to till 2 - 4 inches deep. Take care during tilling to keep the tilled (MOCCASIN II PLUS Herbicide-treated) soil on the beds.

**Preemergence:** Apply MOCCASIN II PLUS Herbicide after planting. Water with sprinkler or flood irrigation within 7 - 10 days.

**Fall Application for Spring Weed Control (Only in IA, MN, ND, SD, WI, and portions of NE and IL - See specific instructions in the BEANS, PEAS, AND LENTILS; CORN; and SOYBEANS sections of this label for timing of application and other information):** Do not apply to frozen ground. Use on medium and fine soils with greater than 2.5% organic matter that will be planted to corn or soybeans the next spring. Ground may be tilled before or after application. Do not incorporate deeper than 2 to 3 inches if tilled after treatment.

### Restriction

- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for the specific crop, or illegal residues may result.

**Fall Application for Italian Ryegrass Control (Corn, Cotton, Grain and Forage Sorghum, and Soybean Only - See specific instructions in the CORN, COTTON, GRAIN OR FORAGE SORGHUM, and SOYBEAN sections of this label for timing of application and other information):** MOCCASIN II PLUS Herbicide may be applied in the fall (September 1 - December 1) for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). A tillage operation may precede the application. Do not incorporate deeper than 2 - 3 inches if tillage follows the application of MOCCASIN II PLUS Herbicide.

### Restrictions

- **DO NOT** apply MOCCASIN II PLUS Herbicide to frozen ground.
- All crops on the MOCCASIN II PLUS Herbicide label may be planted the following spring after application.
- If a spring application is made, the combined total amount of MOCCASIN II PLUS Herbicide applied in the fall plus the spring must not exceed the maximum seasonal S-metolachlor rate for the specific crop planted, or illegal residues may result.
- Refer to the crop sections on this label for specific directions.

**Ground Application:** Apply MOCCASIN II PLUS Herbicide alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For MOCCASIN II PLUS Herbicide tank mixtures with wettable powder or dry flowable formulations, use screens and strainers no finer than 50-mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

For information on applying in lower volumes of carrier, see **LOW CARRIER APPLICATION** section.

For application by air or through center pivot systems, see **SPRAY DRIFT MANAGEMENT** sections.

For information on impregnating dry fertilizer, see **Dry Bulk Granular Fertilizer** section.

For information on application using variable-rate technologies, see **Variable-Rate Application** section.

## SPRAY EQUIPMENT

### LOW CARRIER APPLICATION

#### For Broadcast Ground Application Only

Use sprayers, such as Ag-Chem RoGator®, Hagie, John Deere Hi-Cycle™, Melroe Spra-Coupe, Tyler Patriot™, or Willmar Air Ride®, that provide accurate and uniform application. **Only water may be used as a carrier.** Screens in suction and in-line strainers should be 50-mesh. Manufacturers may require that tip screens as fine as 100-mesh be used with some nozzles. Use a pump with capacity to: (1) maintain up to 35 - 40 psi at the nozzles, and (2) provide sufficient agitation in tank to keep mixture in suspension. Use a minimum of 5.0 gallons of spray mixture per acre. Maximum sprayer speed is 15 mph. Rinse sprayer thoroughly with clean water immediately after each use.

#### Precautions

- Use low pressure nozzles to reduce drift and increase application accuracy.
- Take care when using automatic rate controlling devices to spray the material within the rated working pressure and flow ranges of the nozzles selected.
- Use nozzle screens when specified by the manufacturer.
- Place all nozzles on 20-inch centers, except flooding types which should be placed on 40-inch centers. When Flat Fan-type nozzles are used, angles of 80° or 110° are recommended.
- Always read and follow the manufacturer's directions for optimum setup and performance of their nozzles or tips.

### AERIAL APPLICATION

Apply MOCCASIN II PLUS Herbicide in water alone or in tank mixtures metribuzin in a minimum total volume of 2.0 gals/A by aircraft. MOCCASIN II PLUS Herbicide may be applied by air in combination with atrazine, linuron, benefin, pendimethalin, trifluralin, or metribuzin. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. Apply at a maximum height of 10 ft, using low-drift nozzles at a maximum pressure of 40 psi, and only when wind speed does not exceed 10 mph. To ensure that spray will not adversely affect adjacent sensitive nontarget plants, apply MOCCASIN II PLUS Herbicide alone or MOCCASIN II PLUS Herbicide + atrazine by aircraft at a minimum upwind distance of 400 ft from sensitive plants, or apply MOCCASIN II PLUS Herbicide + linuron or metribuzin at a minimum upwind distance of 300 ft from sensitive plants.

#### Sensitive Areas

The pesticide must only be applied when the potential for drift to adjacent sensitive areas (e.g., residential areas, bodies of water, known habitat for threatened or endangered species, nontarget crops) is minimal (e.g., when wind is blowing away from the sensitive areas).

Avoid application to humans or animals. Flagmen and loaders must avoid inhalation of spray mist and prolonged contact with skin.

### CENTER PIVOT IRRIGATION APPLICATION

MOCCASIN II PLUS Herbicide alone or in tank mixture with other herbicides on this label, which are labeled for center pivot application, may be applied in irrigation water preemergence (after planting, but before weeds or crop emerge) at rates indicated on this label. MOCCASIN II PLUS Herbicide also may be applied postemergence to the crop and preemergence to weeds in crops where postemergence applications are allowed on this label. Follow all restrictions (height, timing, rate, etc.) to avoid illegal residues. Apply this product only through a center pivot irrigation system. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from nonuniform distribution of treated water. If you have questions about calibration, contact State Extension specialists, equipment manufacturers, or other experts. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

## Restrictions

- **DO NOT** apply this product through any other type of irrigation system.
- **DO NOT** connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system, unless the pesticide label-prescribed safety devices for public water systems are in place.

## Operating Instructions

1. The system must contain a functional check-valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water-source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check-valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump or piston pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.
8. Prepare a mixture with a minimum of 1 part water to 1 part herbicide(s) and inject this mixture into the center pivot system. Injecting a larger volume of a more dilute mixture per hour will usually provide more accurate calibration of metering equipment. Maintain sufficient agitation to keep the herbicide in suspension.
9. Meter into irrigation water during entire period of water application.
10. Apply in 1/2 - 1 inch of water. Use the lower water volume (1/2 inch) on *coarse-textured soils* and the higher volume (1 inch) on *fine-textured soils*. More than 1 inch of water at application may reduce weed control by moving the herbicide below the effective zone in the soil.

**Precaution for center pivot applications:** Where sprinkler distribution patterns do not overlap sufficiently, unacceptable weed control may result. Where sprinkler distribution patterns overlap excessively, crop injury may result.

## DRY BULK GRANULAR FERTILIZERS

Dry bulk granular fertilizers may be impregnated or coated with MOCCASIN II PLUS Herbicide alone or selected MOCCASIN II PLUS Herbicide tank mixtures provided that they are registered for preplant incorporated or preplant surface applications which are used to control weeds in crops on the MOCCASIN II PLUS Herbicide label; and that they are not prohibited from use on dry bulk granular fertilizers.

When applying MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide mixtures with dry bulk granular fertilizers, follow all directions for use, restrictions, and precautions on the respective product labels regarding target crops, rates, soil texture, application methods (including timing of application), and rotational crops restrictions.

Compliance with individual state regulations relating to dry bulk granular fertilizer blending, registration, labeling, and application is the responsibility of the individual and/or company selling the herbicide/fertilizer mixture.

Prepare the herbicide/fertilizer mixtures by using any commonly used dry bulk fertilizer blender (such as closed drum, belt, ribbon). Nozzles used to spray MOCCASIN II PLUS Herbicide and MOCCASIN II PLUS Herbicide mixtures onto the fertilizer must be placed to provide uniform spray coverage. Take care to aim the spray directly onto the fertilizer only and to avoid spraying the walls of the blender.

If the herbicide/fertilizer mixture is too wet, add a highly absorptive material such as Agsorb® or Celatom MP-79, or similar granular clay or diatomaceous earth materials, to obtain a dry, free-flowing mixture. Add absorptive materials only after the herbicide has been thoroughly blended into the fertilizer mixture. Best application results will be obtained by using a granule of 6/30 particle size or of a size similar to that of the fertilizer material being used. Generally, less than 2% by weight of absorptive material will be needed. Avoid using more than 5% absorptive material by weight.

Amounts of MOCCASIN II PLUS Herbicide, atrazine, atrazine + simazine, isoxaflutole, simazine, metribuzin, or ethalfluralin can be calculated by using the following formula:

$$\frac{2,000}{\text{lbs of fertilizer per acre}} \times \text{pts/A of liquid or flowable product} = \text{pts of liquid or flowable product per ton of fertilizer}$$
$$\frac{2,000}{\text{lbs of fertilizer per acre}} \times \text{lbs/A of dry product} = \text{lbs of dry product per ton of fertilizer}$$

**Pneumatic (Compressed Air) Application (MOCCASIN II PLUS Herbicide Alone):** High humidity, high urea concentrations, low fertilizer use rates, and dusty fertilizer may cause fertilizer mixture to build up or plug the distributor head, air tubes, or nozzle deflector plates. To minimize buildup, premix MOCCASIN II PLUS Herbicide with Exxon Aromatic 200 at a rate of 1.0 - 4.0 pts/gals of MOCCASIN II PLUS Herbicide. Aromatic 200 is a noncombustible/nonflammable petroleum product. Aromatic 200 may be used in either a fertilizer blender or through direct injection systems. Do not use drying agents when using Aromatic 200.

## Precautions

- Use mixtures of MOCCASIN II PLUS Herbicide and Aromatic 200 on dry fertilizer only. Poor results or crop injury may result if these mixtures are used in water or liquid fertilizer solutions for spraying applications.
- When impregnating MOCCASIN II PLUS Herbicide in a blender before application, a drier mixture can be attained by substituting a drying agent for Aromatic 200. The use of Agsorb FG or drying agents of 6/30 particle size are recommended.

## Restrictions

- To avoid potential for explosion, **DO NOT** impregnate MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide mixtures on ammonium nitrate, potassium nitrate, or sodium nitrate, either alone or in blends with other fertilizers.
- **DO NOT** use MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide mixtures on straight limestone, since absorption will not be achieved. Fertilizer blends containing limestone can be impregnated.
- **DO NOT** use drying agents with On-The-Go impregnation equipment.

## Application

Apply 200 - 700 lbs of the herbicide/fertilizer mixture per acre. For best results, apply the mixture uniformly to the soil with properly calibrated equipment immediately after blending. Uniform application of the herbicide/fertilizer mixture is essential to prevent possible crop injury. Nonuniform application may also result in unsatisfactory weed control. In areas where conventional tillage is practiced, a shallow incorporation of the mixture into the soil may improve weed control. On fine- or medium-textured soils in areas where soil incorporation is not planned, i.e., reduced tillage situations or in some conventional till situations, make applications approximately 30 days before planting to allow moisture to move the herbicide/fertilizer mixture into the soil. On coarse-textured soils, make applications approximately 14 days prior to planting.

## Precaution

- To avoid crop injury, do not use the herbicide/fertilizer mixture on crops where bedding occurs.



## MIXING INSTRUCTIONS

**MOCCASIN II PLUS Herbicide Alone:** Mix MOCCASIN II PLUS Herbicide with water or fluid fertilizer and apply as a spray. Fill the spray tank 1/2 - 3/4 full with water or fluid fertilizer, add the proper amount of MOCCASIN II PLUS Herbicide, then add the rest of the water or fluid fertilizer. Provide sufficient agitation during mixing and application to maintain a uniform emulsion.

### Tank Mixtures – follow this order:

Fill the spray tank 1/4 full with water, and start agitation;

Next add 2,4-D, atrazine, benefin, isoxaflutole, dicamba + atrazine, bentazon, 2,4-DB, metribuzin + chlorimuron-ethyl, prometryn, clomazone, fluometuron, EPTC, glufosinate, linuron, MSMA, simazine, pendimethalin, imazethapyr, imazaquin, metribuzin, ethalfuralin, or trifluralin, and allow it to become dispersed;

Add MOCCASIN II PLUS Herbicide;

Then add paraquat, glyphosate + 2,4-D, or glyphosate if these products are being used;

And finally add the rest of the water.

For tank mixtures with atrazine, isoxaflutole, dicamba + atrazine, metribuzin + chlorimuron-ethyl, prometryn, clomazone, fluometuron\*, EPTC, linuron, simazine, pendimethalin\*, imazethapyr, imazaquin, metribuzin, ethalfuralin, or trifluralin, fluid fertilizers may replace all or part of the water as carrier, except in the atrazine postemergence and the dicamba + atrazine postemergence tank mixes. For tank mixtures with atrazine, see additional mixing instructions on the atrazine label.

For each mixture, check compatibility with fluid fertilizer, as described below, before mixing in spray tank. For all tank mixtures, agitate during mixing and application to maintain a uniform suspension.

\* See **Special Mixing Instructions** for tank mixtures with fluometuron and with atrazine or simazine + pendimethalin under the appropriate tank mixture section.

For directions on how to conduct a compatibility test, see the **COMPATIBILITY TEST** section.

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### COMPATIBILITY TEST

A jar test is beneficial before tank mixing to ensure compatibility of MOCCASIN II PLUS Herbicide with other pesticides. The following test assumes a spray volume of 25 gals/A. For other spray volumes, make appropriate changes in the ingredients.

**Precautions:** Nitrogen solutions or complete fluid fertilizers may replace all or part of the water in the spray. Because liquid fertilizers vary, even within the same analysis, **always check compatibility with pesticide(s) before use.** Incompatibility of tank mixtures is more common with suspensions of fertilizer and pesticides.

### Test Procedure

1. Add 1.0 pt of carrier (fertilizer or water) to each of 2 one qt jars with tight lids. **Restrictions:** Use the same source of water that will be used for the tank mix and conduct the test at the temperature the tank mix will be applied.
2. To one of the jars, add 1/4 tsp or 1.2 milliliters of a compatibility agent approved for this use, such as Compex® or Unite® (1/4 tsp is equivalent to 2.0 pts/100 gals spray). Shake or stir gently to mix.
3. To both jars, add the appropriate amount of pesticide(s) in their relative proportions based on label rates. If more than one pesticide is used, add them separately with dry pesticides first, flowables next, and emulsifiable concentrates last. After each addition, shake or stir gently to thoroughly mix.

4. After adding all ingredients, put lids on and tighten, and invert each jar ten times to mix. Let the mixtures stand 15 - 30 minutes and then look for separation, large flakes, precipitates, gels, heavy oily film on the jar, or other signs of incompatibility. Determine if the compatibility agent is needed in the spray mixture by comparing the two jars. If either mixture separates, but can be remixed readily, the mixture can be sprayed as long as good agitation is used. If the mixtures are incompatible, test the following methods of improving compatibility: (a) Slurry the dry pesticide(s) in water before addition, or (b) add 1/2 the compatibility agent to the fertilizer or water and the other 1/2 to the emulsifiable concentrate or flowable pesticide before addition to the mixture. If incompatibility is still observed, do not use the mixture.

5. After compatibility testing is complete, dispose of any pesticide wastes in accordance with the **STORAGE AND DISPOSAL** section in this label.

**Example Tank Mix Partner Products** To tank mix, fill the solution tank 1/4 full with water and commence agitation. If the product to be tank mixed is underlined in the table, add it first and allow to disperse. Next add MOCCASIN II PLUS Herbicide, followed by any products NOT underlined in the table. Then add the remaining amount of water.

Active Ingredient	Brand Name(s) <i>Alternative brands may be used</i>	EPA Registration Number(s) <i>Alternative brands may be used</i>
2,4-D	Amine 4	42750-14-55467, 55467-14, 71368-1-55467
	Lo-Vol 4	228-139-55467
	Lo-Vol 6	42750-20-55467, 71368-11-55467
2,4-DB	Butoxone	2749-126 & 2749-516
	Butyrac	42750-39 & 42750-38
Acifluorfen	STORM	70506-59
	ULTRA BLAZER	70506-60
Atrazine	Atrazine 4L	55467-13, 100-497-55467, 19713-11-55467
	Atrazine 90DF	100-585-55467, 35915-3-55467
	AAtrex	100-497 & 100-585
Benfluralin	Balan 2.5G	8378-35
	Balan Dry Flowable	34704-746
Bentazon	Basagran	7969-45-(multiple), 7969-112, 70506-434
	BROADLOOM	70506-306
Chlorimuron	Classic	5481-681
Clethodim	Select	59639-3
Clopyralid	Stinger	62719-73
Cloransulam	FirstRate	5481-676
Desmedipham	Betanex	254-620
Dicamba	Detonate	7969-137-55467, 42750-209-55467
	Banvel	55467-38
Dicamba + Atrazine	Marksman	7969-136

(continued)

Active Ingredient	Brand Name(s) <i>Alternative brands may be used</i>	EPA Registration Number(s) <i>Alternative brands may be used</i>
Ethalfuralin	Sonalan	10163-355 & 10163-356
EPTC	Eptam	10163-281 & 10163-283
Flometuron	Cotoran 4L	66222-181
Fomesafen	Flexstar	100-1101
	Flexstar GT	100-1325
	Reflex	100-993
Fluazifop	Fusilade	100-1070
Fluazifop + Fenoxaprop	Fusion	100-1059
Flumetsulam	Python	5481-677
Glufosinate	Autonomy	7969-448-55467
	INTERLINE	70506-310
	Liberty	264-829 & 7969-448
Glyphosate	Roundup	524-549-(multiple)
	Buccaneer	55467-9, 55467-10, 55467-15
Glyphosate + 2,4-D	Landmaster BW	42750-62
Imazethapyr	Pursuit	241-310
Isoxaflutole	Balance Pro	264-600
Linuron	Linuron DF	19713-251
	Linex	61842-21
	Lorox DF	61842-23
Metribuzin	Lexone	352-382
	Sencor	432-1469 & 264-738
	TRICOR	70506-68 & 70506-103
Metribuzin + Chlorimuron	Canopy	352-444
Metribuzin + Sulfentrazone	Authority MTZ	279-3340
s-Metolachlor + Fomesafen	Prefix	100-1268
MSMA	MSMA 6 Plus	19713-42
	MSMA 6.6	19713-41
	Target 6 Plus	42519-3
	Target 6.6	42519-1
Nicosulfuron	Accent XP	352-817
Paraquat	Gramoxone	100-1431 & 100-1652
Pendimethalin	ACUMEN	241-337-55467, 70506-318
	Prowl	241-337 & 241-418
	SATELLITE 3.3	70506-318
	SATELLITE HYDROCAP	70506-230

(continued)

Active Ingredient	Brand Name(s) <i>Alternative brands may be used</i>	EPA Registration Number(s) <i>Alternative brands may be used</i>
Prometryn	Caparol	100-620
Quizalofop	Assure II	5481-646
Rimisulfuron-methyl	Beacon	100-705
Saflufenacil	Sharpen	7969-278
Saflufenacil + Dimethenamide-P	Verdict	7969-279
Sethoxydim	Poast	7969-58-(multiple)
Simazine	Princep	100-526 & 100-603
Sulfentrazone + Cloransulam	Sonic	62719-680
	Gauntlet	279-3246
Sulfentrazone + Chlorimuron	Authority Maxx	279-9560
Sulfosate	Touchdown	100-1117
Trifluralin	Treflan	34704-853-(multiple)
Trisulfuron	Upbeet	279-9584

## CROP USE DIRECTIONS

### CORN (ALL TYPES) MOCCASIN II PLUS HERBICIDE ALONE

Apply either preplant surface, preplant incorporated, preemergence, postemergence, or lay-by, using the rate specified below.

#### PREPLANT SURFACE-APPLIED

Refer to instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**.

#### Fall Application for Spring Weed Control:

Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts/A on *medium-textured* and 2.0 pts/A on *fine-textured soils*. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not incorporate deeper than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

**Fall Application for Italian Ryegrass Control:** MOCCASIN II PLUS Herbicide may be applied for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*). Apply MOCCASIN II PLUS Herbicide at 1.33 - 1.67 pts/A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower MOCCASIN II PLUS Herbicide rate for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of MOCCASIN II PLUS Herbicide. For fall applications after emergence of glyphosate-resistant Italian ryegrass, paraquat can be tank mixed with MOCCASIN II PLUS Herbicide to control emerged ryegrass. Refer to the paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with MOCCASIN II PLUS Herbicide for control or improved control of other weeds present at the time of application.

### Restrictions

- **DO NOT** apply MOCCASIN II PLUS Herbicide to frozen ground.
- **DO NOT** apply more than 3.9 pts/A (3.77 lbs ai) per year (depending on soil texture) through any combination of applications.

**Fall Application for Control or Suppression of Yellow Nutsedge (ID, OR, and WA only):** For preemergent control or suppression of yellow nutsedge the following spring, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide in the fall after the harvest of the previous crop but before freeze-up. Fall applications of MOCCASIN II PLUS Herbicide can be surface-applied or incorporated.

### Restrictions

- **DO NOT** make more than one fall application per crop.
- **DO NOT** apply more than 1.33 pts/A in a single fall preplant application.
- **DO NOT** apply to frozen ground.
- **DO NOT** apply more than 3.9 pts/A (3.77 lbs ai) per year (depending on soil texture) through any combination of applications.

## EARLY PREPLANT APPLICATIONS

Use on medium- and fine-textured soils with minimum-tillage or no-tillage systems in CO, IA, IL, IN, KS, KY, MN, MO, MT, ND, NE, SD, TN, WI, and WV. Apply 2/3 the labeled rate of MOCCASIN II PLUS Herbicide (1.67 pts/A on *medium soils* and 2.0 pts/A on *fine soils*) as a split treatment 30 - 45 days before planting and the remainder at planting. Applications made less than 30 days prior to planting may be as either a split or single treatment. Apply 1.33 pts/A on *coarse soils* not more than 2 weeks before planting.

### Restriction

- **DO NOT** apply more than 3.9 pts/A (3.77 lbs ai) per year (depending on soil texture) through any combination of applications.

On medium- and fine-textured soils with minimum- or no-tillage systems in CT, DE, MA, MD, ME, MI, NH, NY, OH, PA, RI, VA, VT, and WV, preplant surface applications may be applied following the directions for use above. If the amount of rainfall results in unsatisfactory length of weed control following the earlier treatment, a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide may be used, i.e., atrazine, primisulfuron-methyl, atrazine + s-metolachlor, nicosulfuron, dicamba + atrazine, bentazon, bromoxynil, or 2,4-D. If the postemergence treatment includes the herbicide used preplant surface-applied, do not exceed the total labeled rate for corn on a given soil texture. For extended residual or control of heavy weed infestations, up to 2.6 pts/A is allowed. Observe all directions for use, precautions, and limitations on the label of the postemergent herbicide.

## PREPLANT INCORPORATED OR PREEMERGENCE

Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**. On *coarse soils*, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.33 pts/A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.67 - 2.0 pts/A if organic matter content is 3% or greater. For extended residual or control of heavy weed infestations, up to 2.6 pts/A is allowed.

## POSTEMERGENCE OR LAY-BY

To extend the duration of weed control in corn, a maximum rate of 2.0 pts/A of MOCCASIN II PLUS Herbicide may be applied after corn emergence until the corn plants reach 40 inches in height, following any preplant surface-applied, preplant incorporated, or preemergence herbicide application, including MOCCASIN II PLUS Herbicide. For best results, apply to soil free of emerged weeds and directed toward the base of corn plants taller than 5 inches. The total MOCCASIN II PLUS Herbicide rate applied on corn during any one year must not exceed 3.9 pts/A, depending on soil texture.

### Restrictions for all applications to corn

- **DO NOT** graze or feed forage from treated areas for 30 days following application.
- PHI (Pre-Harvest Interval): **DO NOT** harvest sweet corn ears from treated areas for 30 days following application.

## PROBLEM WEED CONTROL DIRECTIONS

### Shattercane, Wild Proso Millet, Woolly Cupgrass, and Eclipta – Partial

**Control:** For more consistent partial control of these weeds, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide preplant incorporated followed by 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge. Use the 1.33 pts/A rate when a heavy infestation is expected. A shallow cultivation may be needed to control any late emerging weeds.

**Woolly Cupgrass and Wild Proso Millet Control Program:** For control of these species, use the following 3-step program: (1) Apply MOCCASIN II PLUS Herbicide early preplant, preplant incorporated, or preemergence at 1.67 pts/A on *medium soils* and 2.0 pts/A on *fine-textured soils*, up to the maximum label rate. Lightly incorporate with a rotary hoe if rainfall does not occur within 5 - 7 days; (2) Apply a postemergence tank mix of primisulfuron-methyl at 0.38 oz/A at 1 packet per 4 acres plus nicosulfuron at 0.33 oz/A plus 1.0 qt of crop oil concentrate plus 1.0 gal/A of 28% nitrogen, or the equivalent amount of ammonium sulfate, when grasses are 2 - 3 inches tall and the corn is at least 4 inches tall; and (3) Cultivate 14 - 21 days after the postemergence application.

In the event of escape of annual weeds following a preplant surface, preplant incorporated, or preemergence treatment of MOCCASIN II PLUS Herbicide, follow with a postemergence application of an appropriately labeled broadleaf and/or grass weed herbicide, i.e., atrazine, primisulfuron-methyl, atrazine + s-metolachlor, nicosulfuron, dicamba + atrazine, bentazon, bromoxynil, or 2,4-D. If the postemergence treatment includes the herbicide used in the earlier treatment, do not exceed the total labeled rate for corn on a given soil texture.

### Restrictions

- **DO NOT** apply more than the labeled application rate for a given soil texture per year through any combination of applications.
- In corn, use up to 2.5 pts/A of MOCCASIN II PLUS Herbicide preplant surface, preplant incorporated, or preemergence treatment on soils having an organic matter content between 6% and 20%.
- Bromoxynil may be applied postemergence alone or in tank mix combination with atrazine. **DO NOT** exceed the allowed amount of atrazine in tank mix combination with bromoxynil postemergence. Refer to the atrazine, and bromoxynil labels for specific rates and precautions.
- **DO NOT** use MOCCASIN II PLUS Herbicide on peat or muck soils.

## CORN

### MOCCASIN II PLUS HERBICIDE COMBINATIONS

MOCCASIN II PLUS Herbicide in any tank mixture for corn may be applied in water or fluid fertilizer before corn emerges. Use only water as a carrier when MOCCASIN II PLUS Herbicide is applied after corn emergence.

### Restrictions

- **DO NOT** graze or feed forage from treated areas for 30 days following application.
- Pre-Harvest Interval (PHI): **DO NOT** harvest sweet corn ears from treated areas for 30 days following application.

### IMPORTANT: for tank mixtures with atrazine:

If applying MOCCASIN II PLUS Herbicide in tank mixture with atrazine, follow all restrictions and rate limitations on the atrazine label.

Some states may have established rate limitations for atrazine within specific geographical areas. Consult your state lead pesticide control agency for more information. It is a violation of this label to deviate from state use regulations.

**Table 2**

**MOCCASIN II PLUS Herbicide Tank Mixtures for Corn – Additional Weeds Controlled and Special Instructions**

	MOCCASIN II PLUS Herbicide + atrazine and/or simazine (Preplant Surface, PPI, PRE)	MOCCASIN II PLUS Herbicide + atrazine (Post)	MOCCASIN II PLUS + dicamba + atrazine (Field Corn)	MOCCASIN II PLUS + atrazine + linuron	MOCCASIN II PLUS Herbicide + atrazine or simazine + pendimethalin	MOCCASIN II PLUS Herbicide + atrazine + dicamba	MOCCASIN II PLUS + isoxaflutole
<b>Special Mixing Instructions</b>					1		
<b>Comments</b>	2,3,4,5,7,8	2,3,4,5		2,3,4,5,6	2,3,4,5	7	2,3,7
Browntop panicum	X			X	X		X
Cocklebur	X	0	0	X	X		0-X
Common purslane	X			X	X	X	X
Hairy nightshade	X			X	X		X
Jimsonweed		X	0			X	X
Kochia		X				X	X
Lambsquarters	X	X	X	X	X	X	X
Morningglory	X	0	0	X	X		X
Mustard		X				X	X
Pigweed				X	X	X	X
Prickly sida		X					
Ragweed	X	X	X	X	X	X	X
Smartweed	X	X	X	X	X	X	X
Velvetleaf	X	X	0	X	X	0-X	0-X

X = control; 0 = partial control; 0-X = partial to full control depending on ratio of products used or on weed population

**Comments**

- Special Mixing Instructions for MOCCASIN II PLUS Herbicide + atrazine or simazine and pendimethalin.
  - Fill the spray tank 1/4 full with water or fluid fertilizer and start agitation.
  - To aid compatibility, add a compatibility agent, such as Unite or X-77®, at 4.0 pts/100 gals of spray mixture.
  - Then add the atrazine or simazine and allow it to become dispersed.
  - Then add MOCCASIN II PLUS Herbicide and pendimethalin.
  - Add the rest of the water.
- Although a single formulation for atrazine or simazine is listed in the rate tables, other formulations may be substituted.
- Restriction: DO NOT** apply more than the labeled rate of atrazine per acre per year.
- In Minimum-Tillage and No-Tillage systems, mix with paraquat for control of most emerged annual weeds and suppression of perennial weeds; or with glyphosate + 2,4-D for suppression of emerged field bindweed and control or suppression of annual weeds; or with or glyphosate for control of most emerged annual and perennial weeds.
- Refer to the **CORN – MOCCASIN II PLUS HERBICIDE COMBINATIONS / TANK MIXTURE WITH ATRAZINE; OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA + ATRAZINE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS** sections for specific directions for 2,4-D or dicamba + atrazine burndown combinations in Minimum-Tillage and No-Tillage systems.

MOCCASIN II PLUS Herbicide in any tank mixture for corn may be applied in water or fluid fertilizer, except as noted.

Refer to **CORN (ALL TYPES) – MOCCASIN II PLUS HERBICIDE ALONE** for sequential postemergence treatments if escape weeds develop.

**Restrictions**

- For all applications to corn, **DO NOT** graze or feed forage from treated areas for 30 days following application.
- PHI (Pre-Harvest Interval): **DO NOT** harvest sweet corn ears from treated areas for 30 days following application.
- When applying MOCCASIN II PLUS Herbicide in tank mixture with atrazine, **DO NOT** exceed the allowed amount of atrazine per acre per year.

**TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE PREPLANT SURFACE, PREPLANT INCORPORATED, OR PREEMERGENCE**

In addition to the weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + atrazine or simazine, or MOCCASIN II PLUS Herbicide + atrazine + simazine, applied preplant surface, preplant incorporated, or preemergence, also controls the following weeds: browntop panicum, cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

Apply MOCCASIN II PLUS Herbicide + atrazine or simazine or MOCCASIN II PLUS Herbicide + atrazine + simazine either preplant surface, preplant incorporated, or preemergence.

**Preplant Surface-Applied:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES** and under application instructions for MOCCASIN II PLUS Herbicide alone on corn. Apply the tank mixtures as a split or single treatment as indicated in the **MOCCASIN II PLUS HERBICIDE ALONE / PREPLANT SURFACE-APPLIED** section of the label for corn.

**Preplant Incorporated or Preemergence:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**. Apply MOCCASIN II PLUS Herbicide + atrazine or simazine, or MOCCASIN II PLUS Herbicide + atrazine + simazine, using the rates from **Table 3** and labeled rates of these tank mix partners.

**Restriction**

- **DO NOT** apply more than the labeled rate for a given soil texture per year through any combination of applications.

**Shattercane and Wild Proso Millet – Partial Control**

For more consistent partial control of shattercane or wild proso millet where MOCCASIN II PLUS Herbicide is applied in tank mixture or sequentially with other registered corn herbicides, the following applications may be made:

1. Apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide + the labeled rate of atrazine or simazine preplant incorporated, followed by 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide preemergence. Make the preemergence application during or after planting, but before weeds and corn emerge.
2. Apply MOCCASIN II PLUS Herbicide at 1.33 pts/A alone or in tank mix combination with the labeled rate of atrazine, or simazine, preplant incorporated. Do not exceed the total rate of triazine herbicide labeled in combination with MOCCASIN II PLUS Herbicide for corn grown on a given soil texture. Follow with a post-directed application of ametryn at the labeled rate. Refer to the ametryn label for specific directions for the post-directed application.
3. Apply EPTC or butylate formulation at labeled rates preplant incorporated, followed by a preemergence application of MOCCASIN II PLUS Herbicide at 1.0 - 1.33 pts/A. Do not use EPTC on soils where rapid degradation has been shown to occur. Make the preemergence application during or after planting, but before weeds and corn emerge.

**Note:** When following the application regimes in numbers 1 - 3 above, a shallow cultivation may be needed after the preemergence or postemergence application to help control any late emerging shattercane or wild proso millet plants.

**Restriction**

- **DO NOT** apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide in the preplant incorporated plus preemergence application on soils with less than 6% organic matter, or crop injury may result.

**Table 3**

**MOCCASIN II PLUS Herbicide + atrazine or simazine, or MOCCASIN II PLUS Herbicide + atrazine + simazine, Preplant Incorporated or Preemergence – Corn (All Types)**

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of atrazine and/or simazine.

**Broadcast Rate per Acre**

Soil Texture	Less than 3% Organic Matter	3% Organic Matter or Greater
<b>COARSE</b>	0.8 - 1.0 pt	1.0 pt
<b>MEDIUM</b>	1.0 - 1.33 pts	1.33 pts
<b>FINE</b>	1.33 pts	1.33 - 1.67 pts

Use simazine rather than atrazine when heavy infestations of crabgrass or fall panicum are expected. On soils having between 6% and 20% organic matter, apply MOCCASIN II PLUS Herbicide up to 2.33 pts/A in tank mix combination with atrazine. Refer to the atrazine label for weeds controlled.

When using the tank mixture of MOCCASIN II PLUS Herbicide + atrazine + simazine, use equal rates of each as labeled when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of atrazine + simazine.

For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply the labeled rate atrazine, or the same total amount of atrazine + simazine with 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide.

**TANK MIXTURE WITH ATRAZINE POSTEMERGENCE**

**Weeds Controlled**

- |                            |                 |
|----------------------------|-----------------|
| barnyardgrass (watergrass) | morningglory*   |
| cocklebur*                 | mustard         |
| crabgrass                  | pigweed         |
| crowfootgrass              | prickly sida    |
| fall panicum               | purslane        |
| foxtail, giant             | ragweed         |
| foxtail, green             | smartweed       |
| foxtail, yellow            | velvetleaf      |
| jimsonweed                 | yellow nutsedge |
| kochia                     |                 |
| lambsquarters              |                 |
| * partial control          |                 |

Apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *coarse soils*, 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *medium soils*, or 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *fine soils*. Apply this tank mixture before grass and broadleaf weeds pass the 2-leaf stage and before corn exceeds 5 inches in height. Application to weeds larger than the 2-leaf stage will generally result in unsatisfactory control.

**Lay-by:** Apply to corn plants no taller than 12 inches. Direct applications to corn taller than 5 inches to the base of the corn plants. Applications to corn plants less than 5 inches tall may be made over the top. Occasionally, some corn leaf burn may occur, but this should not affect later growth or yield. Do not apply this postemergence tank mixture in fluid fertilizer, or severe crop injury may occur.

For better control of cocklebur, morningglory, velvetleaf, and yellow nutsedge on *fine-textured soils* above 3% organic matter, apply the labeled rate of atrazine, with 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide.

Tank mixtures of MOCCASIN II PLUS Herbicide + atrazine may be applied following use of any registered preplant surface-applied, preplant incorporated, or preemergence corn herbicide, including MOCCASIN II PLUS Herbicide + atrazine.

## Restriction

- **DO NOT** apply more than 3.9 pts/A (3.77 lbs ai) of MOCCASIN II PLUS Herbicide, or more than the labeled rate of atrazine ai/A during any one year. Refer to the atrazine label for geographic, soil-texture, and rotational restrictions.

### TANK MIXTURE WITH DICAMBA + ATRAZINE

**Preemergence:** Use this tank mixture only on field corn which is flat-planted (no furrows) in CO, IA, IL, IN, KS, MN, NE, OH, SD, and WI.

In addition to the weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + dicamba + atrazine, applied preemergence, also controls lambsquarters, ragweed, and smartweed and will provide partial control of cocklebur, jimsonweed, morningglory, and velvetleaf.

Apply MOCCASIN II PLUS Herbicide + dicamba + atrazine preemergence. Broadcast labeled rate of dicamba + atrazine with 1.33 pts/A of MOCCASIN II PLUS Herbicide on *medium soils*, or with 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide on *fine soils*. Apply this tank mixture to the soil surface at planting or after planting, but before corn emerges. Plant corn at least 1.5 inches deep and apply behind planting equipment, avoiding incorporation by the planter wheel or other seed covering device. Do not incorporate before corn emergence. If it is necessary to rotary hoe to break the soil crust, do not disturb the soil more than 1/2 inch deep.

## Precautions

Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

## Restrictions

- **DO NOT** apply with aircraft.
- **DO NOT** apply on coarse soils or on soils with less than 2.5% organic matter.

**Postemergence for Control of Pigweed (Mid-Atlantic states, including DE, MD, PA, VA, and WV):** Apply 1.0 - 1.5 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of dicamba + atrazine or dicamba by ground equipment when pigweed plants are less than 3 inches tall and before corn is taller than 5 inches in a minimum of 20 gallons of spray per acre. Use the lower rate on coarse-textured and low organic matter soils. Use the higher rate on fine-textured and high organic matter soils.

## Precautions

Avoid drift to sensitive nontarget plants, such as soybeans, during application, or injury may occur.

## Restrictions

- **DO NOT** apply with aircraft.

### TANK MIXTURE WITH ATRAZINE OR SIMAZINE + PENDIMETHALIN FOR PROLONGED CONTROL OF LAMBSQUARTERS AND PIGWEED IN FIELD CORN ONLY (NORTHEAST U.S., INCLUDING MI, IN, KY, AND STATES EAST OF THESE)

For prolonged control of lambsquarters and pigweed, in addition to a broad spectrum of annual broadleaf and grass weeds, MOCCASIN II PLUS Herbicide in tank mix combination with atrazine\* or simazine + pendimethalin may be applied after planting, but before corn or weeds emerge. Apply by ground equipment in a minimum of 10 gallons of water or 20 gallons of liquid fertilizer. Apply by air in a minimum of 5.0 gallons of water. Refer to **Table 3** of this label for rates of MOCCASIN II PLUS Herbicide. Apply tank mix partners as specified on their labels.

\* **DO NOT** apply MOCCASIN II PLUS Herbicide in tank mix combination with atrazine 80W + pendimethalin, as this combination is not compatible. Other atrazine formulations may be used.

**Mixing Instructions:** See **Comment No. 1** following **Table 2**.

Observe all directions for use, restrictions, and limitations on the respective product labels when applying these products in tank mix combination. Refer to the pendimethalin label for replanting instructions in the event of crop loss.

### TANK MIXTURE WITH ATRAZINE OR SIMAZINE, OR ATRAZINE + SIMAZINE WITH PARAQUAT, GLYPHOSATE + 2,4-D, OR GLUFOSINATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, glyphosate + 2,4-D, or glyphosate should be tank mixed with MOCCASIN II PLUS Herbicide + atrazine, MOCCASIN II PLUS Herbicide + simazine, or MOCCASIN II PLUS Herbicide + atrazine + simazine. See **Comment No. 7** following **Table 2**. The MOCCASIN II PLUS Herbicide, MOCCASIN II PLUS Herbicide + atrazine or simazine, or MOCCASIN II PLUS Herbicide + atrazine + simazine portion of the tank mixture provides pre-emergence control of the weeds listed on this label in the tank mixture section for **MOCCASIN II PLUS Herbicide, MOCCASIN II PLUS HERBICIDE + ATRAZINE OR SIMAZINE, or MOCCASIN II PLUS HERBICIDE + ATRAZINE + SIMAZINE / Preplant Surface, Preplant Incorporated, or Preemergence.**

**Application:** Apply before, during, or after planting, but before the corn emerges. Add paraquat, glyphosate + 2,4-D, or glyphosate and apply as directed on the product label.

**Paraquat:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Restrictions:** Do not apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

**Glyphosate + 2,4-D:** See the label for weeds controlled, rates for specific weeds, and other information concerning use.

**Glyphosate:** See the glyphosate label for weeds controlled, rates, and other use directions.

Apply in 20 - 60 gallons of water or fluid fertilizer per acre with ground equipment. Use simazine in preference to atrazine when heavy infestations of crabgrass or fall panicum are expected.

When using the tank mixture of MOCCASIN II PLUS Herbicide + atrazine + simazine, use equal rates of atrazine and simazine as shown when heavy broadleaf weed infestations are expected. When heavy infestations of crabgrass or fall panicum are expected, use a 1:2 ratio of atrazine + simazine instead of the 1:1 ratio given. Refer to **Comment No. 2** following **Table 2** for atrazine 4L and simazine 4L conversions.

For cocklebur, yellow nutsedge, and velvetleaf control on *fine-textured soils* above 3% organic matter, apply the labeled rate of atrazine, or the same total amount of atrazine + simazine, with 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide.

### TANK MIXTURE WITH ATRAZINE; OR ATRAZINE + 2,4-D; OR ATRAZINE + 2,4-D + DICAMBA + ATRAZINE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where corn is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, MOCCASIN II PLUS Herbicide applied in combination with atrazine will kill most emerged small annual weeds. Apply MOCCASIN II PLUS Herbicide + labeled rate of atrazine before, during, or after planting, but before corn emerges, according to the rates in **Table 3**.

Where heavy crop residues exist, add labeled 3.8 lbs ai/gals of 2,4-D amine (such as Weedar 64, Weedar 64A, DMA-4 Herbicide, Weedone® 638 or Formula 40) to the spray tank last and apply in a minimum of 25 gallons of carrier per acre.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds, and therefore, are

recommended instead of water. Add X-77 surfactant at 1.0 - 2.0 qts/100 gals of diluted spray, or another appropriate surfactant at its labeled rate, or add crop oil concentrate plus 28% liquid nitrogen (or equivalent). Apply before weeds are taller than 3 inches. If alfalfa is present, add dicamba + atrazine to the spray mixture at labeled rates and apply before alfalfa is taller than 6 inches.

For fields with existing sod grasses (e.g., bromegrass, orchardgrass, rye, or timothy), when existing weeds exceed 3 inches in height or when very dry conditions exist, add paraquat at the rate indicated on the product label in place of or in addition to 2,4-D, as indicated above. Do not apply paraquat in suspension-type liquid fertilizer. Observe all directions for use, restrictions, and limitations on the respective product labels when applying these products in tank mix combination.

### TANK MIXTURE WITH ATRAZINE + DICAMBA IN CONSERVATION TILLAGE FIELD AND SILAGE CORN

In conservation tillage systems where corn is planted directly into a cover crop or previous crop residue, MOCCASIN II PLUS Herbicide + atrazine + dicamba will kill most emerged small annual weeds. Apply MOCCASIN II PLUS Herbicide + atrazine + dicamba before, during, or after planting, but before corn emergence on *medium* and *fine* soils with greater than 2.5% organic matter. For fields with existing vegetation taller than 3 inches or when very dry conditions exist, add paraquat at its standard rate. MOCCASIN II PLUS Herbicide + atrazine + dicamba may be applied postemergence to corn shorter than 3 inches and before weedy grasses exceed the 2-leaf stage.

As carriers, nitrogen solutions and complete liquid fertilizers, applied before corn emergence, enhance burndown of existing weeds. Do not apply paraquat in suspension-type liquid fertilizer or use on emerged corn.

Refer to the atrazine + dicamba label and follow all directions, limitations, restrictions, and information regarding application and use in corn.

### TANK MIXTURE WITH ISOXAFLUTOLE FIELD CORN ONLY

MOCCASIN II PLUS Herbicide and isoxaflutole have a complementary response and weed control profile which allows various tank mix rate combinations to be considered. The addition of isoxaflutole will improve the control of certain problem weeds, including Texas panicum, woolly cupgrass, and wild proso millet. MOCCASIN II PLUS Herbicide improves both the duration and spectrum of annual grass and small-seeded broadleaf weed control, in particular foxtails (yellow foxtail), witchgrass, and yellow nutsedge.

To reduce the risk of an adverse crop response, the isoxaflutole label does not allow applications to *coarse-textured soils* with less than 1.5% organic matter and warns about applications to all soils with less than 1.5% organic matter or with pH greater than 7.5, as well as applications made to areas in fields with clay knolls, eroded hillsides, and exposed subsoil. MOCCASIN II PLUS Herbicide has no adverse crop response warnings or use restrictions.

Select a rate option for MOCCASIN II PLUS Herbicide plus isoxaflutole by weighing the intensity of problem weed pressure (population presence and density) and your acceptance for risk of an adverse crop response. For example, where Texas panicum, woolly cupgrass, or wild proso millet is a primary target weed, use a tank-mix combination with a higher isoxaflutole rate for the given soil type.

Where the acceptance of an adverse crop response risk is low and/or a more general weed spectrum is targeted (especially yellow foxtail, witchgrass, or yellow nutsedge), use a tank-mix combination with a higher MOCCASIN II PLUS Herbicide rate for the given soil type. Where a target weed is listed as controlled on both product labels, a tank-mix combination option including intermediate rates of both products may be used. Where a target weed is listed as controlled on only one product label, do not apply a rate of that product below what is labeled for that weed on the individual product label, or unacceptable control may result.

Follow all other directions for use, rate limitations, precautions, and restrictions on both the MOCCASIN II PLUS Herbicide and isoxaflutole product labels.

Do not use isoxaflutole on *coarse-textured soils* with less than 1.5% organic matter.

### TANK MIXTURES FOR POSTEMERGENCE SALVAGE WEED CONTROL IN FIELD CORN ONLY

For postemergence control of weeds in specific types of field corn, the MOCCASIN II PLUS Herbicide combinations listed below may be used. Full season weed control from early preplant, preplant incorporated, or preemergence treatments can lead to maximum yield potential under competition-free conditions. However, if control of emerged weeds is needed, a postemergence program listed below can be applied to provide residual control for the remainder of the season.

Follow all label directions, instructions, restrictions, and limitations for each product used. For each tank mixture with MOCCASIN II PLUS Herbicide, apply only to the specific field corn type specified on the tank mix product label.

#### Restrictions

- **DO NOT** use fluid fertilizer with these mixtures or corn injury may occur.

#### Precautions

- In-row weed control may be reduced because of lack of coverage when applied to corn over 4 inches tall.

### MOCCASIN II PLUS Herbicide + Glufosinate Products Registered for Postemergence Use in Glufosinate-Tolerant Corn

These tank mixtures can be applied postemergence to weeds and corn from seed designated as corn warranted as being tolerant to glufosinate. Products containing glufosinate are used to provide postemergence control of a broad spectrum of grass and broadleaf weeds while the MOCCASIN II PLUS Herbicide provides residual control of grasses and certain broadleaf weeds listed in the label section **MOCCASIN II PLUS HERBICIDE APPLIED ALONE / Weeds Controlled**. Refer to section **MOCCASIN II PLUS HERBICIDE ALONE / Preplant Incorporated or Preemergence** and use the minimum rate per soil texture and organic matter classification for season-long residual control. Refer to the label of the glufosinate tank mix product for the postemergence application rates according to weed species and their maximum height at the time of postemergence application. Where multiple weed species are present, use the highest labeled rate to control the species and growth stages present.

Follow all applicable use directions, limitations, restrictions, and precautions regarding application to corn on the MOCCASIN II PLUS Herbicide and tank mix product labels. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

### MOCCASIN II PLUS Herbicide + Glyphosate for Postemergence Application to Glyphosate-Tolerant Corn

The tank mixture of MOCCASIN II PLUS Herbicide + glyphosate can be applied postemergence to weeds and to corn designated as glyphosate-tolerant. Application may be applied postemergence to glyphosate-tolerant corn from emergence until corn reaches 30 inches tall or the V8 stage (8 leaves with collars), whichever comes first. This mixture will provide postemergence control of weed species on the glyphosate label and residual control of weed species on the MOCCASIN II PLUS Herbicide label. Use the minimum MOCCASIN II PLUS Herbicide rate postemergence with glyphosate in glyphosate-tolerant corn as specified in the **CORN – MOCCASIN II PLUS Herbicide Alone / Preplant Incorporated or Preemergence** section of this label according to soil texture and organic matter. Refer to the glyphosate label and follow appropriate use directions, application procedures, precautions, and limitations. Refer to the glyphosate label for directions for control

of problem species. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

### MOCCASIN II PLUS Herbicide + Glyphosate + Atrazine for Postemergence Application to Glyphosate-Tolerant Corn

The tank mixture of MOCCASIN II PLUS Herbicide + atrazine + glyphosate can be applied postemergence to weeds and to corn designated as glyphosate-tolerant. Application may be applied postemergence to glyphosate-tolerant corn from emergence up to 12 inches in height. This mixture will provide postemergence control of weed species on the glyphosate label and residual control of weed species on the MOCCASIN II PLUS Herbicide + atrazine label. Use the minimum MOCCASIN II PLUS Herbicide + atrazine rate postemergence with glyphosate in glyphosate-tolerant corn as specified in the **Corn - MOCCASIN II PLUS Herbicide Combinations / Tank Mixture With atrazine or simazine, or atrazine + simazine / Preplant Incorporated or Preemergence** section and **Table 3** of this label according to soil texture and organic matter.

Follow all applicable use directions, limitations, precautions, and information regarding application to corn on the MOCCASIN II PLUS Herbicide, atrazine, and glyphosate labels for application to glyphosate-tolerant corn. Where difficult species and/or severe weed populations are expected, use the maximum rate where rate ranges are listed.

### COTTON MOCCASIN II PLUS HERBICIDE ALONE

**Application:** Apply MOCCASIN II PLUS Herbicide postemergence to cotton and preemergence to weeds at 0.5 - 1.33 pts/A, according to the state limitations in the following **Postemergence** section.

AR, KS, LA, MS, TN, and Bootheel of MO - Apply preemergence only at 0.5 - 1.0 pt/A on sandy loams, 0.66 - 1.33 pts/A on *medium soils*, or 1.0 - 1.33 pts/A on *fine soils*.

NM, OK, and TX - Apply preplant incorporated or preemergence at 1.0 pt/A on sandy loams, 1.0 - 1.33 pts/A on *medium soils*, or 1.33 pts/A on *fine soils*.

**Fall Application for glyphosate-resistant Italian Ryegrass Control:** Apply MOCCASIN II PLUS Herbicide for residual control at 1.33 - 1.67 pts/A in the fall (September 1 - December 1) after harvest of the previous crop and before Italian ryegrass emergence. Use the lower rate for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. Do not incorporate deeper than 2 - 3 inches if tillage follows the application of MOCCASIN II PLUS Herbicide. For fall applications after emergence of glyphosate-resistant Italian ryegrass, paraquat can be tank mixed with MOCCASIN II PLUS Herbicide to control emerged ryegrass. Refer to the paraquat brands label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with MOCCASIN II PLUS Herbicide for control or improved control of other weeds present at the time of application.

#### Restrictions

- **DO NOT** apply to frozen ground.
- **DO NOT** apply more than the seasonal maximum for cotton (2.6 pts/A, depending on soil texture) through any combination of applications.
- **DO NOT** apply more than a total of 2.0 pts/A (1.9 lbs ai) on *coarse soils* or 2.6 pts/A (2.5 lbs ai) of MOCCASIN II PLUS Herbicide on *medium* and *fine soils* per year.
- **DO NOT** graze or feed forage or fodder from cotton to livestock, or illegal residues may result.
- **DO NOT** apply MOCCASIN II PLUS Herbicide on sand or loamy sand soils, or in areas where water is likely to “pond” over the bed.
- **DO NOT** apply on Taloka silt loam.
- **DO NOT** use in Gaines County, TX

**Preplant Incorporated (NM, OK, and TX Only):** Apply to the soil and incorporate into the top inch of soil immediately before planting, at planting, or after planting, but before crop or weeds emerge. Use a rolling cultivator or

similar implement to uniformly incorporate not more than 1 inch deep. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. Where furrow irrigation is used, wet the top of the bed for best results. If the crop is to be planted on beds, apply and incorporate after bed formation. Plant cotton below the zone of incorporation; i.e., at least 1 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated prior to planting, use a planter that will result in the least soil disturbance.

**Note:** For best control of yellow nutsedge and suppression of seedling johnsongrass, apply MOCCASIN II PLUS Herbicide pre-plant incorporated at the maximum rate for the soil texture, whether applied alone or mixed with prometryn.

**Preemergence:** Apply to the soil surface at planting or after planting, but before weeds or crop emerge.

**Postemergence:** Apply MOCCASIN II PLUS Herbicide broadcast over the top or directed to the soil surface according to the rate limitations listed below by state. Over-the-top postemergence application may be made not later than 100 days before harvest, and directed-postemergence application may be made not later than 80 days before harvest. Application before weeds emerge or after clean cultivation to remove existing weeds is necessary since MOCCASIN II PLUS Herbicide will not control emerged weeds. MOCCASIN II PLUS Herbicide postemergence may be applied over any previous registered herbicide treatment. In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 - 1 inch of water (1/2 inch on *coarse-textured soils* to 1 inch on *fine-textured soils*) to incorporate MOCCASIN II PLUS Herbicide. In furrow-irrigated areas, apply MOCCASIN II PLUS Herbicide, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of MOCCASIN II PLUS Herbicide.

**VA, NC, SC, GA, FL, and AL:** Apply postemergence at 1.0 - 1.33 pts/A.

**TN, AR, KS, MS, MO, and LA:** Apply postemergence at 0.5 - 1.33 pts/A.

**TX, OK, NM, AZ, CA, and Clay Soils in AR:** Apply postemergence at 1.0 - 1.33 pts/A before August 1.

**Multiple Applications:** Apply as a preplant incorporated or preemergence treatment and follow with an application postemergence to cotton before weeds emerge or after clean cultivation to remove existing weeds, since MOCCASIN II PLUS Herbicide will not control emerged weeds. Apply MOCCASIN II PLUS Herbicide postemergence over a previous preplant or preemergence MOCCASIN II PLUS Herbicide application as shown in the following table.

State	Multiple MOCCASIN II PLUS Herbicide Applications to Cotton		
	Preplant Incorporated or Preemergence (Pt/A)	+	Postemergence (Pt/A)
MS, LA, TN, AR, KS, MO	0.5 - 1.33 (Preemergence Only)	+	0.5 - 1.33
TX, OK, NM	1.0 - 1.33	+	1.0 - 1.33 before August 1
NC, VA	1.0 - 1.33 (Preemergence Only)	+	1.0 - 1.33

In sprinkler-irrigated areas, sprinkler irrigate after application with 1/2 - 1 inch of water (1/2 inch on *coarse-textured soils* to 1 inch on *fine-textured soils*) to incorporate MOCCASIN II PLUS Herbicide. In furrow-irrigated areas, apply MOCCASIN II PLUS Herbicide, incorporate with a rolling cultivator or similar implement that provides uniform shallow incorporation (2 inches or less), and then irrigate. In non-irrigated areas, if at least 1/2 inch of rainfall does not occur within 10 days after application, cultivate with a rolling cultivator or similar implement that provides uniform shallow incorporation of MOCCASIN II PLUS Herbicide.



For best control of yellow nutsedge and suppression of seedling johnsongrass, apply MOCCASIN II PLUS Herbicide preplant incorporated, preemergence, or postemergence to cotton and preemergence to weeds at the maximum rate for the soil texture, whether applied alone or in combinations. These treatments may be applied over previous registered herbicide treatments.

**Precautions**

- To avoid crop injury, **DO NOT** apply over the top in fluid fertilizer or any other adjuvant, surfactant, oil, or other pesticide not specified in the **COTTON** section of this label.
- To avoid concentration in the seed furrow, **DO NOT** make broadcast applications of MOCCASIN II PLUS Herbicide to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width must not exceed the width of the bottom of the furrow.
- In furrow-planted cotton, to avoid concentration in the furrow and potential injury, **DO NOT** apply MOCCASIN II PLUS Herbicide postemergence until after first “knifing” or cultivation to level soil surface.

**COTTON**  
**MOCCASIN II PLUS HERBICIDE COMBINATIONS**

It is the pesticide user’s responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

**TANK MIXTURE WITH PROMETRYN**

Apply MOCCASIN II PLUS Herbicide tank mixtures with prometryn preplant incorporated or preemergence in water or fluid fertilizer. When fluid fertilizer is used as a carrier for MOCCASIN II PLUS Herbicide, either alone or in combination with prometryn, mix only the amount that will be sprayed in one operation. These mixtures should not be allowed to stand without agitation. Only use water as a carrier for postemergence-directed application.

In addition to those weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + prometryn, applied preplant incorporated or preemergence, controls the following weeds: junglerice, wild oats, annual morningglory, groundcherry, hairy nightshade, lambsquarters, malva, mustard, prickly sida (teaweed), purslane, ragweed, and shallow-germinating seedlings of cocklebur and coffeeweed. As a postemergence-directed application, prometryn provides postemergence control and residual control of weeds on its label, while MOCCASIN II PLUS Herbicide provides residual control of weed species on its label. MOCCASIN II PLUS Herbicide will not control emerged weeds.

**Preplant Incorporated or Preemergence:** Apply MOCCASIN II PLUS Herbicide + prometryn, either preplant incorporated or preemergence, using the labeled rate of prometryn with the indicated rate of MOCCASIN II PLUS Herbicide from the table below. Plant cotton below the zone of incorporation; i.e., at least 1.0 inch on *fine soils* and 1.5 inches on *coarse* and *medium soils*. If incorporated before planting, use a planter that will cause the least soil disturbance.

**MOCCASIN II PLUS Herbicide + Prometryn – Cotton (NM, OK, TX)**

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of prometryn.

**Broadcast Rate per Acre**

Area	Soil Texture	Rate
ALL	Sand, loamy sand	DO NOT USE
OK, and Blacklands and Gulf Coast of TX Rio Grande Valley of TX	Loams	0.8 - 1.33 pts
	Clays	1.33 pts
NM; High Plains, Rolling Plains, Edwards Plateau of TX; and Southwest TX	Sandy loam	0.8 - 1.0 pt
	Loams	0.8 - 1.33 pts
	Sandy clay loams, other clay soils	1.33 pts

**Postemergence Directed (AR, AZ, CA, LA, MS, NM, OK, TN, TX, and MO):**

Tank mix MOCCASIN II PLUS Herbicide with prometryn in water and apply postemergence directed in cotton for control of emerged weeds listed on the prometryn label and residual preemergence control of weeds controlled by MOCCASIN II PLUS Herbicide and prometryn, or application may be made after cultivation for residual preemergence control. These treatments may be applied over previous registered treatments, including MOCCASIN II PLUS Herbicide, provided the maximum label rate of any product is not exceeded. **DO NOT** apply over the top of cotton or injury may occur.

Apply MOCCASIN II PLUS Herbicide + prometryn in a minimum of 20 gallons of spray volume per acre. Follow the directions, restrictions, and precautions on the prometryn label when prometryn is applied as a postemergence-directed application. Refer to the directions, restrictions, and precautions for use of MOCCASIN II PLUS Herbicide under the **Cotton – MOCCASIN II PLUS Herbicide Alone / Postemergence** section.

Refer to the prometryn label for further instructions, restrictions, and limitations.

**Precautions**

To avoid concentration in the seed furrow, do not make broadcast applications of MOCCASIN II PLUS Herbicide + prometryn to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.

**Restrictions**

- **DO NOT** apply on sand or loamy sand soils, or in areas where water is likely to “pond” over the bed.
- **DO NOT** apply in cut areas of newly leveled fields, or in areas of excess salt.
- **DO NOT** apply to glandless cotton varieties.
- **DO NOT** graze or feed forage or fodder from cotton to livestock, or illegal residues may result.
- **DO NOT** apply on Taloka silt loam.
- **DO NOT** use in Gaines County, TX.

**TANK MIXTURE WITH FLUOMETURON**

Apply MOCCASIN II PLUS Herbicide in tank mixture with fluometuron pre-emergence for control of those weeds controlled by MOCCASIN II PLUS Herbicide alone and those as listed on the fluometuron label. This combination will also control spotted spurge, hyssop spurge, nodding spurge, and prostrate spurge. Apply to the soil surface at planting or after planting, but before weeds or crop emerge, using the appropriate rates from the table below. Apply the tank mixture postemergence to cotton but preemergence to weeds, or apply postemergence to both cotton and broadleaf weeds for control of weeds on the fluometuron label. Apply as a directed, semi-directed, or

over-the-top spray. MOCCASIN II PLUS Herbicide will not control emerged weeds but will provide preemergence control of species on its label. Where rate ranges are given for fluometuron, use the higher rate when applying postemergence to weeds that are 2 inches or less. These treatments may be applied over previous registered treatments, including MOCCASIN II PLUS Herbicide, provided the maximum label rate of any product is not exceeded.

**Mixing Instructions:** Incompatibility may occur when tank mixing MOCCASIN II PLUS Herbicide and fluometuron. To help overcome this condition, fill the spray tank 1/4 full with water or fluid fertilizer and start agitation, add the fluometuron and allow it to become dispersed. Add X-77 at 0.5% volume/volume final spray (4.0 pts/100 gals), then add the MOCCASIN II PLUS Herbicide and finally the rest of the water or fluid fertilizer. Agitate during mixing and application to maintain a uniform suspension. **DO NOT** use fluid fertilizer as a carrier for postemergence applications.

Apply labeled rates of fluometuron with the rates of MOCCASIN II PLUS Herbicide below.

### MOCCASIN II PLUS Herbicide + Fluometuron – Cotton

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of fluometuron.

#### Broadcast Rate per Acre

Soil Texture	Area 1: AR, LA, MS, Bootheel of MO and TN	Area 2: Eastern OK, Gulf Coast, RioGrande Valley, Eastern TX
Sand, loamy sand	DO NOT USE	DO NOT USE
Sandy loam	0.5 - 1.0	0.8 - 1.0
Loam, silt loam, silt	0.66 - 1.33	1.0 - 1.33
Fine soil	1.0 - 1.33	1.33

Refer to the fluometuron labels for further instructions, restrictions, and limitations.

#### Precautions

- To avoid crop injury, do not apply MOCCASIN II PLUS Herbicide + fluometuron on sand or loamy sand soils, or in areas where water is likely to “pond” over the bed. **DO NOT** use on Taloka silt loam.
- To avoid concentration in the seed furrow, do not make broadcast applications of MOCCASIN II PLUS Herbicide + fluometuron to cotton planted in furrows more than 2 inches deep. Band applications may be made to cotton planted in furrows deeper than 2 inches, but band width should not exceed the width of the bottom of the furrow.
- The use of fluometuron following the use of a systemic insecticide at planting may result in crop injury.

#### Restrictions

- **DO NOT** feed treated forage or gin trash to livestock, or graze treated areas.
- **DO NOT** use in Gaines County, TX.

### TANK MIXTURE OF MOCCASIN II PLUS HERBICIDE OR MOCCASIN II PLUS HERBICIDE + FLUOMETURON WITH PARAQUAT, OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where cotton is planted directly into a cover crop, stale seedbed, or previous crop residues, the contact herbicides paraquat, or glyphosate may be added to a tank mix of either MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide + fluometuron. When used as directed, the paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the glyphosate label. The MOCCASIN II PLUS Herbicide and MOCCASIN II PLUS Herbicide + fluometuron

portion of the tank mixture provides preemergence control of the weeds listed on this label in the **MOCCASIN II PLUS HERBICIDE** and **MOCCASIN II PLUS HERBICIDE + FLUOMETURON** sections, respectively.

Refer to the label of each product used in combination and observe the planting details, information regarding application, geographical restrictions, and all other precautions and limitations. Refer to **Mixing Instructions** under **TANK MIXTURE WITH FLUOMETURON** section.

**Application:** Apply before, during, or after planting, but before the cotton emerges. Apply the labeled rate of fluometuron with MOCCASIN II PLUS Herbicide at 0.8 - 1.0 pt/A on sandy loams, *medium-*, and *fine-textured soils*.

Apply in 20 - 60 gallons of water or fluid fertilizer per acre with ground equipment.

**Paraquat:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Glyphosate:** See the glyphosate label for weeds controlled, rates, and other use directions.

#### Precautions

- If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.
- Refer to the fluometuron labels and the **TANK MIXTURE WITH FLUOMETURON** section of this label for further instructions, precautions, and limitations.

#### Restrictions

- **DO NOT** apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.
- **DO NOT** apply MOCCASIN II PLUS Herbicide + fluometuron + glyphosate in tank mixture because of compatibility problems.
- **DO NOT** use in Gaines County, TX.

### TANK MIXTURE WITH MSMA, MSMA + PROMETRYN, OR MSMA + FLUOMETURON

Tank mix MOCCASIN II PLUS Herbicide with MSMA in water and applied postemergence directed for control of emerged weeds listed on the MSMA product label and residual preemergence control of weeds controlled by MOCCASIN II PLUS Herbicide. The addition of prometryn or fluometuron will add control of weed species on their respective labels.

**Postemergence Directed (AL, AR, AZ, CA, FL, GA, LA, MS, NC, NM, OK, SC, TN, TX, VA, and Bootheel of MO):** Apply MOCCASIN II PLUS Herbicide + MSMA postemergence directed to cotton at least 3 inches tall according to the directions, limitations, and precautions on the MSMA product label, as well as the directions, limitations, and precautions for use of MOCCASIN II PLUS Herbicide in the section for **COTTON – MOCCASIN II PLUS HERBICIDE ALONE / Postemergence**. Do not apply after first cotton bloom. These treatments may be applied over previous registered treatments, including MOCCASIN II PLUS Herbicide, provided the maximum label rate of any product is not exceeded. Fluometuron or prometryn may be added to the MOCCASIN II PLUS Herbicide + MSMA tank mixture according to the respective label directions for application to cotton at least 3 inches tall. When these mixtures are used, follow the mixing instructions for MOCCASIN II PLUS Herbicide + prometryn or fluometuron and then add the MSMA product.

#### Restriction

- **DO NOT** use MOCCASIN II PLUS Herbicide in tank mix with premixes of MSMA plus herbicides other than those registered for use in tank mixture with MOCCASIN II PLUS Herbicide on cotton.

### TANK MIXTURE WITH TRIFLURALIN FOR POST-DIRECTED FOLLOWED BY SOIL INCORPORATION APPLICATIONS

Apply MOCCASIN II PLUS Herbicide as a tank mixture with trifluralin in cotton for improved late-season weed control when used as an incorporated lay-by type application. This combination may be applied after the cotton is at least

3 inches tall and has reached the 4 true-leaf stage. Direct the application to the soil surface and away from the crop foliage. Incorporate using a sweep or rolling type cultivator to provide uniform and shallow mixing into the top 2 inches of soil. Refer to each product label for the appropriate application rates by soil type and for this application timing, and follow all product use limitations and restrictions.

**TANK MIXTURE WITH GLYPHOSATE FOR USE ON GLYPHOSATE-TOLERANT COTTON ONLY**

Apply MOCCASIN II PLUS Herbicide as a tank mixture with glyphosate in water postemergence over-the-top or postemergence directed for control of emerged weeds listed on the glyphosate labels and for residual preemergence control of weeds listed on the MOCCASIN II PLUS Herbicide label. See the **COTTON – MOCCASIN II PLUS HERBICIDE ALONE / Postemergence** section of this label for rates and timings of MOCCASIN II PLUS Herbicide and follow the glyphosate label for their respective rates, application methods, and application timing restrictions. Do not add additional spray adjuvants, surfactants, fertilizer additives, or pesticides to this tank mixture if applied postemergence over-the-top or unacceptable injury may occur. Refer to the glyphosate label and follow appropriate use directions, application procedures, precautions, and limitations.

**Precautions**

Postemergence over-the-top applications of this tank mixture may cause temporary injury in the form of necrotic spotting to exposed cotton leaves, which will not affect normal plant development.

**Restrictions**

- DO NOT apply this tank mixture postemergence to any cotton variety unless it is designated glyphosate-tolerant and unless the glyphosate formulation being used is registered for postemergence use in glyphosate-tolerant cotton.
- DO NOT apply glyphosate postemergence over-the-top to cotton past the growth stage limit specified on their respective labels.
- DO NOT use on sand or loamy sand soils in Gaines County, TX.

**SOYBEAN, IMMATURE SEED**

Apply MOCCASIN II PLUS Herbicide preplant or preemergence for the control or suppression of grass and small-seeded weeds in immature-seed soybean or other food-grade soybeans. For specific rates, see the rate table listed below.

**Preplant Surface-Applied:** For minimum-tillage or no-tillage systems only, apply MOCCASIN II PLUS Herbicide alone up to 45 days before planting. Use only split applications for treatments made 30 - 45 days before planting, with 2/3 the labeled broadcast rate for the crop and soil texture applied initially and the remaining 1/3 applied at planting. Treatments less than 30 days before planting may be made either as a split or a single application. If weeds are present at the time of treatment, apply in a tank mixture combination with a contact herbicide (e.g., paraquat, or glyphosate). Observe directions for use, precautions, and restrictions on the label of the contact herbicide. To the extent possible, do not move treated soil out of the row or move untreated soil to the surface during planting, or weed control will be diminished.

**Preplant Incorporated:** Apply MOCCASIN II PLUS Herbicide to the soil and incorporate into the top 2 inches of soil within 14 days before planting, using a finishing disk, harrow, rolling cultivator, or similar implement capable of providing uniform 2-inch incorporation. Use a preplant incorporated application if furrow irrigation is used or when a period of dry weather after application is expected. If crop will be planted on beds, apply and incorporate MOCCASIN II PLUS Herbicide after bed formation, unless specified otherwise.

**Preemergence:** Apply MOCCASIN II PLUS Herbicide during planting (behind the planter) or after planting, but before weeds emerge.

**MOCCASIN II PLUS Herbicide Broadcast Rates Per Acre**

Soil Texture	Percent Organic Matter in Soil	
	<3%	≥3%
Coarse	1 - 1.33 pts	1.33 pts
Medium	1.33 - 1.67 pts	1.33 - 1.67 pts
Fine	1.33 - 1.67 pts	1.67 - 2.0 pts

MOCCASIN II PLUS Herbicide will not control emerged weeds.

**Restrictions**

- DO NOT cut for hay within 120 days following a MOCCASIN II PLUS Herbicide application.
- DO NOT use for forage within 60 days following a MOCCASIN II PLUS Herbicide application.
- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide during any one year.

**GRASSES GROWN FOR SEED (ID, OR, WA) MOCCASIN II PLUS HERBICIDE APPLIED ALONE**

To control weeds and volunteer grasses in established grasses grown for seed, apply MOCCASIN II PLUS Herbicide to established stands of tall fescue, orchardgrass, perennial ryegrass, fine fescue, bentgrass, and Kentucky bluegrass just before, during, or immediately following the first fall rains or just before or during a late summer or early fall irrigation, but before target grasses emerge. The seed crop must have had one seed harvest or been established at least one year. The post-harvest residue (straw) should be evenly spread, removed, or burned before applying MOCCASIN II PLUS Herbicide. Rainfall or irrigation is required after application and before weed emergence for best control. MOCCASIN II PLUS Herbicide will provide preemergence control/suppression of volunteer seedlings of perennial ryegrass, fine fescue species, tall fescue, orchardgrass, bentgrass, and Kentucky bluegrass. MOCCASIN II PLUS Herbicide will control those weed species listed in the **MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and will suppress or control rattle fescue, annual bluegrass, Italian ryegrass, California brome, downy brome, and roughstalk bluegrass.

Apply MOCCASIN II PLUS Herbicide by ground equipment in a minimum of 10 gallons of water per acre using the rate listed below according to grass species.

Established Grass Crop Grown for Seed	Pt/A
Fine fescue species	1.0
Perennial ryegrass	1.0
Bentgrass	1.0 - 1.33
Kentucky bluegrass	1.0 - 1.33
Orchardgrass	1.0 - 1.33
Tall fescue	1.0 - 1.33

**Precautions**

- Tank mixtures with other pesticides, or the addition of an adjuvant, can increase the risk of crop injury. Application to perennial ryegrass and fine fescue stands under stress may cause crop injury.
- If weed escapes occur following a MOCCASIN II PLUS Herbicide application, an application of a postemergence herbicide may be necessary to control escapes. When making such an application, follow all directions, precautions, and limitations on the label of the postemergence herbicide.
- Control may be decreased if excessive straw from the previous harvest is present at application and/or insufficient rainfall/irrigation occurs.

### Restrictions

- Apply MOCCASIN II PLUS Herbicide only once per year.
- DO NOT graze forage regrowth for 60 days following application west of the Cascades.
- In areas east of the Cascades, DO NOT graze forage regrowth for 150 days following application.
- Hay may be harvested anytime between seed harvest and the next application of S-metolachlor.
- DO NOT apply after the November 15 or poor control may result.

## HORSERADISH

Apply a single application of MOCCASIN II PLUS Herbicide at a broadcast rate of 1.0 - 1.33 pts/A to the soil surface after planting, but before weed or crop emergence (i.e., preemergence). Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. MOCCASIN II PLUS Herbicide will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical means.

### Restrictions

- Apply MOCCASIN II PLUS Herbicide only once per year.
- DO NOT apply more than 1.33 pts/A (1.3 lbs ai) of MOCCASIN II PLUS Herbicide per year.

## PEANUTS

### MOCCASIN II PLUS HERBICIDE ALONE

Apply MOCCASIN II PLUS Herbicide, either preplant incorporated, postplant incorporated, or preemergence, using the appropriate rate specified below.

**Preplant Incorporated or Preemergence:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**.

**Postplant Incorporated:** Apply and shallowly incorporate MOCCASIN II PLUS Herbicide into the soil after planting, but before peanut germination. Incorporation depth and incorporating implements must be kept above the seed, or seed will be damaged.

Apply MOCCASIN II PLUS Herbicide alone, preplant incorporated, postplant incorporated, or preemergence, at a broadcast rate of 1.0 - 1.33 pts/A in the Southeast\* and 0.8 - 1.33 pts/A in NM, OK, and TX.

\* In the Southeast, use 1.33 - 2.0 pts/A and apply preemergence for partial control of Florida beggarweed.

MOCCASIN II PLUS Herbicide alone may be applied as directed after any of the following preplant incorporated herbicides when used according to their labels: benefin; trifluralin; ethalfluralin; imazethapyr; or pendimethalin.

### Restrictions

- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- DO NOT graze or feed peanut forage or fodder to livestock for 30 days following application.
- Pre-Harvest Interval (PHI): DO NOT apply within 90 days of harvest.

## PEANUTS

### MOCCASIN II PLUS HERBICIDE COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

## TANK MIXTURE WITH BENEFIN

MOCCASIN II PLUS Herbicide + benefin tank mixture applied preplant incorporated controls those weeds listed under **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** and those weeds as listed on the benefin label.

Apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of benefin in a minimum of 10 gallons of spray volume per acre for ground application or in a minimum of 5.0 gallons of spray volume per acre for aerial application. Follow the procedures for benefin on the benefin label for soil preparation and incorporation of this tank mix. Apply and incorporate MOCCASIN II PLUS Herbicide + benefin up to 14 days prior to planting. Follow all restrictions and precautions on the benefin label.

## TANK MIXTURE OR SEQUENTIALLY WITH IMAZETHAPYR

The tank mixture or sequential treatment of MOCCASIN II PLUS Herbicide and imazethapyr controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by imazethapyr alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the imazethapyr label for weeds controlled by imazethapyr.

Refer to the respective labels for application methods, timing, rates, restrictions, and precautions; and use in accordance with the more restrictive label. Do not exceed the label rate of either product. MOCCASIN II PLUS Herbicide will not control emerged weeds.

## TANK MIXTURE WITH ETHALFLURALIN

The tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by ethalfluralin alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the ethalfluralin label for weeds controlled by ethalfluralin.

Apply MOCCASIN II PLUS Herbicide + ethalfluralin preplant incorporated, using the appropriate rate from the table below. Follow labeled soil preparation procedures for ethalfluralin.

### MOCCASIN II PLUS Herbicide + Ethalfluralin – Peanuts

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of ethalfluralin.

#### Broadcast Rate per Acre

Soil Texture	Southeast	NM, OK, TX
<b>COARSE</b>	1.0 - 1.33 pts	0.8 - 1.33 pts
<b>MEDIUM</b>	1.0 - 1.33 pts	0.8 - 1.33 pts
<b>FINE</b>	1.0 - 1.33 pts	0.8 - 1.33 pts

## TANK MIXTURE WITH PENDIMETHALIN

MOCCASIN II PLUS Herbicide + pendimethalin applied preplant incorporated controls all weeds controlled by MOCCASIN II PLUS Herbicide alone plus Texas panicum, field sandbur, johnsongrass from seed, lambsquarters, Kochia, annual spurge, and other species on the pendimethalin label. Apply MOCCASIN II PLUS Herbicide + pendimethalin by ground or by aerial equipment within 14 days before planting. Incorporate into the top 1 - 2 inches of soil before planting and within 7 days of application, using a finishing disk or similar implement capable of providing uniform incorporation. If peanuts will be planted on beds, apply and incorporate after bed formation. Refer to the Incorporation instructions of the respective labels for additional directions.

Apply MOCCASIN II PLUS Herbicide + pendimethalin preplant incorporated, using the appropriate rates from the table below.

## MOCCASIN II PLUS Herbicide + Pendimethalin – Peanuts

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of pendimethalin.

### Broadcast Rate per Acre

Soil Texture	NM, OK, TX	Other Peanut Growing States
Sand, loamy sand	0.8 pt	1.0 - 1.33 pts
Sandy loam	0.8 - 1.0 pt	1.0 - 1.33 pts
Fine soil	1.33 pts	1.33 pts

## TANK MIXTURE WITH PARAQUAT

MOCCASIN II PLUS Herbicide + paraquat applied at ground cracking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section of this label. Apply paraquat plus the appropriate MOCCASIN II PLUS Herbicide rate from the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section in a minimum spray volume of 20 gals/A with ground equipment. Refer to the paraquat label and follow all directions, limitations, and restrictions.

## TANK MIXTURE WITH PARAQUAT + BENTAZON

The addition of bentazon to the MOCCASIN II PLUS Herbicide + paraquat mixture will result in improved control of such problem broadleaf weeds as prickly sida, cocklebur, smartweed, and bristly starbur. MOCCASIN II PLUS Herbicide + paraquat + bentazon applied at ground cracking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section of this label. Apply bentazon + paraquat with the appropriate MOCCASIN II PLUS Herbicide rate from the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section in a minimum spray volume of 20 gals/A with ground equipment. Refer to the paraquat and bentazon labels and follow all directions, limitations, and restrictions.

## TANK MIXTURE WITH PARAQUAT + 2,4-DB

The addition of 2,4-DB to the MOCCASIN II PLUS Herbicide + paraquat mixture will result in improved control of such problem broadleaf weeds as sicklepod, morning glory, and cocklebur. MOCCASIN II PLUS Herbicide + paraquat + 2,4-DB applied at ground cracking will control or suppress small (1- to 6-inch) emerged annual grass and broadleaf weeds and provide residual control of weed species listed in the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section of this label. Apply paraquat + 2,4-DB with the appropriated MOCCASIN II PLUS Herbicide rate from the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section in a minimum spray volume of 20 gals/A with ground equipment. Refer to the Gramoxone brands and 2,4-DB labels and follow all directions, limitations, and restrictions for each product.

## TANK MIXTURE WITH BENTAZON

MOCCASIN II PLUS Herbicide + bentazon applied at ground cracking will control species on the bentazon label and provide residual control of species listed in the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section of this label. Apply labeled rate of bentazon in 20 gals/A, depending on weed species and stage of growth as specified on the bentazon label, with the appropriate MOCCASIN II PLUS Herbicide rate from the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section. A second bentazon application may be made in all peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

## TANK MIXTURE WITH BENTAZON + BUTYRAC 200

MOCCASIN II PLUS Herbicide + bentazon + 2,4-DB applied at ground cracking will control species on their labels, especially morning glories. Apply labeled rates of bentazon and 2,4-DB in 20 gals/A, depending on weed species and stage of growth as specified on the bentazon label, with the appropriate MOCCASIN II PLUS Herbicide rate from the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section. A second bentazon + 2,4-DB application may be made in all peanut-growing areas, if needed. Refer to the respective labels and follow all directions, limitations, and restrictions for each product.

## SEQUENTIALLY WITH ACIFLUORFEN + BENTAZON

Apply MOCCASIN II PLUS Herbicide according to the directions for **MOCCASIN II PLUS HERBICIDE ALONE** and follow with a postemergence treatment of acifluorfen + bentazon as specified on its label for the control of weeds listed on the MOCCASIN II PLUS Herbicide label and on the acifluorfen + bentazon label. Refer to the **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** section and to the acifluorfen + bentazon label and follow all directions, limitations, and restrictions for each product.

## MULTIPLE APPLICATIONS

Where weed pressure is heavy or where species difficult to control are expected, MOCCASIN II PLUS Herbicide is most effective when used as follows:

### Southeast Only (AL, FL, GA, NC, SC, VA)

**Preplant Incorporated:** Apply MOCCASIN II PLUS Herbicide preplant incorporated as directed under **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** or apply MOCCASIN II PLUS Herbicide + benfen preplant incorporated as directed previously in this section. Refer to the respective section for weeds controlled.

### OR

**Preemergence to before “ground cracking”:** Apply MOCCASIN II PLUS Herbicide any time from preemergence to before “ground cracking” at 1.0 - 2.0 pts/A for extended control of weeds not yet emerged. **DO NOT use MOCCASIN II PLUS Herbicide after peanut emergence.** If peanuts have emerged, use Dual Magnum® according to its label: Peanuts – Combinations – Multiple Applications.

Follow the PPI or PRE application by:

**Lay-by:** **DO NOT use MOCCASIN II PLUS Herbicide.** Apply Dual Magnum at lay-by as directed under the Peanuts – Alone section of the Dual Magnum label.

### Southwest Only (NM, OK, TX)

**1st Application:** Apply MOCCASIN II PLUS Herbicide preplant incorporated or preemergence to before “ground cracking” as directed under **PEANUTS – MOCCASIN II PLUS HERBICIDE ALONE** or apply MOCCASIN II PLUS Herbicide + benfen preplant incorporated as directed previously in this section. **DO NOT use MOCCASIN II PLUS Herbicide after peanut emergence.** If peanuts have emerged, use Dual Magnum according to its label.

**2nd Application:** **DO NOT use MOCCASIN II PLUS Herbicide.** Apply s-metolachlor at lay-by as directed under the Peanuts – Alone section of the s-metolachlor label. Use only when late germinating weeds are expected to be a problem. Refer to the product Applied Alone section for a list of weeds controlled.

### Restrictions

- DO NOT apply more than the equivalent of 2.67 lbs of active ingredient of MOCCASIN II PLUS Herbicide per acre during any one year. If s-metolachlor is used as a sequential treatment, the lbs of active ingredient (1.0 pt = 0.95 lb) plus the lbs of active ingredient of MOCCASIN II PLUS Herbicide must not exceed 2.67 lbs.
- **DO NOT use MOCCASIN II PLUS Herbicide or other s-metolachlor products after peanuts have emerged.**
- DO NOT graze or feed peanut forage or fodder to livestock for 30 days following application.
- Pre-Harvest Interval (PHI): DO NOT apply within 90 days of harvest.

## BEANS, PEAS, AND LENTILS MOCCASIN II PLUS HERBICIDE ALONE

For use on beans, peas, and lentils, including garbanzo, great northern beans, kidney beans, lima beans, mung beans, navy beans, peas (English\*; southern peas, including blackeye, pinkeye, crowder, etc.), pinto beans, snap beans (green, wax, string), lentils, and lupines (sweet, white, white sweet, and grain).

### Fall Application:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts/A on *medium-textured* and 2.0 pts/A on *fine-textured soils*. A tillage operation may precede the application. A fall and/or a spring tillage may follow application, but do not incorporate deeper than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

### Restrictions

- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- DO NOT apply to frozen ground.
- If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for beans, peas, and lentils.

### Spring Application:

Apply MOCCASIN II PLUS Herbicide, either preplant incorporated or pre-emergence, using the appropriate rate specified below. **Preplant Incorporated or Preemergence:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**. On *coarse soils* with less than 3% organic matter, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide or 1.33 pts/A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.67 - 2.0 pts/A if organic matter content is 3% or greater.

\* On English peas, use only preemergence applications. If soils are cold and wet during pea germination and emergence, the use of MOCCASIN II PLUS Herbicide may delay maturity and/or reduce yields.

### Restrictions

- DO NOT cut for hay within 120 days following a MOCCASIN II PLUS Herbicide application.
- DO NOT use for forage within 60 days following a MOCCASIN II PLUS Herbicide application.
- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.

## BEANS, PEAS, AND LENTILS MOCCASIN II PLUS HERBICIDE COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

### Restrictions

- When applying MOCCASIN II PLUS Herbicide in combination on beans, peas, and lentils, **DO NOT** cut for hay within 120 days following application.

## TANK MIXTURE AND SEQUENTIAL APPLICATIONS WITH EPTC BEANS (GREEN OR DRY)

This mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by EPTC alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section of this label for weeds controlled by MOCCASIN II PLUS Herbicide alone and to the EPTC label for weeds controlled by EPTC.

**Preplant Incorporated:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**. **Sequential:** Apply EPTC alone preplant incorporated, as specified on that label. Follow with a pre-emergence application of MOCCASIN II PLUS Herbicide, at rates specified for MOCCASIN II PLUS Herbicide alone, during planting (behind the planter) or after planting, but before the weeds or crop emerge.

Refer to the **PRODUCT INFORMATION** section of this label and to the EPTC label for weather, cultural practices, and all other precautions and limitations that affect performance of these products.

Apply the labeled rate of EPTC with MOCCASIN II PLUS Herbicide as specified. On *coarse soils*, apply 0.8 pt/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.0 pt/A if organic matter content is 3% or greater. On *medium soils*, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.33 pts/A if organic matter content is 3% or greater. On *fine soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide if organic matter is less than 3%, or 1.33 - 1.67 pts/A if organic matter is 3% or greater.

\* Refer to the EPTC label for rate limitations depending on geographical area, limitations and restrictions, and species and varietal restrictions.

## TANK MIXTURE WITH TRIFLURALIN BEANS (DRY – KIDNEY, NAVY, PINTO, ETC.; LIMA; AND SNAP)

MOCCASIN II PLUS Herbicide + trifluralin tank mix applied preplant incorporated controls those weeds listed under **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** and those weeds listed for trifluralin alone on the trifluralin label. MOCCASIN II PLUS Herbicide + trifluralin may be applied by ground or by aerial equipment and incorporated up to 14 days prior to planting. Follow the procedures on this label and on the respective trifluralin label using equipment that provides uniform 2-inch incorporation.

Apply MOCCASIN II PLUS Herbicide + trifluralin tank mix using the appropriate MOCCASIN II PLUS Herbicide rate specified for MOCCASIN II PLUS Herbicide alone, and the labeled trifluralin rate from the Dry Beans, and the Lima and Snap Beans sections of the respective trifluralin label. Choose the product rate for the specific soil texture/organic matter classification and weed species expected.

## POTATOES MOCCASIN II PLUS HERBICIDE ALONE

Apply MOCCASIN II PLUS Herbicide, either incorporated, preemergence, or postemergence to potatoes after hilling/lay-by, according to directions specified below for control of weeds listed under the **PRODUCT INFORMATION** section. Within a rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. For application by center pivot irrigation, see the **CENTER PIVOT IRRIGATION APPLICATION** section of this label.

**Incorporated:** Apply MOCCASIN II PLUS Herbicide at 1.0 - 2.0 pts/A to the soil and incorporate into the top 3 inches before planting, using a finishing disk, harrow, rolling cultivator, or similar implement. Planting and later cultural practices should not bring untreated soil to the surface. Postplant incorporated application may be made any time after planting to drag-off, but before potato emergence. Use an implement that evenly distributes MOCCASIN II PLUS

Herbicide in the top 2 inches of soil. Do not damage potato seed pieces or sprouts with incorporation equipment.

**Preemergence:** Apply MOCCASIN II PLUS Herbicide at 1.0 - 2.0 pts/A, either after planting as a preemergence, delayed preemergence, after drag-off or hilling treatment, but before weeds emerge. Up to 2.5 pts/A of MOCCASIN II PLUS Herbicide alone may be used where soil organic matter is between 6% and 20%.

**Postemergence After Hilling/Lay-by:** Apply 1.67 pts/A of MOCCASIN II PLUS Herbicide postemergence to potatoes through after hilling/at lay-by to control MOCCASIN II PLUS Herbicide-sensitive species for remainder of the growing season. This application will not control emerged weeds. It may be applied over a previous MOCCASIN II PLUS Herbicide application, but do not apply more than 3.6 pts/A (3.4 lbs ai) of MOCCASIN II PLUS Herbicide per year.

#### Precautions

If cool, wet soil conditions occur after application, MOCCASIN II PLUS Herbicide may delay maturity and/or reduce yield of Superior and other early maturing potato varieties. These directions for use do not apply to sweet potatoes or yams.

#### Restrictions

- DO NOT apply more than 3.6 pts/A (3.4 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- DO NOT use on muck or peat soils.
- DO NOT apply both as a preemergence and an incorporated treatment.
- Pre-Harvest Interval (PHI): DO NOT harvest potatoes within 40 days after a lay-by application, or within 60 days after the at-planting to drag-off application.

### POTATOES MOCCASIN II PLUS HERBICIDE COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

#### TANK MIXTURE WITH METRIBUZIN

In addition to those weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide applied in tank mix combination with, or sequentially with, any of the registered metribuzin formulations also controls the following broadleaf weeds: hairy, hemp sesbania, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard; and will provide partial control of cocklebur, nightshade, and jimsonweed.

Apply MOCCASIN II PLUS Herbicide at 1.0 - 2.0 pts/A plus the labeled metribuzin use rate preemergence or postemergence to potatoes through after last hilling. Apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide on *coarse soils* and 1.33 - 2.0 pts/A on other soil textures. Within this rate range, use the lower rate on soils relatively coarse-textured or low in organic matter; use the higher rate on soils relatively fine-textured or high in organic matter. Effectiveness will be reduced if later cultural practices expose untreated soil. MOCCASIN II PLUS Herbicide will not control emerged weeds.

Refer to the metribuzin label for precautionary statements, restrictions, application information, center pivot irrigation application, weeds controlled, and varietal limitations.

#### Precautions

- Make postemergence applications to potatoes, except center pivot, only as a directed or semi-directed spray to avoid chlorosis, minor necrosis, or leaf distortion.
- These directions for use do not apply to sweet potatoes or yams.

#### Restrictions

- Pre-Harvest Interval (PHI): DO NOT harvest potatoes treated with MOCCASIN II PLUS Herbicide in tank mixture with metribuzin within 60 days after application.
- Pre-Harvest Interval (PHI): Potatoes may not be harvested within 40 days after a lay-by application of MOCCASIN II PLUS Herbicide.
- DO NOT use this tank mixture on muck or peat soils.

### MOCCASIN II PLUS HERBICIDE + LINURON TANK MIXTURE (EAST OF ROCKY MOUNTAINS)

Apply MOCCASIN II PLUS Herbicide in a tank mix combination with any of the registered linuron formulations as a preemergence broadcast application to potatoes. Apply to the soil surface after planting and before emergence of the crop or after final drag-off according to the rates appropriate to soil texture.

Refer to the **PRODUCT INFORMATION** section of this label and to the linuron label for precautionary statements, restrictions, application information, and weeds controlled.

#### Precautions

- To avoid crop injury, do not use on sands or loamy sands.
- To avoid crop injury, do not incorporate or spray over the top of emerged potatoes.

#### TANK MIXTURE WITH PENDIMETHALIN

In addition to the weeds controlled by MOCCASIN II PLUS Herbicide alone, this tank mixture with pendimethalin controls such problem species as kochia, lambsquarters, purslane, annual spurge, stinging nettle, and others specified on the pendimethalin label. Apply MOCCASIN II PLUS Herbicide + pendimethalin preemergence, preemergence incorporated, or early postemergence, according to the specific directions on the pendimethalin label, using the rates appropriate to soil texture.

Refer to the MOCCASIN II PLUS Herbicide and pendimethalin labels and observe all directions, timings, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

#### TANK MIXTURE WITH PENDIMETHALIN + EPTC

In addition to the weeds controlled by MOCCASIN II PLUS Herbicide alone, this tank mixture will control those species on the pendimethalin and EPTC labels. Refer to the MOCCASIN II PLUS Herbicide + pendimethalin labels for rates of those products, depending on geographical area. Refer to the respective MOCCASIN II PLUS Herbicide, pendimethalin, and EPTC labels and observe all directions, limitations, precautions, and restrictions concerning the use of these products on potatoes and follow the most restrictive.

### PUMPKIN MOCCASIN II PLUS HERBICIDE ALONE

#### Preemergence

Apply MOCCASIN II PLUS Herbicide preemergence (before the weeds have emerged) at 1.0 to 1.33 pts/A as an inter-row or inter-hill application in pumpkin. Leave 1 foot of untreated area over the row, or 6 inches to each side of the planted hill and/or any emerged pumpkin foliage (inter-row or inter-hill means not directly over the planted seed or young pumpkin plants). Use the lower MOCCASIN II PLUS Herbicide rate on soils light in texture (loamy sand or lighter) and low in soil organic matter (less than 3%). MOCCASIN II PLUS Herbicide applied as a broadcast spray over the planted row or hill, or applications made directly to crop foliage, will increase the risk of injury (e.g., stand loss, delayed maturity, and loss of yield) to the pumpkin crop.

Because MOCCASIN II PLUS Herbicide will not control emerged weeds, it must be applied before the weeds emerge. Weeds that are present should be controlled by another means, e.g., by mechanical means or using another herbicide.

#### Restriction

- DO NOT apply more than 1.33 pts/A (1.3 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- Pre-Harvest Interval (PHI): DO NOT apply within 30 days before pumpkin harvest.

### RHUBARB MOCCASIN II PLUS HERBICIDE ALONE

Apply MOCCASIN II PLUS Herbicide at a broadcast rate of 0.67 - 1.33 pts/A to the soil surface in early spring, prior to crop emergence. Use lower rates on soils relatively coarse-textured and higher rates on fine-textured soils. A band application may also be used, applying proportionally less spray mixture on the area actually treated. MOCCASIN II PLUS Herbicide will not control emerged weeds. Control emerged weeds with an appropriate registered foliar herbicide or by mechanical or physical means.

#### Restrictions

- DO NOT apply more than once per year.
- DO NOT apply more than 1.33 pts/A (1.3 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- Pre-Harvest Interval (PHI): DO NOT harvest rhubarb within 62 days of the MOCCASIN II PLUS Herbicide application.

### SAFFLOWER MOCCASIN II PLUS HERBICIDE ALONE

**Preplant Incorporated or Preemergence:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**.

On *coarse soils*, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.33 pts/A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.67 - 2.0 pts/A if organic matter content is 3% or greater.

#### Restrictions

- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.

### GRAIN OR FORAGE SORGHUM (SEED TREATED WITH SAFENER) MOCCASIN II PLUS HERBICIDE ALONE

Apply MOCCASIN II PLUS Herbicide, either preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Apply MOCCASIN II PLUS Herbicide alone only when the sorghum seed has been properly treated with a seed treatment appropriate for s-metolachlor. Preplant or preemergence applications of MOCCASIN II PLUS Herbicide to sorghum not treated with an appropriate seed treatment will result in crop death.

#### Restrictions

- DO NOT apply more than 1.67 pts/A (1.6 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- DO NOT apply MOCCASIN II PLUS Herbicide to frozen ground.
- If a spring application is made, DO NOT apply MOCCASIN II PLUS Herbicide or any other product containing S-metolachlor the following spring to grain or forage sorghum.
- Except for the split preplant surface treatment, do not make more than one application per year.

- Pre-Harvest Interval (PHI): DO NOT apply MOCCASIN II PLUS Herbicide postemergence within 75 days of harvest.

**Fall Application for Italian Ryegrass Control:** Apply MOCCASIN II PLUS Herbicide for residual control of glyphosate-resistant Italian ryegrass (*Lolium multiflorum*) using 1.33 - 1.67 pts/A in the fall (September 1 - December 1) after harvest of the previous crop and before Italian ryegrass emergence. Use the lower rate for coarse-textured soils and the higher rate for fine-textured soils. A tillage operation may precede the application. Do not incorporate deeper than 2 - 3 inches if tillage follows the application of MOCCASIN II PLUS Herbicide. For fall applications after emergence of glyphosate-resistant Italian ryegrass, Gramoxone brands can be tank mixed with MOCCASIN II PLUS Herbicide to control emerged ryegrass. Refer to the paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with MOCCASIN II PLUS Herbicide for control or improved control of other weeds present at the time of application.

**Preplant Surface-Applied:** Refer to instructions for use of MOCCASIN II PLUS Herbicide under **APPLICATION PROCEDURES** section on this label. For minimum-tillage or no-tillage systems only, apply MOCCASIN II PLUS Herbicide up to 45 days before planting in CO, IA, IL, KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts/A of MOCCASIN II PLUS Herbicide on *medium soils* or 1.67 pts/A on *fine soils*. Treatments less than 30 days prior to planting may be made either as a split or single application. Apply 1.33 pts/A of MOCCASIN II PLUS Herbicide on *coarse soils* not more than 2 weeks prior to planting. Under dry conditions, irrigation after application is necessary to move MOCCASIN II PLUS Herbicide into the soil.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of MOCCASIN II PLUS Herbicide under **APPLICATION PROCEDURES** section on this label. Broadcast 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide on *coarse soils*, 1.33 - 1.5 pts/A on *medium soils*, or 1.33 - 1.67 pts/A on *fine soils*.

**Postemergence:** Refer to instructions for use of MOCCASIN II PLUS Herbicide under **APPLICATION PROCEDURES** section on this label. MOCCASIN II PLUS Herbicide may be applied broadcast postemergence at 1.0 - 1.33 pts/A on *coarse soils*, 1.33 - 1.5 pts/A on *medium soils*, or 1.33 - 1.67 pts/A on *fine soils*. MOCCASIN II PLUS Herbicide will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or chemical means. When applied alone, MOCCASIN II PLUS Herbicide will be safe to emerged sorghum. The risk of sorghum injury increases when adjuvants (e.g., non-ionic, crop oil), nitrogen sources (e.g., AMS, UAN), or fertilizers are applied with MOCCASIN II PLUS Herbicide.

#### Precautions

- If sorghum seed is not properly treated with a seed treatment appropriate for s-metolachlor, preplant and preemergence applications of MOCCASIN II PLUS Herbicide will severely injure the crop.
- Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of MOCCASIN II PLUS Herbicide. The crop will normally outgrow this effect.
- Application of MOCCASIN II PLUS Herbicide on sorghum grown under dry mulch tillage may result in crop injury.

### GRAIN OR FORAGE SORGHUM (SEED TREATED WITH SAFENER) MOCCASIN II PLUS HERBICIDE TANK MIXTURES

Apply MOCCASIN II PLUS Herbicide preplant or preemergence (prior to sorghum emergence) tank mixtures with atrazine in water or fluid fertilizer. Apply preplant or preemergence in tank mixtures only when the sorghum seed has been properly treated with a seed treatment appropriate for s-metolachlor. Preplant or preemergence applications of MOCCASIN II PLUS Herbicide to sorghum not treated with an appropriate seed treatment will result in crop death.

**IMPORTANT: FOR TANK MIXTURES WITH ATRAZINE** – If applying MOCCASIN II PLUS Herbicide in tank mixture with atrazine, all the restrictions and rate limitations on the atrazine label must be followed if more restrictive/protective than those on this label. In addition, if atrazine is/must be applied at rates lower than those on this



label, broadleaf weed control may be affected. Refer to the atrazine label for weeds controlled at the reduced rates.

### Precautions

- Applications of MOCCASIN II PLUS Herbicide + atrazine on highly alkaline soils or on eroded areas where calcareous subsoils are exposed may cause sorghum injury.
- If sorghum seed is not properly treated with an appropriate s-metolachlor safener, MOCCASIN II PLUS Herbicide + atrazine may severely injure the crop.
- Under high soil moisture conditions prior to sorghum emergence, injury may occur following preplant and preemergence application of MOCCASIN II PLUS Herbicide + atrazine. The crop will normally outgrow this effect.
- Use of MOCCASIN II PLUS Herbicide + atrazine on sorghum grown under dry mulch tillage may cause crop injury.

### Restriction

- Except for the split preplant surface treatment, do not make more than one application per year.

## TANK MIXTURE WITH ATRAZINE

In addition to the weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + atrazine also controls the following broadleaf weeds when applied either preplant surface, preplant incorporated, or pre-emergence: cocklebur, common purslane, hairy nightshade, lambsquarters, morningglory, ragweed, smartweed, and velvetleaf.

**Preplant Surface-Applied:** Refer to instructions for use of MOCCASIN II PLUS Herbicide under **APPLICATION PROCEDURES** section on this label. For minimum-tillage or no-tillage systems only, MOCCASIN II PLUS Herbicide + atrazine may be applied up to 45 days prior to planting in IA, IL, eastern KS, MO, NE, and SD. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Apply 1.5 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *medium soils* with 1.5% organic matter or greater. Apply 1.5 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *fine soils* with less than 1.5% organic matter, or apply 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine on *fine soils* with 1.5% organic matter or greater. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is necessary to move MOCCASIN II PLUS Herbicide + atrazine into the soil.

### Precautions

- Use on coarse soils or on medium soils with less than 1.5% organic matter may cause crop injury.

**Preplant Incorporated or Preemergence:** Refer to instructions for use of MOCCASIN II PLUS Herbicide under **APPLICATION PROCEDURES** on this label. On *medium soils* with 1.5% organic matter or greater, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine. On *fine soils* with less than 1.5% organic matter, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine; on *fine soils* with 1.5% organic matter or greater, apply 1.2 - 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of atrazine.

### Precautions

Use under the following conditions may cause crop injury: on coarse soils; on medium soils with less than 1.5% organic matter; in NM, OK, or TX, except in northeast OK and the TX Gulf Coast and Blacklands areas; and preplant incorporated in AZ or the Imperial Valley of CA.

## TANK MIXTURE OF MOCCASIN II PLUS HERBICIDE OR MOCCASIN II PLUS HERBICIDE + ATRAZINE, WITH PARAQUAT, GLYPHOSATE + 2,4-D, OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS

In minimum-tillage or no-tillage systems where sorghum (seed treated with appropriate safener for s-metolachlor) is planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, glyphosate + 2,4-D, or glyphosate may be tank mixed with MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide + atrazine. See **Comment No. 7** following **Table 2**. The MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide + atrazine portion of the tank mixture provides preemergence control of the weeds listed on this label under the respective sections.

Refer to the label of each product used in combination and observe the planting details, restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before sorghum emerges. Add paraquat, glyphosate + 2,4-D, or glyphosate and apply as directed on the product labels.

**Paraquat:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

**Glyphosate + 2,4-D:** Apply at labeled rates taking into account weed species and size. See the glyphosate + 2,4-D label for weeds controlled, rates for specific weeds, and other information concerning use.

**Glyphosate:** See the glyphosate labels for weeds controlled, rates, and other use directions.

## SWEET SORGHUM (SEED TREATED WITH SAFENER)

Apply MOCCASIN II PLUS Herbicide preplant surface, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below, only when the sweet sorghum seed has been properly treated with a seed treatment appropriate for s-metolachlor. Preplant or preemergence applications of MOCCASIN II PLUS Herbicide to sweet sorghum not treated with an appropriate seed treatment will result in crop death.

**Soil-Applied:** Apply MOCCASIN II PLUS Herbicide up to 45 days before planting. Use only split applications for treatments made 30 - 45 days prior to planting, with 2/3 of the broadcast rate applied initially and the remaining 1/3 at planting. Treatments less than 30 days prior to planting may be made either as a split or single application. Under dry conditions, irrigation after application is necessary to move MOCCASIN II PLUS Herbicide into the soil.

### MOCCASIN II PLUS Herbicide Rates for Soil Applications to Sweet Sorghum

Soil Type	30 - 45 Days Prior to Planting <sup>1</sup>	<30 Days Prior to Planting	At Planting <sup>2</sup>
Coarse	Do not use	1.33 pts/A	1.0 - 1.33 pts/A
Medium	1.5 pts/A	1.5 pts/A	1.33 - 1.5 pts/A
Fine	1.67 pts/A	1.67 pts/A	1.33 - 1.67 pts/A

<sup>1</sup> Use only as a split application with 2/3 of the broadcast rate applied initially and the remaining 1/3 applied at planting.

<sup>2</sup> Preplant incorporated or preemergence.

**Post-Applied:** MOCCASIN II PLUS Herbicide may be applied postemergence to sweet sorghum for residual control of grasses and small-seeded broadleaf weeds. Postemergence application to sweet sorghum may be made to crop up to 5 inches in height. MOCCASIN II PLUS Herbicide will not control emerged weeds. Therefore, emerged weeds must be controlled by cultural or other chemical methods. When applied alone, MOCCASIN II PLUS Herbicide will be safe to emerged sweet sorghum. Use of adjuvants is prohibited on sweet sorghum.

## MOCCASIN II PLUS Herbicide Rates for Postemergence Applications to Sweet Sorghum

Soil Type	Postemergence Rate
Coarse	1.0 - 1.33 pts/A
Medium	1.33 pts/A
Fine	1.33 pts/A

### Precautions

- If sweet sorghum seed is not properly treated with a seed treatment appropriate for s-metolachlor, soil applications of MOCCASIN II PLUS Herbicide prior to sorghum emergence will severely injure the crop.
- Under high soil moisture conditions prior to sweet sorghum emergence, injury may occur following soil applications of MOCCASIN II PLUS Herbicide. The crop will normally outgrow this effect.
- To avoid crop injury, do not use MOCCASIN II PLUS Herbicide on sorghum grown under dry mulch tillage.

### Restrictions

- DO NOT apply more than once per year. MOCCASIN II PLUS Herbicide may be applied either as a soil-applied treatment or a postemergence treatment, but not both.
- Pre-Harvest Interval (PHI): DO NOT apply MOCCASIN II PLUS Herbicide postemergence within 90 days of harvest.

## SOYBEANS MOCCASIN II PLUS HERBICIDE ALONE

### NOT FOR USE IN CALIFORNIA

Apply MOCCASIN II PLUS Herbicide preplant surface-applied, preplant incorporated, preemergence, or postemergence using the appropriate rate specified below. Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES** section of this label.

### Restrictions

- DO NOT apply this product to frozen ground.
- DO NOT apply more than 2.6 pts/A (2.5 lbs ai) in a single preemergence application.
- DO NOT apply more than 2.0 pts/A (1.9 lbs ai) in a single postemergence application.
- The total MOCCASIN II PLUS Herbicide rate applied preplant, preemergence or postemergence to soybeans during any one year must not exceed 3.9 pts/A (3.77 lbs ai).
- The combined total amount of the active ingredient (s-metolachlor) from all applications to soybeans must not exceed 3.77 lbs ai/A.
- Pre-Harvest Interval (PHI): Make postemergence application at least 75 days before harvest.
- DO NOT graze or feed treated soybean forage, hay, or straw to livestock within 30 days after a preplant surface, preplant incorporated, or preemergence application of this product.
- DO NOT graze or feed treated forage or hay from soybeans to livestock following a postemergence application of MOCCASIN II PLUS Herbicide.

### Fall Application for Spring Weed Control:

1. Apply after September 30 in ND, SD, MN, WI, and north of Route 30 in IA.
2. Apply after October 15 north of Route 91 in NE and south of Route 30 in IA.
3. Apply after October 31 north of Route 136 in IL.

In all locations, apply to crop stubble after harvest when the sustained soil temperature at a 4-inch depth is less than 55°F and falling. In minimum-till or no-tillage systems on soils having greater than 2.5% organic matter, use 1.67 - 2.0 pts/A of MOCCASIN II PLUS Herbicide on *medium-textured* and 2.0 pts/A of MOCCASIN II PLUS Herbicide on *fine-textured soils*. A tillage operation may precede the application. A fall and/or a spring tillage may follow

application, but do not incorporate deeper than 2 - 3 inches. Minimize furrow and ridge formation in the tillage operations.

**Restrictions:** If a spring application is made, the total rate of the fall plus spring applications must not exceed the maximum total rate for soybeans of 2.5 pts/A depending on soil texture. DO NOT apply to frozen ground.

**Fall Application for Italian Ryegrass Control:** MOCCASIN II PLUS Herbicide may be applied for residual control of Italian ryegrass (*Lolium multiflorum*), including glyphosate-resistant populations. Apply MOCCASIN II PLUS Herbicide at 1.33 - 1.67 pts/A in the fall (September 1 - December 1) after harvest of the previous crop and prior to Italian ryegrass emergence. Use the lower MOCCASIN II PLUS Herbicide rate for *coarse-textured soils* and the higher rate for *fine-textured soils*. A tillage operation may precede the application. Do not incorporate to a depth greater than 2 - 3 inches if tillage follows the application of MOCCASIN II PLUS Herbicide. For fall applications after emergence of glyphosate-resistant Italian ryegrass, paraquat can be tank mixed with MOCCASIN II PLUS Herbicide for control emerged ryegrass. Refer to the paraquat label for specific rates, application instructions, and restrictions. Other registered herbicides may be tank mixed with MOCCASIN II PLUS Herbicide for control of improved control of other weeds present at the time of application.

**Preplant Surface – Spring Application:** Use on medium and fine soils with minimum-tillage or no-tillage systems. Apply 2/3 the labeled rate of MOCCASIN II PLUS Herbicide (1.67 pts/A on *medium soils* and 2.0 pts/A on *fine soils*). Applications made less than 30 days before planting may be as either a split or single treatment. Apply 1.33 pts/A of MOCCASIN II PLUS Herbicide on *coarse soils* not more than 2 weeks prior to planting. On soils with 6 - 20% organic matter or for extended residual or control of heavy weed infestations, up to 2.5 pts/A is allowed.

**Preplant Incorporated or Preemergence:** On *coarse soils*, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.33 pts/A if organic matter content is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide if organic matter content is less than 3%, or 1.67 - 2.0 pts/A if organic matter content is 3% or greater. On soils with 6 - 20% organic matter or for extended residual or control of heavy weed infestations, up to 2.5 pts/A is allowed.

**Postemergence:** Apply 1.0 - 2.0 pts/A as a postemergence treatment to soybeans from emergence up through the third trifoliolate leaf stage. MOCCASIN II PLUS Herbicide will not control emerged weeds so it must be applied to a weed-free soil surface or in a tank mixture with products that provide postemergence control of weeds present at the time of application.

MOCCASIN II PLUS Herbicide can also be applied as part of a sequential soybean weed control program. If MOCCASIN II PLUS Herbicide was applied as a preplant surface, preplant incorporated, or a preemergence treatment, a second treatment of MOCCASIN II PLUS Herbicide can be applied postemergence provided that the total MOCCASIN II PLUS Herbicide rate during any one crop does not exceed 3.7 pts/A.

## SOYBEANS MOCCASIN II PLUS HERBICIDE COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

Water or fluid fertilizer may be used as carrier for MOCCASIN II PLUS Herbicide in combination with metribuzin, linuron, metribuzin + chlorimuron-ethyl, imazethapyr, imazaquin, ethalfluralin, or clomazone.

### Restrictions

- For all of the following combinations, on soybeans use up to 2.5 pts/A MOCCASIN II PLUS Herbicide preplant incorporated or preemergence treatment on soils having an organic matter content between 6% and 20%. The total MOCCASIN II PLUS Herbicide rate applied to soybeans must not exceed 3.9 pts/A (3.77 lbs ai) per year.

## TANK MIXTURE WITH METRIBUZIN

In addition to those weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + metribuzin, when applied as directed, also controls hairy nightshade, hemp sesbania, lambsquarters, prickly sida, ragweed, smartweed, velvetleaf, Venice mallow, and wild mustard, and provides partial control of cocklebur and jimsonweed.

Apply MOCCASIN II PLUS Herbicide and metribuzin preplant incorporated or preemergence, using the appropriate rates from the table below. **Preplant Incorporated or Preemergence:** Follow instructions for use of MOCCASIN II PLUS Herbicide alone under **APPLICATION PROCEDURES**.

**Sequential:** Apply MOCCASIN II PLUS Herbicide alone **Preplant Incorporated**, as specified in the table below for this tank mixture. Follow with a preemergence application of metribuzin during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

Refer to the metribuzin label for planting details and soybean variety restrictions.

### MOCCASIN II PLUS Herbicide + Metribuzin – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of metribuzin.

#### Broadcast Rate per Acre

Soil Texture*	0.5 to less than 3% organic matter MOCCASIN II PLUS Herbicide	3% organic matter or greater MOCCASIN II PLUS Herbicide
<b>COARSE</b> Loamy sand (over 2% organic matter), sandy loam	0.8 - 1.0 pt	1.0 pt
<b>MEDIUM</b>	1.0 - 1.33 pts	1.33 pts
<b>FINE</b>	1.33 pts	1.33 - 1.67 pts
<b>Mississippi Delta only</b> Silty clay, clay	1.33 pts	1.33 - 1.67 pts
<b>Muck or Peat</b> (soils with more than 20% organic matter) – DO NOT APPLY		

\* On all sand and on loamy sand with less than 2% organic matter, do not use this tank mixture preemergence, or the sequential treatment. Do not use the tank mixture preplant incorporated on any sand, loamy sand, or sandy loam, or crop injury may occur.

#### Restrictions

Follow most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the metribuzin label.

#### Precautions

To avoid crop injury, do not use the tank mix or sequential application on soil with less than 0.5% organic matter or on alkaline soil with a pH over 7.4, or crop injury may occur. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days.

## TANK MIXTURE WITH LOROX

In addition to those weeds controlled by MOCCASIN II PLUS Herbicide alone, MOCCASIN II PLUS Herbicide + linuron, applied preemergence, also controls the following broadleaf weeds: lambsquarters, prickly sida, ragweed, smartweed, Venice mallow, and wild mustard, and provides partial control of cocklebur, jimsonweed, morningglory, and velvetleaf.

**Preemergence:** Apply during planting (behind planter) or after planting, but before weeds or soybeans emerge. Refer to the linuron label for planting details. Apply the appropriate rates from the table below.

#### Precaution

To avoid crop injury, do not use on soil with less than 0.5% organic matter.

### MOCCASIN II PLUS Herbicide + Linuron – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of linuron.

#### Broadcast Rate per Acre

Soil Texture*	0.5% to Less Than 3% Organic Matter	3% Organic Matter or Greater
<b>COARSE**</b>	0.8 pt	1.0 pt
<b>MEDIUM</b>	1.0 pt	1.33 pts
<b>FINE</b>	1.33 pts	1.33 - 1.67 pts
Muck or Peat (soils with more than 20% organic matter) – <b>DO NOT USE</b>		

\* DO NOT use on sand, gravelly soils, or exposed subsoils.

\*\* DO NOT use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter.

## TANK MIXTURE WITH TRIFLURALIN

MOCCASIN II PLUS Herbicide + trifluralin tank mix applied preplant incorporated controls those weeds listed under **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** and those weeds listed for trifluralin alone on the trifluralin label. MOCCASIN II PLUS Herbicide + trifluralin may be applied by ground or aerial equipment and incorporated up to 14 days before planting. Follow the procedures on the trifluralin and MOCCASIN II PLUS Herbicide labels, using equipment that provides uniform 2-inch incorporation.

Apply MOCCASIN II PLUS Herbicide + trifluralin tank mix using the appropriate rate from the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of this label and the trifluralin alone section of the trifluralin label for the specific soil texture/organic matter classification and weed species expected.

To control DNA-resistant goosegrass\* and other species on the respective labels where the soil organic matter is 3% or less, apply the rate in the table below.

### MOCCASIN II PLUS Herbicide + Trifluralin – Organic Matter Content Less Than 3%

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of trifluralin.

#### Broadcast Rate per Acre

	Organic Matter Less Than 3%
<b>COARSE*</b>	0.8 - 1.0 pt
<b>MEDIUM</b>	1.0 pt
<b>FINE</b>	1.33 pts

\* Where a range of rates is given for MOCCASIN II PLUS Herbicide, use the minimum rate where DNA-resistant goosegrass is the predominant species.

Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the trifluralin labels.

## TANK MIXTURE WITH IMAZAQUIN

This tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by imazaquin alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the imazaquin label for weeds controlled by imazaquin. Refer to the imazaquin label for geographical locations where this tank mixture may be applied.

Apply MOCCASIN II PLUS Herbicide + imazaquin preplant incorporated or preemergence, using rates in the table below. Follow use directions under Application Instructions on the imazaquin label. For preplant incorporated applications, apply and incorporate within 30 days before planting. Observe all other precautions and limitations on the imazaquin labels.

### MOCCASIN II PLUS Herbicide + Imazaquin – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of imazaquin.

#### Broadcast Rate per Acre

Soil Texture	Less Than 3% Organic Matter	3% or More Organic Matter
COARSE	0.8 pt	1.0 pt
MEDIUM	1.0 pt	1.33 pts
FINE	1.33 pts	1.33 - 1.67* pts
Muck or Peat (soils with more than 20% organic matter) – DO NOT USE		

\* Use the higher rate of MOCCASIN II PLUS Herbicide if heavy weed infestations are expected.

**Restrictions:** Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the imazaquin label.

### TANK MIXTURE WITH METRIBUZIN + CHLORIMURON-ETHYL

This tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by metribuzin + chlorimuron-ethyl alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the metribuzin + chlorimuron-ethyl label for weeds controlled by metribuzin + chlorimuron-ethyl.

Apply preplant incorporated or preemergence, using the appropriate rates from the table below. **Preplant Incorporated:** Apply within 2 weeks of planting. Uniformly incorporate into the top 1 - 2 inches of soil before planting soybeans. **Preemergence:** Apply after planting, but before soybeans emerge.

Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the metribuzin + chlorimuron-ethyl label, including varietal restrictions.

#### MOCCASIN II PLUS Herbicide + Metribuzin + Chlorimuron-ethyl – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of metribuzin + chlorimuron-ethyl.

#### Broadcast Rate per Acre

Soil Texture	Less Than 3% Organic Matter	3% or More Organic Matter
COARSE	0.8 pt	1.0 pt
MEDIUM	1.0 pt	1.33 pts
FINE	1.33 pts	1.33 - 1.67* pts
Muck or Peat (soils with more than 20% organic matter) – DO NOT USE		

\* Refer to the metribuzin + chlorimuron-ethyl label for appropriate rate according to geographical location, soil and organic matter classification, and pH limitations.

**Restriction:** DO NOT apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the metribuzin + chlorimuron-ethyl label.

### TANK MIXTURE WITH CLOMAZONE

This tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by clomazone alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the clomazone label for weeds controlled by clomazone.

Apply MOCCASIN II PLUS Herbicide + clomazone preplant incorporated, using rates in the table below. Follow all clomazone application instructions as

to incorporation interval, geographical location, equipment operation, soil moisture conditions, etc.

Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the clomazone label, including rotational restrictions.

#### MOCCASIN II PLUS Herbicide + Clomazone – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of clomazone.

#### Broadcast Rate per Acre

Soil Texture	0.5 - 3% Organic Matter	Greater Than 3% Organic Matter
COARSE	0.8 pt	1.0 pt
MEDIUM	1.0 pt	1.33 pts
FINE	1.33 pts	1.33 - 1.67 pts

### TANK MIXTURE WITH ETHALFLURALIN

This tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by ethalfluralin alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the ethalfluralin label for weeds controlled by ethalfluralin.

Apply MOCCASIN II PLUS Herbicide and ethalfluralin preplant incorporated, using the appropriate rates from the table below.

**Preplant Incorporated:** Follow soil preparation procedures for ethalfluralin.

**Sequential:** Apply ethalfluralin alone preplant incorporated as specified on the ethalfluralin label. Follow with a preemergence application of MOCCASIN II PLUS Herbicide during planting (behind the planter) or after planting, but before weeds or soybeans emerge.

#### MOCCASIN II PLUS Herbicide + Ethalfluralin – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of ethalfluralin.

#### Broadcast Rate per Acre

Soil Texture	Less Than 3% Organic Matter	3% or More Organic Matter
COARSE	1.0 - 1.33 pts	1.33 pts
MEDIUM	1.33 - 1.67 pts	1.33 - 1.67 pts
FINE	1.33 - 1.67 pts	1.67 - 2.0 pts
Muck or Peat (soils with more than 20% organic matter) – DO NOT USE		

Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the ethalfluralin label.

### TANK MIXTURE WITH IMAZETHAPYR

This tank mixture controls all weeds controlled by MOCCASIN II PLUS Herbicide alone and by imazethapyr alone. Refer to the **MOCCASIN II PLUS HERBICIDE APPLIED ALONE** section for weeds controlled by MOCCASIN II PLUS Herbicide and to the imazethapyr label for weeds controlled by imazethapyr. Refer to the imazethapyr label for geographical locations where this tank mixture may be applied.

Apply MOCCASIN II PLUS Herbicide + imazethapyr early preplant, preplant incorporated, or preemergence after planting, using rates in the table below. Application can be made in water or liquid fertilizer. Follow all use directions

under Soil Applications on the imazethapyr label. For early preplant and preplant incorporated applications, apply within 30 days before planting.

Follow the most restrictive limitations and precautions on the **SOYBEANS – MOCCASIN II PLUS HERBICIDE ALONE** section of the MOCCASIN II PLUS Herbicide label and the Soybean directions on the imazethapyr label, including rotational restrictions.

#### MOCCASIN II PLUS Herbicide + Imazethapyr – Soybeans

Apply indicated rates of MOCCASIN II PLUS Herbicide with labeled rates of imazethapyr.

#### Broadcast Rate per Acre

Soil Texture	Less Than 3% Organic Matter	3% or More Organic Matter
<b>COARSE</b>	0.8 pt	1.0 pt
<b>MEDIUM</b>	1.0 pt	1.33 pts
<b>FINE</b>	1.33 pts	1.33 - 1.67 pts

**Sequential:** Apply MOCCASIN II PLUS Herbicide early preplant, preplant incorporated, or preemergence after planting at 0.8 pt/A on *coarse soils* and 1.0 pt/A on *medium-* and *fine-textured soils*. Follow with a sequential postemergence application of imazethapyr to control emerged weeds according to the imazethapyr label. MOCCASIN II PLUS Herbicide will improve the consistency and level of control from imazethapyr on most grass species. Refer to the imazethapyr postemergence label for a listing of weeds controlled, application rate, and growth stage limitations.

**TANK MIXTURE WITH METRIBUZIN, IMAZAQUIN, LINURON, METRIBUZIN + CHLORIMURON-ETHYL, OR IMAZETHAPYR, PLUS PARAQUAT, OR GLYPHOSATE FOR MINIMUM-TILLAGE OR NO-TILLAGE SYSTEMS**

In minimum-tillage or no-tillage systems where soybeans are planted directly into a cover crop, stale seedbed, established sod, or previous crop residues, the contact herbicides paraquat, or glyphosate may be added to a tank mix of either MOCCASIN II PLUS Herbicide + metribuzin, MOCCASIN II PLUS Herbicide + imazaquin, MOCCASIN II PLUS Herbicide + linuron, MOCCASIN II PLUS Herbicide + metribuzin + chlorimuron-ethyl, or MOCCASIN II PLUS Herbicide + imazethapyr. When used as directed, the paraquat portion of the tank mixture controls most emerged weeds and suppresses many perennial weeds. Glyphosate combinations will control emerged annual and perennial weeds when applied as directed on the glyphosate label. The MOCCASIN II PLUS Herbicide + metribuzin, imazaquin, linuron, metribuzin + chlorimuron-ethyl, or imazethapyr portion of the tank mixture provides preemergence control of the weeds listed on this label in the tank mixture section for MOCCASIN II PLUS Herbicide + metribuzin, MOCCASIN II PLUS Herbicide + imazaquin, MOCCASIN II PLUS Herbicide + metribuzin + chlorimuron-ethyl, and MOCCASIN II PLUS Herbicide + imazethapyr, respectively.

Refer to the label of each product used in combination and observe the planting details, soybean variety restrictions, information regarding application to soybeans, geographical restrictions, and all other precautions and limitations.

**Application:** Apply before, during, or after planting, but before the soybeans emerge. Add paraquat or glyphosate and apply as directed on the product labels.

**Paraquat:** Apply as directed on the product label. This treatment will not control weeds taller than 6 inches.

#### Restriction

- DO NOT apply combinations containing paraquat in suspension-type liquid fertilizers, as the activity of paraquat will be reduced.

**Glyphosate:** See the glyphosate label for weeds controlled, rates, and other use directions.

Apply in 20 - 60 gallons of water or fluid fertilizer per acre with ground equipment.

#### MOCCASIN II PLUS Herbicide + Metribuzin + Paraquat, or Glyphosate

On loamy sand with over 2% organic matter, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of metribuzin. On *medium soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of metribuzin. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of metribuzin.

#### Precautions

To avoid crop injury, do not use this tank mixture on soil with less than 0.5% organic matter, on alkaline soil with a pH over 7.4, or on all sand and on loamy sand with less than 2% organic matter. If heavy rain occurs soon after application, crop injury may result, especially in poorly drained areas where water stands for several days, or where the seeding slit has not been properly closed.

#### MOCCASIN II PLUS Herbicide + Imazaquin + Paraquat, or Glyphosate

On *coarse soils*, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of imazaquin. On *medium soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of imazaquin. On *fine soils*, apply 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of imazaquin.

#### Restrictions

- Pre-Harvest Interval (PHI): DO NOT apply within 90 days of harvest.
- DO NOT graze or feed treated soybean forage, hay, or straw to livestock.

#### MOCCASIN II PLUS Herbicide + Linuron + Paraquat, Glyphosate

On *coarse soils*\*, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide + labeled rate of linuron. On *medium soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of linuron. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of linuron.

\* DO NOT use on loamy sand, except in the northeastern U.S. on loamy sand with over 1% organic matter, or injury may occur. DO NOT use on sand, gravelly soils, or exposed subsoils, or injury may occur.

#### Precaution

- To avoid crop injury, do not use on soil with less than 0.5% organic matter.

#### MOCCASIN II PLUS Herbicide + Metribuzin + chlorimuron-ethyl, + Paraquat, Glyphosate

Use only where soils have 0.5 - 5% organic matter. On *coarse soils* (except sand), apply 1.0 pt/A of MOCCASIN II PLUS Herbicide, on *medium soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide, and on *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. Refer to the metribuzin + chlorimuron-ethyl label for appropriate rate, according to geographical location, soil and organic matter classification, pH limitations, and all other use directions.

#### Restriction

- DO NOT apply to sand, or to any soil with less than 0.5% organic matter, or to any soil with pH greater than 7.0, except as noted on the metribuzin + chlorimuron-ethyl label.

#### MOCCASIN II PLUS Herbicide + Imazethapyr + Paraquat, Glyphosate

On *coarse soils*, apply 1.0 pt/A of MOCCASIN II PLUS Herbicide, on *medium soils*, apply 1.33 pts/A of MOCCASIN II PLUS Herbicide, on *fine soils*, apply 1.67 pts/A of MOCCASIN II PLUS Herbicide + labeled rate of imazethapyr. Refer to the imazethapyr + paraquat and glyphosate labels for appropriate rate, according to geographical location, soil and organic matter classification, pH limitations, and all other use directions.

## POSTEMERGENCE USE ON SOYBEANS MOCCASIN II PLUS HERBICIDE TANK MIXTURES

### Tank Mixture with Glyphosate

MOCCASIN II PLUS Herbicide at 1.0 - 1.33 pts/A may be tank mixed with glyphosate products at labeled rates and applied from emergence up through the third trifoliolate leaf stage of glyphosate-tolerant soybeans. MOCCASIN II PLUS Herbicide alone will not control emerged weeds. Use this treatment only on soybeans designated for use with glyphosate (e.g., glyphosate-tolerant soybeans). The glyphosate product must be registered for postemergence use in glyphosate-tolerant soybeans.

### Tank Mixture with Glufosinate

**Use this treatment only on soybeans designated for use with glufosinate.** MOCCASIN II PLUS Herbicide at 1.0 - 1.33 pts/A may be tank mixed with glufosinate herbicides at labeled rates and applied from emergence up through the third trifoliolate leaf stage of soybeans. MOCCASIN II PLUS Herbicide alone will not control emerged weeds.

Follow the tank mix product label for adjuvant suggestions. The use of COC or UAN with MOCCASIN II PLUS Herbicide may result in temporary crop injury.

#### Restrictions

- DO NOT apply more than 1.33 pts/A (1.3 lbs ai) postemergence.
- Pre-Harvest Interval (PHI): Make postemergence application at least 90 days before harvest.
- DO NOT graze or feed treated forage or hay from soybeans to livestock following a postemergence application of MOCCASIN II PLUS Herbicide.

## SUGAR BEETS MOCCASIN II PLUS HERBICIDE ALONE

### Postemergence Applications

MOCCASIN II PLUS Herbicide may be applied postemergence to sugar beets after the sugar beets have reached the first true-leaf stage. However, because MOCCASIN II PLUS Herbicide is primarily a soil-active herbicide, it must be applied prior to weed emergence in order to provide consistent control of listed weeds. As such, weeds that are emerged with or before the crop, or that are present at the time MOCCASIN II PLUS Herbicide is applied, must be controlled with another appropriately labeled herbicide. Apply MOCCASIN II PLUS Herbicide at 1 pt/A on *coarse soils*, 1.33 pts/A on *medium soils*, and 1.67 pts/A on *fine soils*. More than one postemergence application may be applied, but the total must not exceed 2.6 pts/A. Weeds present at the time of application will not be controlled.

#### Precaution

In coarse soils, MOCCASIN II PLUS Herbicide applied before emergence of sugar beets (i.e., other than postemergence) may cause injury.

#### Restrictions

- DO NOT apply more than 2.6 pts/A (2.5 lbs ai) postemergence.
- Pre-Harvest Interval (PHI): DO NOT harvest within 60 days after the last application.

## SUGAR BEETS MOCCASIN II PLUS HERBICIDE TANK MIX COMBINATIONS

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and directions for use on all product labels involved in tank mixing. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

MOCCASIN II PLUS Herbicide may be tank mixed with quizalofop p-ethyl, sethoxydim, clethodim, clopyralid, or flusulfuron methyl and applied to sugar beets. Tank mixtures of these products with MOCCASIN II PLUS Herbicide will increase the risk of crop injury over that of either product applied alone, as the MOCCASIN II PLUS Herbicide formulation has some adjuvant properties. The addition of a spray adjuvant such as crop oil concentrates (COC's) or methylated seed oils (MSO's) can further increase the risk of crop injury. Injury risk can be reduced by using the lowest effective rate of the tank mix partner(s) and/or adjuvant and by avoiding applications under adverse growing conditions or high soil or air humidity.

## SUGARCANE

Apply MOCCASIN II PLUS Herbicide as a preplant, preemergence, or postemergence treatment for weed control in sugarcane. It may also be used in a treatment program that includes a preplant/preemergence application followed by a postemergence/post-directed application.

#### Restrictions:

- **DO NOT** apply more than 1.71 qts/A (3.26 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- **DO NOT** apply more than 1.22 qts/A (2.34 lbs ai) of MOCCASIN II PLUS Herbicide as a preplant or preemergence application per year.
- **DO NOT** apply more than 0.98 qt/A (1.88 lbs ai) of MOCCASIN II PLUS Herbicide as a postemergence application per year.
- **DO NOT** apply more than 0.49 qt/A (0.94 lb ai) of MOCCASIN II PLUS Herbicide as a postemergence application if a preplant or preemergence application was made.
- **DO NOT** make more than two applications per year.
- **DO NOT** make applications less than 14 days apart.
- **DO NOT** apply to sugarcane that is taller than 60 inches.
- **DO NOT** make a postemergence application within 100 days of harvest.
- **DO NOT** exceed the total combined maximum annual sugarcane rates for s-metolachlor container products.

Apply MOCCASIN II PLUS Herbicide at a rate of 0.89 - 1.22 qts/A before planting, preemergence after new plantings, or after harvest, but before re-emergence of ratoon-cane. Use the higher rate on heavier soils and soils with higher organic matter content.

### Postemergence Applications

MOCCASIN II PLUS Herbicide will not control emerged weeds so apply only to a weed-free soil surface or in tank mixture with products that provide postemergence control of weeds present at the time of application. Apply MOCCASIN II PLUS Herbicide at 0.49 - 0.98 qt/A postemergence before the sugarcane reaches 60 inches in height. Postemergence applications may be made as broadcast post-over-the-top or as post-directed spray to soil between the rows and the base of the sugarcane.

If a preemergence application was made earlier in the season (not to exceed 1.22 qts/A), only 0.49 qt/A may be applied postemergence. The total amount of MOCCASIN II PLUS Herbicide applied (preemergence + postemergence) may not exceed 1.71 qts/A/year.

## SUGARCANE MOCCASIN II PLUS HERBICIDE TANK MIX COMBINATIONS

It is the pesticide user's responsibility to ensure that all products in the listed mixtures are registered for the intended use. Users must follow the most restrictive directions for use and precautionary statements of each product in the tank mixture.

This product may be tank mixed with other registered sugarcane herbicides, insecticides, and fungicides. The tank mix partners are to be applied by the same methods and the same timings as MOCCASIN II PLUS Herbicide unless

otherwise specified in the tank mix product label. Perform a compatibility test before spraying the tank mix application.

## SUNFLOWERS MOCCASIN II PLUS HERBICIDE ALONE

### Preplant Incorporated or Preemergence

Within the rate ranges given below, use the higher rate of MOCCASIN II PLUS Herbicide if heavy weed infestations are expected. On *coarse soils* with organic matter less than 3%, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide; apply 1.33 pts/A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils* with organic matter of less than 3%, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide; apply 1.67 - 2.0 pts/A if organic matter is 3% or greater.

#### Restrictions

- **DO NOT** apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.
- **DO NOT** allow livestock to graze or feed in treated area.
- **DO NOT** exceed the maximum label rates for sunflowers for the soil type.

## TOMATOES MOCCASIN II PLUS HERBICIDE ALONE

### Transplanted

Apply MOCCASIN II PLUS Herbicide preplant incorporated or preplant before transplanting. If the latter method is used, keep soil disturbance to a minimum during the transplanting operation. Application may also be post directed to transplants after the first settling rain or irrigation. When an application is made post-directed, apply in a minimum of 20 gallons of water per acre and minimize contact with tomato plants. MOCCASIN II PLUS Herbicide will not control emerged weeds. In bedded transplanted tomatoes, apply MOCCASIN II PLUS Herbicide preplant non-incorporated to the top of the pressed bed as the last step prior to laying plastic. MOCCASIN II PLUS Herbicide may also be used to treat row-middles in bedded tomatoes, as long as the total amount of MOCCASIN II PLUS Herbicide does not exceed the maximum allowed per crop.

### Seeded

MOCCASIN II PLUS Herbicide may be applied post-directed to direct-seeded tomatoes. Tomato plants must be at least 4 inches tall at the time of application, and the product must be applied in a minimum of 20 gallons of water per acre. Minimize spray contact with tomato plants. MOCCASIN II PLUS Herbicide will not control emerged weeds.

**Tomato Use Rates:** On *coarse soils*, apply 1.0 - 1.33 pts/A of MOCCASIN II PLUS Herbicide if organic matter is less than 3% or 1.33 pts/A if organic matter is 3% or greater. On *medium soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide. On *fine soils*, apply 1.33 - 1.67 pts/A of MOCCASIN II PLUS Herbicide if organic matter is less than 3% or 1.67 - 2.0 pts/A if organic matter is 3% or greater.

#### Precautions

- May damage transplants that have been weakened by any cause. To prevent damage, plant only healthy transplants. Do not plant when wet, cool, or unfavorable growing conditions exist.
- In transplanted tomatoes, if MOCCASIN II PLUS Herbicide is applied preplant incorporated, incorporate to a depth less than the depth of transplanting, and use the lower end of the rate range for the given soil type, or damage may occur.
- For row-middle applications where tomatoes are grown on sandy soils and where high soil moisture conditions can exist (e.g., low binding and high evaporation conditions), as may be found in the States of Florida, Georgia, Maryland, and Virginia, there is potential for crop injury in the form of leaf

epinasty. The risk of this type of injury can be reduced by: a) incorporating the MOCCASIN II PLUS Herbicide immediately following application, b) applying the MOCCASIN II PLUS Herbicide seven or more days before transplanting (but only after the beds have been formed), c) minimizing the application of MOCCASIN II PLUS Herbicide onto the plastic of the bed, or d) any combination of the above.

- Applications may be made using ground equipment, in concentrated spray volumes.
- Applications may be made as a foliar broadcast spray to the soil within 1 week of transplanting and again at blooming/fruitletting to the row middles as a banded/directed application 38 - 77 days after the first treatment.

#### Restrictions

- **DO NOT** apply more than 2.0 pts/A (1.9 lbs ai) of MOCCASIN II PLUS Herbicide per year.
  - **DO NOT** apply to varieties or cultivars with unknown tolerance to MOCCASIN II PLUS Herbicide.
  - **DO NOT** exceed the maximum label rate for the soil texture per year.
  - Apply only by ground application.
  - 90-Day PHI – If the single application rate of MOCCASIN II PLUS Herbicide is greater than 1.33 pts/A (up to 2.0 pts/A), do not harvest tomatoes within 90 days of application.
  - 30-Day PHI – If the application of MOCCASIN II PLUS Herbicide does not exceed 1.33 pts/A, do not harvest tomatoes within 30 days of application.
- When applying at 1.33 pts/A with a 30-day PHI, the following restrictions apply:
- **DO NOT** exceed two applications per year.
  - The use of adjuvants is prohibited.

## MOCCASIN II PLUS HERBICIDE FOR USE ON SOD FARMS

**DO NOT USE IN GREENHOUSES OR OTHER ENCLOSED STRUCTURES.**

### APPLICATION PROCEDURES

**Ground Application:** Apply MOCCASIN II PLUS Herbicide alone or in tank mixtures by ground equipment in a minimum of 10 gallons of spray mixture per acre, unless otherwise specified.

Use sprayers that provide accurate and uniform application. For MOCCASIN II PLUS Herbicide tank mixtures with wettable powder or dry flowable formulations, screens and strainers should be no finer than 50 - mesh. Rinse sprayer thoroughly with clean water immediately after use.

Calculate the amount of herbicide needed for band treatment by the following formula:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \text{broadcast rate per acre} = \text{amount needed per acre of field}$$

**Aerial Application (Sod Farms Only):** Apply MOCCASIN II PLUS Herbicide in water alone or in tank mixtures with atrazine, simazine, or other herbicides registered for use on sod farms in a minimum total volume of 2 gals/A by aircraft. See **Turfgrass** section for a listing of applicable warm-season grasses. Avoid application under conditions where uniform coverage cannot be obtained or where excessive spray drift may occur. In order to assure that spray will be controllable within the target area when used according to label directions, apply at a maximum height of 10 ft, using low-drift nozzles at a maximum pressure of 40 psi, and restrict application to periods when wind speed is 10 mph or less. To assure that spray will not adversely affect adjacent sensitive nontarget plants, apply MOCCASIN II PLUS Herbicide or MOCCASIN II PLUS Herbicide mixtures at a minimum upwind distance of 400 ft from sensitive plants.

Avoid application to humans or animals. Flagmen and loaders must avoid inhalation of spray mist and prolonged contact with skin.

## Turfgrass

### Warm Season Grasses (Bermudagrass, Centipedegrass, St. Augustinegrass, Bahiagrass, and Zoysiagrass) including Commercial St. Augustinegrass Sod Production

Apply **MOCCASIN II PLUS Herbicide before weeds emerge**. Since soil moisture is necessary for activation, irrigate with 1/2 inch of water if rainfall does not occur within 7 days after treatment (See following **Precautions**).

#### Weeds Controlled

Common Name	Scientific Name	Rate of MOCCASIN II PLUS Herbicide*
Annual sedge	<i>Cyperus compressus</i>	2.6 pts/A
Yellow nutsedge	<i>Cyperus esculentus</i>	
Smooth crabgrass	<i>Digitaria ischaemum</i>	
Large crabgrass	<i>Digitaria sanguinalis</i>	
Bearded sprangletop	<i>Leptochloa fascicularis</i>	1.3 - 2.6 pts/A
Mexican sprangletop	<i>Leptochloa uninervia</i>	
Doveweed	<i>Murdannia nudiflora</i>	
Annual bluegrass	<i>Poa annua</i>	

\* 1.0 pt/A = 0.3 ml/1,000 sq ft; 1.3 pts/A = 0.4 ml/1,000 sq ft; 2.6 pts/A = 0.9 ml/1,000 sq ft.

#### Restrictions:

- **DO NOT** use MOCCASIN II PLUS Herbicide on turfgrasses in New York State.
- **DO NOT** use on golf greens, tees, or aprons.
- Split rate of applications can be made at rates not less than 1 pt/A (0.9 lb ai/A).
- **DO NOT** apply more than 2.6 pts/A (2.4 lbs ai/A) in a single application.
- **DO NOT** apply more than once every 6 weeks.
- For commercial sod production, **DO NOT** apply more than 4.2 pts/A (4.0 lbs ai/A) per year to the same area used for sod production.
- For commercial sod production, **DO NOT** make more than 4 applications per acre per year (not to exceed 4.2 pts/A (4.0 lbs ai/A) per year).
- **DO NOT** graze or feed turf clippings to animals.

#### Precautions for all uses on sod farms:

Delayed spring green-up, temporary slowing of growth and yellowing may occur following application. To avoid turf injury:

- Application of a nitrogen-containing fertilizer at or soon after applying MOCCASIN II PLUS Herbicide will minimize delay in spring green-up and any temporary yellowing;
- Use only on turfgrass not under stress from infestations of insects, nematodes, or diseases;
- Do not seed or overseed with desirable turfgrass 4 months before or after treatment, and
- Do not apply this product to newly seeded grasses until they have overwintered and have a well-developed rhizome system.
- Before using MOCCASIN II PLUS Herbicide in the tank mix with fluid fertilizer or other registered pesticides, determine the tolerance of the turf species by applying the combination to a limited area during a period of active growth.
- In turfgrass areas which have heavy thatch, the weed control of MOCCASIN II PLUS Herbicide may be reduced.

## STORAGE AND DISPOSAL

### Pesticide Storage

This product may be stored at temperatures down to 30 degrees below 0°F.

### Pesticide Disposal

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Wastes resulting from the use of this product are toxic. Improper disposal of unused pesticide, spray mixture, or rinsate is a violation of federal law. Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to federal, state, or local procedures. For guidance in proper disposal methods, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office.

### Container Handling [equal to or less than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use and disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### Container Handling [greater than 5 gallons]

Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the person refilling. To clean container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container about 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

### Container Handling [greater than 5 gallons]

Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) or pressure rinse promptly after emptying.

Triple rinse: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

Pressure rinse: Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after flow begins to drip. Then offer for recycling if available, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures allowed by state and local authorities.



**IMPORTANT INFORMATION  
READ BEFORE USING PRODUCT  
CONDITIONS OF SALE AND LIMITATION  
OF WARRANTY AND LIABILITY**

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product reflect the opinion of experts based on field use and tests, and must be followed carefully. It is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of UPL NA Inc. or Seller. Handling, storage, and use of the product by Buyer or User are beyond the control of UPL NA Inc. and Seller. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and User, and Buyer and User agree to hold UPL NA Inc. and Seller harmless for any claims relating to such factors.

**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, UPL NA INC. AND SELLER MAKE NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ON THIS LABEL.**

To the extent consistent with applicable law, UPL NA Inc. or Seller shall not be liable for any incidental, consequential or special damages resulting from the use or handling of this product and **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF UPL NA INC. AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF UPL NA INC. OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

UPL NA Inc. and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by the duly authorized representative of UPL NA Inc.

UPL, the UPL logo, MOCCASIN, ACUMEN, BROADLOOM, INTERLINE, SATELLITE, STORM, TRICOR and ULTRA BLAZER are trademarks of a UPL Corporation Limited Group Company.

®/™ All other products are trademarks of their respective companies.

©2022 UPL Corporation Limited Group Company

ESL072221-7780-042722