

# STERGO™

## GR

Contains mefenoxam, the active ingredient used in Subdue® GR.

### Granular Fungicide

For the control of certain diseases in ornamentals, turf, nonbearing citrus, conifers, and nonbearing fruit and nut plants in nurseries

<b>ACTIVE INGREDIENT:</b>	(% by weight)
Mefenoxam* .....	1.0%
<b>OTHER INGREDIENTS:</b> .....	99.0%
<b>TOTAL:</b> .....	100.0%

\*CAS Nos. 70630-17-0 and 69516-34-3

Stergo™ GR is a granular product that contains 0.16 ounces of mefenoxam per pound of formulated product.

EPA Reg. No.: 91234-371

## KEEP OUT OF REACH OF CHILDREN CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

See below for additional Precautionary Statements.

FIRST AID	
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 - 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact SafetyCall at <b>1-844-685-9173</b> for emergency medical treatment information.	

For Chemical Emergency: Spill, Leak, Fire, Exposure, or Accident,  
Call CHEMTREC Day or Night Within USA and Canada: 1-800-424-9300 or +1 703-527-3887 (collect calls accepted)

Stergo™ GR is not manufactured, or distributed by Syngenta Professional Products, seller of Subdue® GR.

# PRECAUTIONARY STATEMENTS

## Hazards to Humans and Domestic Animals

### CAUTION

Causes moderate eye irritation. Harmful if inhaled or absorbed through skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash hands before eating, drinking, chewing gum, using tobacco or using toilet. Remove and wash contaminated clothing before reuse.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils.
- Shoes plus socks

### User Safety Requirements

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

### Engineering Controls

When handlers use closed systems or enclosed cabs in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d) (4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

### User Safety Recommendations

#### Users should:

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### Environmental Hazards

**DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment wash water.

#### Groundwater Advisory:

Mefenoxam is known to leach through soil into groundwater under certain conditions as a result of label use. This chemical may leach into groundwater if used in areas where soils are permeable, particularly where the water table is shallow.

### Physical and Chemical Hazards

**DO NOT** mix or allow coming in contact with oxidizing agent. Hazardous Chemical reaction may occur.

# DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**Stergo GR** must be used only in accordance with the directions on this label or on EPA approved supplemental labeling.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR DISEASE CONTROL OR CROP INJURY.**

**DO NOT** apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**Maximum usage when applying both metalaxyl- and mefenoxam-containing products to the same crop within the same growing year: DO NOT** apply more than the maximum yearly total application rate for the active ingredient as stated on the label of the product containing the lowest yearly total on that crop. **DO NOT** apply more than 12.3 lbs. mefenoxam per acre per year.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE); notification to workers, and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 48 hours.** Exception: If the product is soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, including plant, soil, or water is:

- Coveralls
- Chemical-resistant gloves made of barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, natural rubber ≥ 14 mils, polyethylene, polyvinyl chloride (PVC) ≥ 14 mils, or Viton® ≥ 14 mils.
- Shoes plus socks

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

**DO NOT enter or allow others to enter treated areas without protective clothing until dust has settled.**

## Product Information

**Stergo GR** is a systemic fungicide for use in ornamentals; turf; nonbearing citrus grown in nurseries and as landscape plantings; conifers grown in nurseries and plantations, including Christmas trees; and nonbearing deciduous fruit and nut plants in nurseries.

### Resistance Management

MEFENOXAM    GROUP 4    FUNGICIDE

For resistance management, **Stergo GR** contains a Group 4 phenylamide fungicide. Any fungal population may contain individuals naturally resistant to **Stergo GR** and other Group 4 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly at the same use site. Follow appropriate resistance-management strategies.

Mefenoxam is a systemic fungicide belonging to the phenylamide class of chemistry which adversely affects fungal RNA synthesis.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of **Stergo GR** or other Group 4 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with a fungicide from a different group that is equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, uses historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM guidelines for specific crop and pathogens.
- For information or to report suspected resistance contact Atticus, LLC at 984-465-4800. You can also contact your pesticide distributor or university extension specialist to report resistance.

Atticus, LLC encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label.

### Calibration Instructions

It is important to calibrate the granule applicator before use. Even after calibration, it is best to check the actual application rate under your operating conditions. Application rates can vary due to changing moisture conditions, equipment wear, normal variation in product bulk density, and particle size distribution.

Calibrating a granule applicator can be done by following these simple steps:

1. Mark off a known area, usually 1,000 sq. feet.
2. At normal walking speed, simulate an application over the 1,000 sq. ft. area and record the time taken to accomplish this.

3. Fill the applicator hopper with product. Set the applicator meter at a random setting.
4. Operate the applicator for the amount of time obtained in step 2 and collect the output of granules in a plastic bag, catch pans, or other collection device.
5. Weigh the granules collected in ounces and compare this weight with the required application rate in ounces per 1,000 sq. ft. from the product label. Repeat steps 1 - 4 until the correct meter setting is found that will deliver the required application rate.

### Application Instructions

For banded applications, calculate the amount of **Stergo GR** needed as follows:

$$\frac{\text{band width in inches}}{\text{row width in inches}} \times \frac{\text{broadcast rate}}{\text{per acre}} = \frac{\text{amount needed}}{\text{per acre}}$$

### ORNAMENTALS

**Stergo GR** is a systemic fungicide that controls diseases caused by downy mildew, fungi and damping-off, root and stem diseases caused by *Pythium* and *Phytophthora* spp. Apply **Stergo GR** to ornamentals grown in greenhouses, lath and shade-houses, outdoor field and container nurseries (including non-bearing fruit and nut plants), conifer plantations, interiorscapes and in commercial and residential landscapes. Apply **Stergo GR** as a soil surface application or as a pre-potting growing media incorporation treatment at the time of seeding and transplanting.

#### Soil and Growing Media Surface Application:

When applying **Stergo GR** to the soil, surface, irrigate with enough water to completely wet the root zone after application. **DO NOT** exceed specified rates or apply more frequently than directed, or plant injury may occur.

#### Growing Media Incorporation Treatment:

When incorporating **Stergo GR** into growing media, the mix must be dry and free of lumps to ensure thorough mixing of the granules throughout the media. When making subsequent applications at designated intervals, other registered formulations may be applied. Refer to the respective product label for rates, precautions, and limitations relative to its use.

#### Rate Selection:

Within a rate range given for a specific group of ornamentals, use lower rates for young plants with shorter crop production times and higher rates for plants with longer crop production times. Under severe disease conditions, use the highest rate and the shortest reapplication interval.

**Notice to User:** Due to the large number of species and varieties of ornamentals and nursery plants, it is impossible to test every one for tolerance to **Stergo GR**. Neither the manufacturer nor the seller has determined whether or not **Stergo GR** can be used safely on ornamentals and nursery plants not specified on this label. The professional user must determine if **Stergo GR** can be used safely prior to commercial use. In a small area, test the specified rates for a particular group of unlabeled plants, i.e., foliage plants, bedding plants, etc., for plant tolerance prior to widespread use.

Foliage Plants	Soil or Growing Media Surface Application:
Aglaonema Aphelandra Dieffenbachia Peperomia Philodendron* Pothos Schefflera Sedum Sempervivum Zygocactus	Distribute 8 - 40 oz. (0.005 - 0.025 lb. a.i.) uniformly over 1,000 sq. ft. of bed or bench area. Irrigate with enough water to thoroughly wet the root zone after application. Repeat applications at 2-to-3-month intervals, if necessary. *On Philodendron, use 12 - 60 oz. (0.008 - 0.038 lb. a.i.) per 1,000 sq. ft. <b>IMPORTANT:</b> To minimize the potential for injury to Pothos, <b>DO NOT</b> use more than 22 oz. (0.014 lb. a.i.) per 1,000 sq. ft. and <b>DO NOT</b> apply more frequently than once every 3 months.
	Growing Media Incorporation Treatment:
	Apply to growing media mix just before planting. Mix only enough for current use. <b>DO NOT</b> store. Thoroughly mix 1.6 - 8 oz. (0.001 - 0.005 lb. a.i.) per cu. yd. in growing media. After transplanting, irrigate with sufficient water to thoroughly wet the root zone. After a 2-to-3-month interval, <b>Stergo GR</b> may be reapplied as a soil surface application.
<b>Use Restrictions:</b>	
<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 60 oz. (0.038 lb. a.i.) per 1,000 sq. ft. per single application.</li> <li>• <b>DO NOT</b> apply more than 600 lbs. (6.0 lbs. a.i.) of <b>Stergo GR</b> per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).</li> <li>• <b>DO NOT</b> make more than 6 applications per year when using reduced application rates.</li> <li>• <b>Minimum Retreatment Interval:</b> 2 months</li> </ul>	

Bedding Plants	Soil or Growing Media Surface Application:
Ageratum Algerian ivy Artemisia Aster Begonia Caladium Carnation Chrysanthemum Coleus Daisy English ivy Foxglove Gaillardia Geranium Impatiens Marigold Pansy Petunia Phlox Pinks Primrose Prostrate Rosemary Salvia Snapdragon Verbena Vinca Zinnia	Uniformly apply 6 - 30 oz. (0.004 - 0.019 lb. a.i.) to 1,000 sq. ft. of bed area. For control of downy mildew use 30 oz. (0.019 lb. a.i.) per 1,000 sq. ft. Irrigate with enough water to thoroughly wet soil after application. Repeat applications at 1-to-2-month intervals, if necessary. <b>DO NOT</b> apply rates of 22 oz. (0.014 lb. a.i.) per 1,000 sq. ft. more than once every 6 weeks.
	Growing Media Incorporation Treatment:
	Apply to growing media mix just before planting. Mix only enough for current use. <b>DO NOT</b> store. Thoroughly mix 1.6 - 8 oz. (0.001 - 0.005 lb. a.i.) per cu. yd. in growing media. After transplanting, irrigate with sufficient water to thoroughly wet the root zone. <b>Stergo GR</b> may be reapplied after 6 - 8 weeks as a soil surface application.
<b>Use Restrictions:</b>	
<ul style="list-style-type: none"> <li>• <b>DO NOT</b> apply more than 30 oz. (0.019 lb. a.i.) per 1,000 sq. ft. per single application.</li> <li>• <b>DO NOT</b> apply more than 600 lbs. (6.0 lbs. a.i.) of <b>Stergo GR</b> per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).</li> <li>• <b>DO NOT</b> make more than 12 applications per year when using reduced application rates.</li> <li>• <b>Minimum Retreatment Interval:</b> 1 month</li> </ul>	

SPECIMEN

Flowers	Soil or Growing Media Surface Application:
African violet Anthurium Baby's breath Carnation Chrysanthemum Columbine Delphinium Easter lily Geranium Gloxinia Poinsettia Rose	Uniformly apply 12 - 60 oz. (0.008 - 0.04 lb. a.i.) per 1,000 sq. ft. Repeat application at 1-to-2-month intervals, if necessary. Irrigate with enough water to thoroughly wet soil after application. <b>DO NOT</b> apply rates of 37 oz. (0.0231 lb. a.i.) more often than every 6 weeks.
	Growing Media Incorporation Treatment:
	Apply to growing media mix just before planting. Mix only enough for current use. <b>DO NOT</b> store. Thoroughly add 1.6 - 8 oz. (0.001 - 0.01 lb. a.i.) per cu. yd. in growing media. After transplanting, irrigate with sufficient water to thoroughly wet the root zone. <b>Stergo GR</b> may be reapplied after 6 - 8 weeks as a soil surface application.
<b>Use Restrictions:</b>	
<ul style="list-style-type: none"> <li>- <b>DO NOT</b> apply more than 60 oz. (0.038 lb. a.i.) per 1,000 sq. ft. per single application.</li> <li>- <b>DO NOT</b> apply more than 600 lbs. (6.0 lbs. a.i.) of <b>Stergo GR</b> per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).</li> <li>- <b>DO NOT</b> make more than 12 applications per year when using reduced application rates.</li> <li>- <b>Minimum Retreatment Interval:</b> 1 month</li> </ul>	
<b>Easter Lilies:</b>	
<ul style="list-style-type: none"> <li>- For surface applications for Easter lilies, <b>DO NOT</b> apply more than 30 oz. (0.019 lb. a.i.) per 1,000 sq. ft. and only make one at-planting application.</li> <li>- For growing media incorporation treatments for Easter lilies, <b>DO NOT</b> use a pre-potting growing media mix application.</li> </ul>	

Azaleas	Soil or Growing Media Surface Application:
Azaleas	Uniformly apply 14.8 - 75 oz. (0.009 - 0.047 lb. a.i.) per 1,000 sq. ft. of bed area. Irrigate with enough water to thoroughly wet the soil after application. Repeat applications at 2-to-4-month intervals, if necessary.
	Growing Media Incorporation Treatment:
	Apply to growing media mix just before planting. Mix only enough for current use. <b>DO NOT</b> store. Thoroughly mix 1.6 - 10 oz. (0.001 - 0.006 lb. a.i.) per cu. yd. in growing media. After transplanting, irrigate with sufficient water to thoroughly wet the root zone. After a 2-to-4-month interval, <b>Stergo GR</b> may be reapplied as a soil surface application. <b>IMPORTANT:</b> (1) To minimize the potential for injury to azaleas, <b>DO NOT</b> apply repeat applications at rates greater than 75 oz. (0.047 lb. a.i.) per 1,000 sq. ft. at application intervals less than every 3 months, and <b>DO NOT</b> exceed a total of 125 oz. (0.078 lb. a.i.) per 1,000 sq. ft. in six months. (2) Use the lower rate for "Coral Bell" variety.
<b>Use Restrictions:</b>	
<ul style="list-style-type: none"> <li>- <b>DO NOT</b> apply more than 75 oz. (0.047 lb. a.i.) per 1,000 sq. ft. per single application.</li> <li>- <b>DO NOT</b> apply more than 600 lbs. (6.0 lbs. a.i.) of <b>Stergo GR</b> per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).</li> <li>- <b>DO NOT</b> make more than 6 applications per year when using reduced application rates.</li> <li>- <b>Minimum Retreatment Interval:</b> 2 months</li> </ul>	

Woody Ornamentals Other Than Azaleas	Soil and Growing Media Surface Application:
<i>Aucuba japonica</i> Arborvitae Boxwood Ceanothus Cotoneaster Dogwood Ficus "Halls" Honeysuckle Ilex <i>Juniperus</i> spp. Photinia <i>Pieris japonica</i> <i>Pinus</i> spp. Pittosporum Rhododendron White cedar White pine Yew	Uniformly apply 26 - 125 oz. (0.016 - 0.078 lb. a.i.) per 1,000 sq. ft. Irrigate with enough water to thoroughly wet the soil after application. Repeat applications at 2-to-3-month intervals, if necessary. <b>DO NOT</b> apply rates greater than 100 oz. (0.063 lb. a.i.) per 1,000 sq. ft. more frequently than every 10 weeks.
	Growing Media Incorporation Treatment:
	Apply to growing media mix just before planting. Mix only enough for current use. <b>DO NOT</b> store. Thoroughly mix 1.6 - 12.5 oz. (0.001 - 0.008 lb. a.i.) per cu. yd. in growing media. After transplanting, irrigate with sufficient water to thoroughly wet the root zone. After a 2-to-3-month interval, <b>Stergo GR</b> may be reapplied as a soil surface application.
<b>Use Restrictions:</b>	
<ul style="list-style-type: none"> <li>- <b>DO NOT</b> apply more than 125 oz. (0.078 lb. a.i.) per 1,000 sq. ft. per single application.</li> <li>- <b>DO NOT</b> apply more than 600 lbs. (6.0 lbs. a.i.) of <b>Stergo GR</b> per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).</li> <li>- <b>DO NOT</b> make more than 6 applications per year when using reduced application rates.</li> <li>- <b>Minimum Retreatment Interval:</b> 2 months</li> </ul>	

### NONBEARING CITRUS IN NURSERIES

Use **Stergo GR** on nonbearing citrus for control of citrus foot rot, root rot, and trunk canker caused by *Phytophthora* spp. Apply to the soil surface as a broadcast application under the canopy of the trees. For best efficacy, 1/2 inch of irrigation or rainfall is required within 24 hours after application. For banded applications, use the formula in the **Product Information** section of the label to calculate the amount of **Stergo GR** needed per acre of trees.

Make the first application of **Stergo GR** at the time of planting and make 2 additional applications at 3 month intervals throughout the growing season. Apply 200 lbs. (2.0 lbs. a.i.) per acre as a soil surface application broadcast or banded to the seedbeds, liners, or bedded stock. For banded applications, use a band wide enough to cover the root systems of the plants. Calculate the amount of **Stergo GR** needed for a banded treatment by using the formula in the **Product Information** section of the label.

#### Use Restrictions:

- **DO NOT** apply more than 200 lbs. (2.0 lbs. a.i.) per acre per single application.
- **DO NOT** apply more than 600 lbs. (6.0 lbs. a.i.) of **Stergo GR** per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).
- **DO NOT** make more than 3 applications per year.
- **Minimum Retreatment Interval:** 3 months
- **DO NOT** use in greenhouse citrus nursery stock intended for commercial fruit production.



## CONIFERS IN NURSERIES AND PLANTATIONS (INCLUDING CHRISTMAS TREES)

**Stergo GR** provides control of Phytophthora root rot of conifers. For best efficacy, 1/2 inch of irrigation or rainfall is required within 24 hours after application.

	Soil Surface Application:
<b>Seedbeds and Plug-Plantings</b>	Uniformly apply 6 - 30 lbs. (0.06 - 0.30 lb. a.i.) per acre (2.2 - 11 oz./1,000 sq. ft.) (0.001 - 0.007 lb. a.i./1,000 sq. ft.) in the spring and again in the fall.
<b>2-0 Transplants</b>	Uniformly apply 12 - 60 lbs. (0.12 - 0.60 lb. a.i.) per acre (4.4 - 22 oz./1,000 sq. ft.) (0.003 - 0.014 lb. a.i./1,000 sq. ft.) in the spring and again in the fall.

### Conifers in Plantations

Use of **Stergo GR** will aid in the control of Phytophthora root rot when used in conjunction with good cultural practices. The use of **Stergo GR** will not overcome poor management practices including planting on sites that are prone to flooding or are poorly drained. **Stergo GR** will not revitalize trees showing moderate to severe disease symptoms.

**Soil Surface Application:** Apply 50 - 250 lbs. (0.50 - 2.5 lbs. a.i.) of **Stergo GR** per acre (1.25 - 5.75 lbs./1,000 sq. ft.) (0.013 - 0.058 lb. a.i./1,000 sq. ft.) to the soil surface. Applications must be made in early spring before growth starts and in the fall before the ground freezes. Calculate the amount of **Stergo GR** needed for a banded treatment by using the formula in the **Product Information** section of the label.

#### Use Restrictions:

- **DO NOT** apply more than 250 lbs. (2.5 lbs. a.i.) per acre per single application (5.75 lbs./1,000 sq. ft.) (0.058 lb. a.i./1,000 sq. ft.).
- **DO NOT** apply more than 600 lbs. (6.0 lbs. a.i.) of **Stergo GR** per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).
- **DO NOT** make more than 100 applications per year when using the reduced application rate of 6 lbs. (0.06 lb. a.i.).

### NONBEARING FRUIT AND NUT PLANTS IN NURSERIES

**Stergo GR** provides control of Pythium root rot and Phytophthora root, crown, and collar rot of nonbearing deciduous fruits and nuts.

Make the first application of **Stergo GR** at the time of planting. Make additional applications at 3 month intervals throughout the growing year. For best efficacy, 1/2 inch of irrigation or rainfall is required within 24 hours after application.

**Soil Surface Application:** Apply 200 - 400 lbs. (2.0 - 4.0 lbs. a.i.) per treated acre (4.75 - 9.25 lbs./1,000 sq. ft.) (0.048 - 0.093 lb. a.i./1,000 sq. ft.) uniformly to the soil surface under the canopy of the trees. Treat sufficient surface area in nurseries to cover the root zone of the trees. Use the high rate in areas known to be infested with Phytophthora.

#### Use Restrictions:

- **DO NOT** apply more than 400 lbs. (4.0 lbs. a.i.) per treated acre per single application (9.25 lbs./1,000 sq. ft.) (0.093 lb. a.i./1,000 sq. ft.).
- **DO NOT** apply more than 600 lbs. (6.0 lbs. a.i.) of **Stergo GR** per acre per year (13.77 lbs./1,000 sq. ft.) (0.14 lb. a.i./1,000 sq. ft.).
- **DO NOT** make more than 3 applications per year when using reduced application rates.
- **Minimum Retreatment Interval:** 3 months
- **DO NOT** apply to trees that will bear harvestable fruit within 12 months of application, or illegal residues may result.

### TURF

**Stergo GR** controls Pythium blight, Pythium damping-off and yellow tuft (downy mildew) in turfgrasses. **Within the rate range given for turf, use the lower rate for the shortest interval listed and the higher rate for the longest interval. Under severe disease conditions, use the highest rate and shortest interval.**

Established Turf	Soil Surface Application:
Pythium Blight Yellow Tuft Downy Mildew	Uniformly apply as a preventative treatment at 12.5 - 25 oz. (0.008 - 0.016 lb. a.i.) per 1,000 sq. ft. Retreat at 10- to 14-day intervals during periods favorable for disease development.
Newly Seeded Areas	Soil Surface Application:
Pythium Damping-Off Pythium Blight Yellow Tuft Downy Mildew	Uniformly apply 12.5 - 25 oz. (0.008 - 0.016 lb. a.i.) per 1,000 sq. ft. immediately after seeding. <b>Irrigate with 1/4 - 1/2 inch of water after application.</b> Retreat at 7- to 14-day intervals if conditions remain favorable for disease.

#### Use Restrictions:

- **DO NOT** apply more than 25 oz. (0.016 lb. a.i.) per 1,000 sq. ft. per single application.
- **DO NOT** apply more than 270 lbs. (2.7 lbs. a.i.) per acre per year (6.2 lbs./1,000 sq. ft.) (0.062 lb. a.i./1,000 sq. ft.).
- **DO NOT** make more than 7 applications per year when using reduced application rates.
- **Minimum Retreatment Interval:** 7 days

**IMPORTANT:** To minimize the potential for insensitivity, (1) Make no more than 3 applications per growing year of any product in which the **Stergo GR** active ingredient is applied alone, and (2) Apply an alternate EPA-registered fungicide for Pythium control at least once during the season.

### Rotational Crops

**DO NOT** plant any food crop which is not registered for use with the **Stergo GR** active ingredient in soil treated with this active ingredient for a period of 12 months.

## STORAGE AND DISPOSAL

**DO NOT** contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a tightly closed container in a cool, dry place. Store in original container and out of reach of children, preferably in a locked storage area.

**PESTICIDE DISPOSAL:** Pesticide spray mixture or rinsate that cannot be used must be disposed of in a landfill approved for pesticides. Improper disposal of excess pesticide spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by the use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER HANDLING: Bag:** Nonrefillable outer bag. **DO NOT** reuse or refill the outer bag. Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## LIMITATION OF WARRANTY AND LIABILITY

**IMPORTANT: READ BEFORE USE.** Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If these terms and conditions are not acceptable, return the unopened product container at once. By using this product, user or buyer accepts the following Disclaimer of Warranties and Limitations of Liability. **CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, injury, and other unintended consequences may result because of such factors as manner of use or application (including misuse), the presence of other materials, weather conditions, and other unknown factors, all of which are beyond the control of ATTICUS, LLC. To the extent consistent with applicable law, all such risks shall be assumed by the user or buyer. **DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, ATTICUS, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond statements on this label. **LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, neither ATTICUS, LLC the manufacturer, nor the Seller shall be liable for any indirect, special, incidental or consequential damages resulting from the use, handling, application, storage, or disposal of this product. To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use, handling, application, or storage of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid.

**Stergo** is a trademark of Atticus, LLC.

Subdue® GR is a registered trademark of a Syngenta Group Company.

20240626a