



**POISON**

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm

**Tally® 800WG**

Aphicide

**ACTIVE CONSTITUENT: 800g/kg PIRIMICARB**  
(an anticholinesterase compound)

GROUP **1A** INSECTICIDE

For control of certain aphids on crops and pastures as per the Directions for Use.

**IMPORTANT: READ THE ATTACHED LEAFLET BEFORE USING THIS PRODUCT**

© Tally is a registered trademark.

[nufarm.com.au](http://nufarm.com.au)

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001

Net Contents: 1 – 20kg

APVMA Approval No.: 69374/60630

#### STORAGE AND DISPOSAL

Store in the closed, original container in a dry, cool, well ventilated area. Do not store for prolonged periods in direct sunlight. Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

#### SAFETY DIRECTIONS

Harmful if absorbed by skin contact, inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When opening the container and preparing the product for use, wear elbow-length chemical resistant gloves and goggles. If applying by aerial spraying equipment, wear gloves. If applying by high pressure hand wand, wear elbow-length chemical resistant gloves and goggles. If applying by low pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and half face piece respirator with organic vapour/gas cartridge or canister. When opening the container, preparing the product for use and applying by open-cab boomspray equipment or open-cab airblast equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves and goggles. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product or spray on skin, immediately wash area with water. If product or spray in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles, contaminated clothing and respirator, if rubber wash with detergent and warm water.

#### FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

#### MATERIAL SAFETY DATA SHEET


For further information refer to the Material Safety Data Sheet (MSDS), which can be obtained from your supplier or the Nufarm website – [www.nufarm.com.au](http://www.nufarm.com.au)

#### CONDITIONS OF SALE

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising

(including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

BN / DOM / Barcode / /



<b>CARBAMATE PESTICIDES, SOLID, TOXIC, N.O.S. (CONTAINS PIRIMICARB)</b>
<b>UN No: 2757</b>
<b>PACKING GROUP: III</b>
<b>HAZCHEM: 2X</b>
<b>IN AN EMERGENCY DIAL: 000 POLICE OR FIRE BRIGADE</b>
<b>IN AN EMERGENCY, SPECIALIST ADVICE NUFARM AUSTRALIA LIMITED 1800 033 498</b>



# POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Nufarm

# Tally® 800WG

Aphicide

**ACTIVE CONSTITUENT: 800g/kg PIRIMICARB**  
(an anticholinesterase compound)

GROUP **1A** INSECTICIDE

For control of certain aphids on crops and pastures as per the Directions for Use.

READ COMPLETE DIRECTIONS FOR USE BEFORE USING THIS PRODUCT.

nufarm.com.au

APVMA Approval No.: 69374/60630

## DIRECTIONS FOR USE

### Restrictions

DO NOT apply when the wind speed is less than 3 km/hr or more than 20km/hr.

Rate					Critical Comments	
In the following table: "Fruit Crops – Tree and Vine Crops" all rates are given for dilute spraying. For concentrate spraying, refer to the Mixing/Application Section.					For all uses in the table "Fruit Crops – Tree and Vine Crops": Apply by dilute or concentrate spraying equipment. Apply the same total amount of product to the target crop whether applying this product by dilute or concentrate spraying methods.	
<b>FRUIT CROPS – TREE AND VINE CROPS</b>						
CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			Per 100L	Per ha		
Apples	Woolly aphid	NSW, Vic, Tas, SA & WA only	30g	700g to 1.1kg	2 days	Use the higher rate when applying less than 2200L of spray/ha. Add a non-ionic surfactant such as Activator™ or Chemwet™ 1000 at the registered rate to the spray tank.
Citrus	Citrus aphid	Qld & WA only	1.5g	700g		Apply when aphids appear. Use at least 2200L spray/ha.
	Aphid			-		Apply only where pest management is being implemented and only when it is apparent that parasites and/or predators will not control the infestation within a reasonable time.
Stonefruit	Green peach aphid, Black peach aphid, Cherry aphid	All States	30g	700g		Apply at pink bud or when aphids appear. Use at least 1100L spray/ha.
<b>FRUIT CROPS – NON TREE AND VINE CROPS</b>						
CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			Per 100L	Per ha		
Blueberries	Aphids	Qld & WA only	30g	-	2 days	-
Pepinos						
Strawberries		Qld, NSW & WA only	22g			

VEGETABLES						
CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			Per 100L	Per ha		
Artichokes (globe)	Aphids	All States	30g	-	2 days	-
Asparagus						
Beetroot	Green peach aphid		-	625g		Use at least 500L spray/ha plus a non-ionic surfactant such Activator™ or Chemwet™ 1000 at the registered rate.
Beans	Cowpea aphid ( <i>Aphis craccivora</i> )	Vic, Tas & WA only	-	315g		Apply as beans emerge and thereafter whilst aphid attack continues. Spray thoroughly.
Broad beans Lima beans	Aphids	All States	30g	-		
Broccoli Brussels sprouts Cabbages Cauliflowers	Cabbage aphid		65 to 125g	315 or 625g		Apply every 10 to 14 days commencing 1 week after planting. Use 500L spray/ha. Add a non-ionic surfactant such Activator™ or Chemwet™ 1000 at the registered rate to the spray tank. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Brussels sprouts Cabbages Cauliflowers	Green peach aphid					
Chinese cabbages	Cabbage aphid, Green peach aphid					
Capsicum	Aphids					Spray when Aphids are detected and repeat if necessary. Use the higher rates if applied in conditions lower than 20°C.
Cucurbits						Apply every 5 to 10 days during periods of rapid plant growth to delay virus buildup. Use the higher rates if applied in conditions lower than 20°C.
Endive Garden cress			30g	-		-
Kale	Cabbage aphid, Green peach aphid		65 to 125g	315 to 625g		Apply every 10 to 14 days commencing 1 week after planting. Use 500 L spray/ha. Add a non-ionic surfactant such Activator™ or Chemwet™ 1000 at the registered rate to the spray tank. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rates if applied in conditions lower than 20°C.
Kiwano	Aphids					Apply every 5 to 10 days during period of rapid plant growth to delay virus buildup.
Leeks			30g	-		-
Lettuce			65 to 125g	315 to 625g		Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions lower than 20°C.
Okra			30g	-		For use in IPM programs where Helicoverpa resistance strategy is being practiced.
Peas		NSW only	65 to 125g	315 to 625g		-
Potatoes	Potato Aphid	All States	160 to 315g			Apply in 200L spray/ha when aphids first appear. If higher spray volumes per ha are used, decrease the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Radishes	Cabbage aphid, Green peach aphid		65 to 125g			Apply every 10 to 14 days commencing 1 week after planting. Use 500L spray/ha. Add a non-ionic surfactant such Activator™ or Chemwet™ 1000 at the registered rate to the spray tank. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Shallots Spinach	Aphids		30g	-		-
Swedes Turnips	Cabbage aphid, Green peach aphid		65 to 125g	315 to 625g		Apply every 10 to 14 days commencing 1 week after planting. Use 500L spray/ha. Add a non-ionic surfactant such Activator™ or Chemwet™ 1000 at the registered rate to the spray tank. If lower spray volumes per ha are used, increase the rate per 100L to maintain the equivalent rate of product per ha. Use the higher rate if temperature falls below 20°C.
Tomatoes	Aphids					Spray when aphids are detected and repeat if necessary. Use the higher rates if applied in conditions lower than 20°C.
Watercress			30g	-		-

ORNAMENTALS						
CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			Per 100L	Per ha		
Roses Chrysanthemums	Rose aphid, Chrysanthemum aphid	All States	30g	-	-	Apply when aphids appear. Spray to give complete coverage of plants.
Ornamentals	Aphids	NSW & WA only	22g			Apply when aphids appear. Spray to give complete coverage of plants.
		Qld & WA only				If applied in conjunction with predatory mites ( <i>Phytoseiulus persimilis</i> ) apply before release or only after mites have become fully established. DO NOT apply more than 2 sprays in quick succession where IPM is practiced. Where roses are grown in dense hedges and IPM is used, spray only the upper half of the hedge where the aphids are attacking the buds. Allow the lower half of the hedge to serve as a refuge for <i>P. persimilis</i> . In such situations Tally 800WG should NOT be mixed with a fungicide which requires very careful coverage. <b>NOTE:</b> Because of the wide range of crops and pests encompassed under Ornamentals, the phytotoxicity and efficacy of this compound cannot be guaranteed under all uses. Test the chemical for phytotoxicity on a small area of crop before widespread usage.
BROADACRE CROPS						
CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			Per 100L	Per ha		
Canola	Cabbage aphid ( <i>Brevicoryne brassicae</i> ), Green peach aphid ( <i>Myzus persicae</i> )	All States	-	315 or 625g	14 days harvest/ grazing	Apply when Aphids first appear using 30 to 100L water/ha (ground rig) or 20 to 50L/ha (aerial). Use higher rate if temperature falls below 20°C. Add a non-ionic surfactant such as Activator™ or Chemwet™ 1000 at the registered rate to the spray tank.
Cotton	Cotton aphid ( <i>Aphis gossypii</i> ), Green peach aphid ( <i>Myzus persicae</i> )	Qld, NSW & WA only	-	315 or 470g  <b>Aerial Application</b> Spray volume of 30 to 50L/ha  <b>Ground Application</b> Spray volume of 100L/ha minimum	21 days harvest/ grazing	Thorough spray coverage is essential to produce best results. Stage I - Apply when aphids exceed 90% plants infested Stage II - Apply when aphids exceed 95% plants infested Stage III - Apply when aphids exceed 50% plants infested, or - if 1% bolls are open and honeydew is present apply when aphids exceed 10% of plants infested.  If a spray is required for another pest, the aphid threshold for each stage should be reduced by half to determine the need to spray for aphids; e.g. 50% becomes 25% plants infested.  Use Tally 800WG no more than twice per season. Adhere to the resistant management strategy and rotate chemistry. DO NOT use Tally 800WG as consecutive sprays.
Duboisia	Aphids	Qld & WA only	30g	700g	-	Apply when populations warrant.

Lucerne Medic pastures	Spotted alfalfa aphid, Blue green aphid	All States	-	65 to 95g in 200 to 300L water	3 days grazing	<p><b>Application</b> High rate to be used when there is a high aphid population density or with dense crop growth. Complete coverage and penetration of the crop is essential.</p> <p><b>Timing</b> <b>Queensland only:</b> On seedling lucerne, treatment is necessary when 1 or 2 aphids are present per plant. Established lucerne and medics (spring and summer) – spray when there are 20 to 40 aphids per stem. <b>Spotted Alfalfa Aphid (NSW, Tas, SA, WA only):</b> Apply when aphids begin to build up on stems and repeat as necessary. <b>Spotted Alfalfa Aphid and Pea Aphid (Vic only):</b> Seedling lucerne and medics – treat when 1 or 2 aphids are present per plant. Established lucerne and medics (spring and summer) – spray when there are 20 to 40 aphids per stem as above. <b>Tasmania only:</b> A cold weather adapted species which may be found at most times of the year. Spraying should be carried out when aphids begin to increase and be repeated as necessary. For the control of both of the above lucerne and medic pasture pests Tally 800WG at the recommended rate should be applied in 200 to 300L of water/ha. Complete coverage and penetration of the crop is essential. <b>Blue Green Aphid</b> - This aphid is cooler weather tolerant and will be found at all times of the year. Spraying should be carried out as aphids are observed. In some areas Spotted Alfalfa Aphid has developed tolerance to primicarb and in those areas Tally 800WG will not give satisfactory control. However, apparent tolerance should not be confused with poor application. Consult your local Nufarm or Department of Agriculture representative for local information.</p>
	Pea aphid	Vic & WA only	-	65 to 95g		
Lupins	Green peach aphid Cowpea aphid	NSW, Vic & WA only	-	160g	6 weeks harvest/ grazing	<p>Apply when aphids first appear using 20 to 100L/ha (ground rig) or 4 to 15L/ha (aerial). Repeat as necessary. If low relative humidity conditions prevail add 1% or 10% Summer Spray Oil (DC-Tron™ or Caltex) to spray solution for ground and aerial application respectively. <b>Note:</b> For best coverage and hence best results with aerial application, apply when a slight crosswind with a minimum speed of 3km/hr is present.</p>
	Green peach aphid	WA only	-	190g		
Winter cereals	Aphids ( <i>Rhopalosiphum maidis</i> )	All States	-	95g		<p>Apply when 20 or more aphids are present per tiller and when the crop is between emergence of last leaf and flowering stages.</p> <p>Use 20 to 100L/ha (ground rig) or 4 to 15L/ha (aerial). If low relative humidity conditions prevail add Summer Spray Oil as for lupins. Where there is concern for infection by Barley Yellow Dwarf virus, spraying should be carried out as soon as aphids are located in the crop.</p>
	Aphids ( <i>R. maidis</i> , <i>R. padi</i> )	WA only	-	160 to 190g		

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

#### WITHHOLDING PERIODS

Fruit, Vegetables:

DO NOT HARVEST FOR 2 DAYS AFTER APPLICATION.

Canola:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 14 DAYS AFTER APPLICATION.

Cotton:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 21 DAYS AFTER APPLICATION.

Lucerne, Medic pastures:

DO NOT GRAZE, CUT FOR STOCK FOOD OR FEED TO POULTRY FOR 3 DAYS AFTER APPLICATION.

Lupins, Winter cereals:

DO NOT HARVEST, GRAZE OR CUT FOR STOCK FOOD FOR 6 WEEKS AFTER APPLICATION.

**GENERAL INSTRUCTIONS****INSECTICIDE RESISTANCE WARNING**

<b>GROUP</b>	<b>1A</b>	<b>INSECTICIDE</b>
--------------	-----------	--------------------

For insecticide resistance management Nufarm Tally 800WG Aphicide is a Group 1A insecticide. Some naturally occurring insect biotypes resistant to Tally 800WG and other Group 1A insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if Tally 800WG or other Group 1A insecticides are used repeatedly. The effectiveness of Tally 800WG on resistant individuals could be significantly reduced. Since occurrence of resistant individuals is difficult to detect prior to use, Nufarm Australia Ltd accepts no liability for any losses that may result from the failure of Tally 800WG to control insects. Tally may be subject to specific resistance management strategies. For further information contact your local supplier, Nufarm representative or local agricultural department agronomist.

**Resistance Management for Green Peach Aphid Control**

Some Green Peach Aphids are resistant to all commonly used insecticides including Tally 800WG. If resistance is known or suspected, DO NOT use Tally 800WG and seek advice on resistance management. If there is no Tally 800WG resistance, use a maximum of 2 non-consecutive aphid sprays per season. If Tally 800WG was used as the last aphid spray for the season DO NOT use it as the first aphid spray next season.

**MIXING**

Fill tank 50 to 75% full then start agitation. Slowly pour granules into tank and maintain agitation throughout spraying.

**APPLICATION**

If this product is applied when the temperature is less than 15°C it is essential to have thorough spray coverage for quick and total aphid control.

**Fruit Crops – Tree and Vine Crops***Dilute spraying*

- Use a sprayer designed to apply high volumes of water up to the point of run-off and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy. Apply sufficient water to cover the crop to the point of run-off. Avoid excessive run-off.
- The required water volume may be determined by applying different test volumes, using different settings on the sprayer, from industry guidelines or expert advice.
- Add the amount of product specified in the Direction for Use table for each 100L of water. Spray to the point of run-off.
- The required dilute spray volume will change and the sprayer set up and operation may also need to be changed, as the crop grows.

*Concentrate spraying*

- Use a sprayer designed and set up for concentrate spraying (that is a sprayer which applies water volumes less than those required to reach the point of run-off) and matched to the crop being sprayed.
- Set up and operate the sprayer to achieve even coverage throughout the crop canopy using your chosen water volume.
- Determine an appropriate dilute spray volume (see Dilute spraying above) for the crop canopy. This is needed to calculate the concentrate mixing rate.
- The mixing rate for concentrate spraying can then be calculated in the following way:

**EXAMPLE ONLY**

1. Dilute spray volume as determined above: for example 1500L/ha
2. Your chosen concentrate spray volume: for example 500L/ha
3. The concentration factor in this example is: 3 x (ie  $1500L \div 500L = 3$ )
4. If the dilute label rate is 30g/100 L, then the concentrate rate becomes 3 x 30, that is 90g/100L of concentrate spray.

- The chosen spray volume, amount of product per 100L of water, and the sprayer set up and operation may need to be changed as the crop grows.
- For further information on concentrate spraying, users are advised to consult relevant industry guidelines, undertake appropriate competency training and follow industry Best Practices.

**COMPATIBILITY**

Tally 800WG may be mixed in the spray vat with any one of the following products: Activator™, Chemwet™ and Betamite™. A mixture of Tally with more than one of these products or with any other product may be ineffective or may cause serious damage. The use of such a mixture is not recommended and would therefore be entirely at the user's risk.

**PRECAUTIONS**

Do not apply by low pressure hand wand applications to apples, beetroot, broccoli, Brussels sprouts, cabbages, cauliflowers, Chinese cabbages, capsicum, cucurbits, kale, Kiwano, lettuce, peas, potatoes, radishes, swedes, turnips, tomatoes, lucerne, medic pastures, lupins, canola, duboisia and winter cereals.

Do not apply by back pack/knapsack to apples, citrus, stonefruit, blueberries, pepinos, strawberries, artichokes (globe), asparagus, broad beans, lima beans, endive, garden cress, leeks, okra, shallots, spinach, watercress, beetroot, broccoli, Brussels sprouts, cabbages, cauliflowers, Chinese cabbages, capsicum, cucurbits, kale, Kiwano, lettuce, peas, potatoes, radishes, swedes, turnips, tomatoes, roses, chrysanthemums and ornamentals and all broad acre crops.

**PROTECTION OF LIVESTOCK**

Animals and poultry should be kept away from orchards and other treated areas during the horticultural spraying season. Keep animals and poultry away from spray and treated areas for a minimum of 3 days.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or waterways with the chemical or used containers.

DO NOT apply under meteorological conditions (eg crosswinds less than 3km/hr) or from spraying equipment which could be expected to cause spray drift onto adjacent areas, particularly wetlands, waterbodies or watercourses.

**STORAGE AND DISPOSAL**

Store in the closed, original container in a dry, cool, well ventilated area. Do not store for prolonged periods in direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush or puncture and deliver empty packaging for appropriate disposal to an approved waste management facility. If an approved waste management facility is not available bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers or product.

**SAFETY DIRECTIONS**

Harmful if absorbed by skin contact, inhaled or swallowed. May irritate the nose and throat. Will irritate the eyes. Avoid contact with eyes and skin. Do not inhale dust or spray mist. When opening the container and preparing the product for use, wear elbow-length chemical resistant gloves and goggles. If applying by aerial spraying equipment, wear gloves. If applying by high pressure hand wand, wear elbow-length chemical resistant gloves and goggles. If applying by low pressure hand wand, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves, goggles and half face piece respirator with organic vapour/gas cartridge or canister. When opening the container, preparing the product for use and applying by open-cab boomspray equipment or open-cab airblast equipment, wear cotton overalls, over normal clothing, buttoned to the neck and wrist and a washable hat, elbow-length chemical resistant gloves and goggles. If clothing becomes contaminated with product or wet with spray remove clothing immediately. If product or spray on skin, immediately wash area with water. If product or spray in eyes, wash it out immediately with water. Wash hands after use. After each day's use wash gloves, goggles, contaminated clothing and respirator, if rubber wash with detergent and warm water.

**FIRST AID**

If swallowed, splashed on skin or in eyes, or inhaled, contact the Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

**MATERIAL SAFETY DATA SHEET**

If additional hazard information is required refer to the Material Safety Data Sheet.

For a copy phone 1800 067 108, or visit our website at [www.nufarm.com](http://www.nufarm.com).

**In case of emergency: Phone 1800 033 498** Ask for shift supervisor. Toll free 24 hours.

**CONDITIONS OF SALE**

"Any provisions or rights under the Competition and Consumer Act 2010 or relevant state legislation which cannot be excluded by those statutes or by law are not intended to be excluded by these conditions of sale. Subject to the foregoing, all warranties, conditions, rights and remedies, expressed or implied under common law, statute or otherwise, in relation to the sale, supply, use or application of this product, are excluded. Nufarm Australia Limited and/or its affiliates ("Nufarm") shall not accept any liability whatsoever (including consequential loss), or howsoever arising (including negligence) for any damage, injury or death connected with the sale, supply, use or application of this product except for liability which cannot be excluded by statute."

Nufarm Australia Limited  
ACN 004 377 780  
103-105 Pipe Road  
Laverton North Victoria 3026  
Tel: (03) 9282 1000  
Fax: (03) 9282 1001

© Tally is a registered trademark.

™ Other trademarks.