



Trimec[®] PLATINUM

ALL-IN-ONE WEED CONTROL CONCENTRATE

ACTIVE INGREDIENTS:

2,4-D, DMA salt	5.57%
Quinclorac	2.81%
Dicamba, DMA salt	0.70%
Sulfentrazone	0.35%
Pyrimisulfan	0.25%

OTHER INGREDIENTS:90.32%
TOTAL 100.00%	

THIS PRODUCT CONTAINS:

- 0.400 lb. 2,4-D acid equivalent per gallon or 4.63%
- 0.243 lb. Quinclorac acid per gallon or 2.81%
- 0.050 lb. Dicamba acid equivalent per gallon or 0.58%
- 0.030 lb. Sulfentrazone per gallon or 0.35%
- 0.022 lb. Pyrimisulfan per gallon or 0.25%

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

First Aid	
If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p>	

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to

ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

Trimec[®] Platinum All-In-One Weed Control Concentrate is a post-emergent, selective herbicide specifically designed to target a broad spectrum of sedges, crabgrass, and broadleaf weeds. Engineered with a proprietary formulation of five active ingredients, Trimec Platinum All-In-One Weed Control Concentrate controls more than 250 listed weeds found in your lawn:

1. **broadleaf weeds**, including dandelion, clover, henbit, chickweed, and plantain
2. **grassy weeds**, including crabgrass and foxtail
3. **sedges**, including yellow and purple nutsedge

Trimec Platinum All-In-One Weed Control Concentrate may be applied to lawns containing Kentucky bluegrass, fine and tall fescues, perennial ryegrass, Bermudagrass, zoysiagrass, and bentgrass.

2. Use Restrictions

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application.
- Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants.
- Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not use on food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants (flowers, trees, shrubs, hedges, woody ornamentals, groundcovers, groundcovers established in landscape plantings).
- Do not use on Carpetgrass, Dichondra, Buffalograss, St. Augustinegrass, and turfgrass species that are not listed on this label. Do not apply this product to bentgrass mowed under 1/2 inch.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- The maximum application rate is 6.4 fl.oz. of product per 1000 sq.ft per application.
- The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.
- The maximum annual rate is 12.8 fl.oz. of product per 1000 sq.ft. excluding spot treatments.
- Do not collect grass clippings from treated turfgrass for use as mulch around plants or in compost for 3 weeks after treatment.

3. Where to Use

Trimec Platinum All-In-One Weed Control Concentrate may be used on residential lawns without harming lawn grasses when used as directed. Turfgrass species which may be treated include Kentucky Bluegrass, Perennial Ryegrass, Tall Fescue, Fine Fescue, Bentgrass, Common and Hybrid Bermudagrasses, and Zoysiagrass.

4. How Much To Apply

Over-application or rates above those specified on this label can cause turf injury.

For Kentucky Bluegrass, Fine Fescues, Tall Fescue, Perennial Ryegrass, Bentgrass, Bermudagrass, and Zoysiagrass

Spot Treatments: Mix 3.2 fl.oz. (6.4 Tbsp) of product in 1 gallon of water to treat 500 sq.ft.

Entire Lawn Treatments:

- Pump Style Sprayer: Mix 6.4 fl.oz. of product in 0.5 to 1 gallon of water to treat 1000 sq.ft.
- Hose-End Sprayer: Apply 6.4 fl.oz. of product per 1000 sq.ft.

- Pressure Sprayer [for example, 12 volt & Tow-Behind Sprayers]: Mix 6.4 fl.oz. of product in 0.5 to 1 gallon of water to treat 1000 sq.ft.

5. When To Use

Spray when broadleaf weeds are young and actively growing for best results. This product may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Use a broadcast (entire area) application, or follow-up application at a 30 day interval for more mature weeds, for dense infestations, and for adverse environmental conditions. Spot treatments during the summer may be appropriate for sparse infestations, as a follow-up treatment, or any time broadleaf weeds are susceptible.

- **Temperature:** Make broadcast applications when air temperatures are below 90°F. For spot treatments, some turf injury may be expected when air temperatures exceeds 90°F.
- **Seeding before application:** This product may be applied to newly seeded grasses approximately 6 weeks after seedling emergence or after the third mowing.
- **Seeding, sodding, sprigging, and plugging after application:** Treated areas may be reseeded 3 weeks after treatment. Apply this product 3 to 4 weeks after sodding, sprigging, or plugging operations.

6. How To Apply

6.1 Pump Style Sprayers

Ideal for spraying individual weeds

1. Add the appropriate amount of water and concentrate to the sprayer tank (based on rates in How Much to Apply section).
2. Close sprayer, shake well, and pump handle to pressurize.
3. Adjust nozzle to deliver a coarse spray pattern.
4. To broadcast over large areas: apply evenly to treated area
5. To treat individual weeds: spray to wet weed leaf surfaces.
6. Re-pressurize the sprayer as needed to maintain a good spray pattern.

6.2 Dial Style Hose-End Sprayers

Ideal for treating medium-sized lawns (ex. 1,000 – 20,000 sq.ft.)

1. Measure total area to be treated.
2. Fill sprayer with sufficient amount of product to cover the total area to be treated (based on rates in How Much to Apply section). DO NOT ADD WATER.
3. Close sprayer and set sprayer dial to 2.5 oz (rate per gallon).
4. Connect sprayer to hose
5. Extend hose to furthest point from faucet. Walking back towards the faucet, begin spraying, avoiding contact with the treated area.
6. Pour any unused product back into this container.

6.3 Pressure Sprayers

Ideal for treating large country and suburban lawns

1. To calibrate your application equipment consult your equipment owner's manual for speed and pressure settings required to deliver 0.5 to 1 gallon of spray solution to treat 1000 sq.ft. See How Much to Apply section for appropriate rate.
2. To apply evenly, drive at a constant speed that delivers the prescribed amount of spray required for the area to be treated.
3. Mix the appropriate amount of product and water in the sprayer. When mixing, first fill sprayer half full with water.
4. Add correct amount of herbicide for the intended area (based on rates in How Much to Apply section). Continue filling with water to desired level.

7. For Best Results

- **Mowing:** Delay mowing 1 day before and until 1 day after the application of this product.
- **Irrigation:** Delay watering lawn until 2 hours after treatment. Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigation (watering) before and after the application.
- **Turf injury:** Injury may result if treatment is made when turf is under stress. Injury may result if treatment is made to warm season grasses during spring transition.

8. Weeds Controlled

Broadleaf Weeds		
Amaranth, purple	False sunflower	Pineapple weed
Amaranth, slender	Falseflax	Plains coreopsis
Artichoke	Fiddleneck	(*tickseed)
Aster, heath	Field violet	Plantain, blackseed
Aster, purple	(pre-emergent)	Plantain, bracted
Aster, white heath	Filaree, redstem	Plantain, broadleaf
Aster, white prairie	Filaree, whitestem	Plantain, buckhorn
Austrian fieldcress	Fleabane, daisy	Plantain, hoary
Bedstraw, catchweed	(*annual)	Plantain, narrowleaf
Bedstraw, smooth	Fleabane, rough	Plantain, slender
Bedstraw, Northern	Galinsoga, hairy	Plantain, woolly
Bedstraw, pineywoods	Galinsoga, smallflower	Poison ivy
Beggarticks, hairy	Goldenrod	Poison oak
Betony, Florida	Ground ivy	Pokeweed
Bindweed, Field	Groundsel	Prairie Sunflower
Bindweed, Hedge	Gumweed	Prickly lettuce
Bitter Wintercress	Hairy fleabane	(*compass plant)
Bittercress, hairy	Hawkweed, meadow	Prickly sida
Bitterweed	Hawkweed, mouseear	Prostrate knotweed
Black-eyed Susan	Hawkweed, orange	Puncturevine
Black Medic	Hawkweed, yellow	Purslane, common
Blood flower milkweed	Healall	Pusley, brazil
Blue lettuce	Heartleaf drymary	Pusley, Florida
Brassbuttons	Hemp	Pusley, large flower
Bristly oxtongue	Henbit	Ragweed, bur
Broomweed	Hoary cress	Ragweed, common
Buckhorn	Horsenettle	Ragweed, lanceleaf
Bullnettle	Horseweed	Ragweed, western
Burclover	Innocence	Redstem filaree
Burdock	(*Blue-eyed Mary)	Redweed
Burweed, lawn	Jimsonweed	Rough cinquefoil
(*spurweed)	Johnny-Jump-up violet	Scarlet pimpernel
Buttercup, bulbous	Knawel	Shepard's purse
Buttercup, creeping	Knotweed, prostrate	Smartweed, ladysthumb
Buttonweed, common	Kochia	Smartweed, pale
(*poorjoe)	Lambsquarters	Smartweed,
Buttonweed, Virginia	Lespedeza	Pennsylvania
Carolina geranium	Mallow, alkali	Smooth chaff-flower
(*wild)	Mallow, bristly	Smoothdock
Carpetweed	Mallow, common	Sorrel, red (*sheep)
Catnip	Mallow, dwarf	Sowthistle, annual
Catsear, spotted (*false dandelion)	Mallow, venice	Sowthistle, spiny
Chamber bitter	Marcela	Spanish needles
Chickweed, common	Matchweed	Speedwell, bi-lobed
Chickweed, mouseear	Mexicanweed	Speedwell, common
Chickweed, sticky	Moneywort	Speedwell, corn
Chicory	Morningglory, bigroot	Speedwell, ivyleaf
Cinquefoil	Morningglory, ivyleaf	Speedwell, persian
Clover, aslison	Morningglory, mexican	Speedwell, purslane
Clover, crimson	Morningglory, red	Speedwell, slender
Clover, hop	Morningglory, tall	Speedwell, snow
Clover, rabbit foot	Morningglory, wolly	Speedwell, thymeleaf
Clover, red	Mugwort	Spiny amaranth
Clover, strawberry	Mustard, ball	Spurge, prostrate
Clover, white	Mustard, black	Spurge, spotted
Clover, white sweet	Mustard, blue	Spurweed
Clover, yellow sweet	Mustard, hedge	Star of Bethlehem
Cockle	Mustard, Indian	Strawberry, India mock
Cocklebur	Mustard, tansy	Tansy ragwort
Cocklebur, oriental	Mustard, tumble	Tanweed
Cocklebur, spiny	Mustard, white	Thistle, blessed
Common mullein	Mustard, wormseed	Thistle, bull
Corn Spurry	Nettle, stinging	Thistle, Canada
Creeping Beggarweed	(*burning)	Thistle, flodman
Creeping Jenny	Nettle, tall	(*prairie)
(*creeping charlie)	Nettle, wood	Thistle, musk
Cudweed	Old world	Thistle, platte
Cupid shaving brush	diamond flower	Thistle, plumeless
Daisy, English ¹	Oxalis (*yellow	Thistle, Russian
Daisy, oxeeye	woodsorrel &	Thistle, scotch
Dandelion, common	creeping woodsorrel)	Thistle, wavyleaf
Dandelion, false	Parsley-piert	Thistle, yellow (*horrible)
Day flower	Parsnip	Trailing crownvetch
Deadnettle	Pearlwort	Velvetleaf
Dichondra	Pennycress	Vervain, blue
Dock, broadleaf	(*frenchweed)	Vervain, hoary
Dock, curly	Pepperweed, perennial	Vervain, prostrate
Dogbane	Pepperweed, Virginia	Vervain, tall
Dogfennel	(*Peppergrass)	Vetch, bird
Dollarweed	Pigweed, prostrate	Vetch, hairy
(*Pennywort)	Pigweed, redroot	Vetch, milk
Dove weed	Pigweed, Russian	Vetch, narrowleaf
Elderberry	Pigweed, smooth	Vetch, wild
	Pigweed, tumble	(cont. on next page)

Broadleaf Weeds (cont.)		
Virginia creeper	Wild mustard	Woodsorrel, violet
Western clematis	(*yellow rocket)	Woodsorrel, yellow
Western salsify	Wild onion	(*oxalis)
Wild aster	Wild parsnip	Woolly croton
Wild buckwheat	Wild radish	Wormseed
Wild carrot	Wild rape	Yarrow, common
Wild four-o'clock	Wild strawberry	Yarrow, western
Wild garlic	Wild sweet potato	Yarrow, yellow
Wild lettuce	Wild violet ¹	Yellow rocket
Wild marigold	Woodsorrel, creeping	
Grass Weeds		
Crabgrass ^{1,2} (large and smooth)	Foxtail ¹ (green, yellow and giant)	Japanese stiltgrass ³ Signalgrass ¹ (Broadleaf)
Sedges, Kyllinga, and Rushes (including ALS resistant annual sedges)		
Yellow Nutsedge (*watergrass) Purple Nutsedge	Green Kyllinga False Green Kyllinga	Cockscomb Kyllinga Path Rush

*Synonyms

¹ Follow-up application in 30 days may be required.

² Biotypes of large and smooth crabgrass in California have shown varied response to quinclorac. If weed control is not achieved with a sequential (or follow-up) application, do not reapply this product.

³ Not for use in California

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

For questions or comments call toll-free 800.884.3179 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit: GordonsUSA.com

Gordon's® logo and Trimec® are registered trademarks of PBI-Gordon Corporation.

8977-2023 AP071423
EPA REG. NO. 2217-1078



MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.