



Trimec[®] PLATINUM

ALL-IN-ONE WEED CONTROL READY SPRAY

ACTIVE INGREDIENTS:

| | |
|---------------------------|----------------------|
| 2,4-D, DMA salt | 5.57% |
| Quinclorac | 2.81% |
| Dicamba, DMA salt | 0.70% |
| Sulfentrazone | 0.35% |
| Pyrimisulfan | 0.25% |
| OTHER INGREDIENTS: | 90.32% |
| | TOTAL 100.00% |

THIS PRODUCT CONTAINS:

- 0.400 lb. 2,4-D acid equivalent per gallon or 4.63%
- 0.243 lb. Quinclorac acid per gallon or 2.81%
- 0.050 lb. Dicamba acid equivalent per gallon or 0.58%
- 0.030 lb. Sulfentrazone per gallon or 0.35%
- 0.022 lb. Pyrimisulfan per gallon or 0.25%

KEEP OUT OF REACH OF CHILDREN CAUTION



**READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY.**

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear safety glasses. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

| First Aid | |
|---|--|
| If swallowed: | <ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a poison control center or doctor. • Do not give anything by mouth to an unconscious person. |
| If in eyes: | <ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice. |
| Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice. | |

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Rinsing application equipment over the treated area will help avoid run off to water bodies or drainage systems.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. Product Description

Trimec[®] Platinum All-In-One Weed Control Ready Spray is a post-emergent, selective herbicide specifically designed to target a broad spectrum of sedges, crabgrass, and broadleaf weeds. Engineered with a proprietary formulation of five active ingredients, Trimec Platinum All-In-One Weed Control Ready Spray controls more than 250 listed weeds found in your lawn:

1. **broadleaf weeds**, including dandelion, clover, henbit, chickweed, and plantain
2. **grassy weeds**, including crabgrass and foxtail
3. **sedges**, including yellow and purple nutsedge

Trimec Platinum All-In-One Weed Control Ready Spray may be applied to lawns containing Kentucky bluegrass, fine and tall fescues, perennial ryegrass, Bermudagrass, zoysiagrass, and bentgrass.

2. Use Restrictions

- For residential lawn use only.
- Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until sprays have dried.
- Do not apply as a fine mist because of potential for injury to desirable plants. Do not use this product on or near desirable plants, including contact of spray on exposed root systems or adventitious shoots within the drip line of desirable trees and shrubs, since injury may result.
- Do not use on food crops (gardens, fruits, and vegetables), forage crops, or ornamental plants (flowers, trees, shrubs, hedges, woody ornamentals, groundcovers, groundcovers established in landscape plantings).
- Do not use on Carpetgrass, Dichondra, Buffalograss, St. Augustine-grass, and turfgrass species that are not listed on this label. Do not apply this product to bentgrass mowed under 1/2 inch.
- Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays, and oceans. Do not apply when windy. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day. Rinse applicator over lawn area only.
- The maximum application rate is 6.4 fl.oz. of product per 1000 sq.ft per application.
- The maximum number of broadcast applications is limited to 2 per year with a minimum of 30 days between applications.
- The maximum annual rate is 12.8 fl.oz. of product per 1000 sq.ft. excluding spot treatments.
- Do not collect grass clippings from treated turfgrass for use as mulch around plants or in compost for 3 weeks after treatment.

3. Where to Use

Trimec Platinum All-In-One Weed Control Ready Spray may be used on residential lawns without harming lawn grasses when used as directed. Turfgrass species which may be treated include Kentucky Bluegrass, Perennial Ryegrass, Fine Fescues, Tall Fescue, Bentgrass, Common and Hybrid Bermudagrasses, and Zoysiagrass.

4. How Much To Apply

Over-application or rates above those specified on this label can cause turf injury.

For Kentucky Bluegrass, Fine Fescues, Tall Fescue, Perennial Ryegrass, Bentgrass, Bermudagrass, and Zoysiagrass

Entire Lawn Treatments: Apply 6.4 fl.oz. of product per 1000 sq.ft. This 32 fl.oz. container covers 5000 sq.ft. Measure the size of your yard. If the area to be treated is larger than 5000 sq.ft., you may need to buy more containers.

5. When To Use

Spray when broadleaf weeds are young and actively growing for best results. This product may be applied in the spring, summer, or fall. Spring and fall treatments under adequate soil moisture conditions are preferred to summer treatments. Generally, summer broadcast applications to older, drought stressed weeds are less effective.

Use a broadcast (entire area) application, or follow-up application at a 30 day interval for more mature weeds, for dense infestations, and for adverse environmental conditions.

- **Temperature:** Make broadcast applications when air temperatures are below 90°F. For spot treatments, some turf injury may be expected when air temperatures exceeds 90°F.
- **Seeding before application:** This product may be applied to newly seeded grasses approximately 6 weeks after seedling emergence or after the third mowing.
- **Seeding, sodding, sprigging, and plugging after application:** Treated areas may be reseeded 3 weeks after treatment. Apply this product 3 to 4 weeks after sodding, sprigging, or plugging operations.

6. How To Apply

1. Connect your garden hose to the hose-end sprayer. With your thumb, snap off the safety locking tab that prevents accidental discharge of spray solution. Leave control valve in the OFF position.
2. To BEGIN spraying: Turn on water at faucet. Extend hose to the furthest point from the faucet. Point nozzle toward the lawn to be treated and turn the control valve to the ON position. Water will automatically mix with the product. Walk backwards as you apply in order to avoid contact with the treated area. For best results, spray evenly over the measured area.
3. To STOP spraying: Turn control valve to the OFF position. Turn off water at faucet. To relieve pressure, turn control valve to ON position and wait until water stops spraying before removing sprayer from hose.

7. For Best Results

- **Mowing:** Delay mowing 1 day before and until 1 day after the application of this product.
- **Irrigation:** Delay watering lawn until 2 hours after treatment. Adequate soil moisture and favorable growing conditions enhance the performance of this product. If dry conditions exist, schedule irrigation (watering) before and after the application.
- **Turf injury:** Injury may result if treatment is made when turf is under stress. Injury may result if treatment is made to warm season grasses during spring transition.

8. Weeds Controlled

| Broadleaf Weeds | | |
|-------------------------------------|-----------------------------|-----------------------------|
| Amaranth, purple | Chamber bitter | Fiddleneck |
| Amaranth, slender | Chickweed, common | Field violet (pre-emergent) |
| Artichoke | Chickweed, mouseear | Filaree, redstem |
| Aster, heath | Chickweed, sticky | Filaree, whitestem |
| Aster, purple | Chicory | Fleabane, daisy (*annual) |
| Aster, white heath | Cinquefoil | Fleabane, rough |
| Aster, white prairie | Clover, alsike | Galinsoga, hairy |
| Austrian fieldcress | Clover, crimson | Galinsoga, smallflower |
| Bedstraw, catchweed | Clover, hop | Goldenrod |
| Bedstraw, smooth | Clover, rabbit foot | Ground ivy |
| Bedstraw, Northern | Clover, red | Groundsel |
| Bedstraw, pineywoods | Clover, strawberry | Gumweed |
| Beggarticks, hairy | Clover, white | Hairy fleabane |
| Betony, Florida | Clover, white sweet | Hawkweed, meadow |
| Bindweed, Field | Clover, yellow sweet | Hawkweed, mouseear |
| Bindweed, Hedge | Cockle | Hawkweed, orange |
| Bitter Wintercress | Cocklebur | Hawkweed, yellow |
| Bittercress, hairy | Cocklebur, oriental | Healall |
| Bitterweed | Cocklebur, spiny | Heartleaf drymary |
| Black-eyed Susan | Common mullein | Hemp |
| Black Medic | Corn Spurry | Henbit |
| Blood flower milkweed | Creeping Beggarweed | Hoary cress |
| Blue lettuce | Creeping Jenny | Horsenettle |
| Brassbuttons | (*creeping charlie) | Horseweed |
| Bristly oxtongue | Cudweed | Innocence (*Blue-eyed Mary) |
| Broomweed | Cupid shaving brush | Jimsonweed |
| Buckhorn | Daisy, English ¹ | Johnny-Jump-up violet |
| Bullnettle | Daisy, oxeye | Knawel |
| Burclover | Dandelion, common | Knotweed, prostrate |
| Burdock | Dandelion, false | Kochia |
| Burweed, lawn (*spurweed) | Day flower | Lambsquarters |
| Buttercup, bulbous | Deadnettle | Lespedeza |
| Buttercup, creeping | Dichondra | Mallow, alkali |
| Buttonweed, common (*poorjoe) | Dock, broadleaf | Mallow, bristly |
| Buttonweed, Virginia | Dock, curly | Mallow, common |
| Carolina geranium (*wild) | Dogbane | Mallow, dwarf |
| Carpetweed | Dogfennel | Mallow, venice |
| Catnip | Dollarweed (*Pennywort) | Marcela |
| Catsear, spotted (*false dandelion) | Dove weed | Matchweed |
| | Elderberry | (cont. on next page) |
| | False sunflower | |
| | Falseflax | |

| Broadleaf Weeds (cont.) | | |
|--|--|--------------------------------------|
| Mexicanweed | Poison ivy | Thistle, Canada |
| Moneywort | Poison oak | Thistle, flodman (*prairie) |
| Morningglory, bigroot | Pokeweed | Thistle, musk |
| Morningglory, ivyleaf | Prairie Sunflower | Thistle, platte |
| Morningglory, mexican | Prickly lettuce (*compass plant) | Thistle, plumeless |
| Morningglory, red | Prickly sida | Thistle, Russian |
| Morningglory, tall | Prostrate knotweed | Thistle, scotch |
| Morningglory, wolly | Puncturevine | Thistle, wavyleaf |
| Mugwort | Purslane, common | Thistle, yellow (*horrible) |
| Mustard, ball | Pusley, brazil | Trailing crownvetch |
| Mustard, black | Pusley, Florida | Velvetleaf |
| Mustard, blue | Pusley, large flower | Vervain, blue |
| Mustard, hedge | Ragweed, bur | Vervain, hoary |
| Mustard, Indian | Ragweed, common | Vervain, prostrate |
| Mustard, tansy | Ragweed, lanceleaf | Vervain, tall |
| Mustard, tumble | Ragweed, western | Vetch, bird |
| Mustard, white | Redstem filaree | Vetch, hairy |
| Mustard, wormseed | Redweed | Vetch, milk |
| Nettle, stinging (*burning) | Rough cinquefoil | Vetch, narrowleaf |
| Nettle, tall | Scarlet pimpernel | Vetch, wild |
| Nettle, wood | Shepard's purse | Virginia creeper |
| Old world diamond flower | Smartweed, ladysthumb | Western clematis |
| Oxalis (*yellow woodsorrel & creeping woodsorrel) | Smartweed, pale | Western salsify |
| Parsley-piert | Smartweed, Pennsylvania | Wild aster |
| Parsnip | Smooth chaff-flower | Wild buckweed |
| Pearlwort | Smoothdock | Wild carrot |
| Pennycress (*frenchweed) | Sorrel, red (*sheep) | Wild four-o'clock |
| Pepperweed, perennial | Sowthistle, annual | Wild garlic |
| Pepperweed, Virginia (*Peppergrass) | Sowthistle, spiny | Wild lettuce |
| Pigweed, prostrate | Spanish needles | Wild marigold |
| Pigweed, redroot | Speedwell, bi-lobed | Wild mustard (*yellow rocket) |
| Pigweed, Russian | Speedwell, common | Wild onion |
| Pigweed, smooth | Speedwell, corn | Wild parsnip |
| Pigweed, tumble | Speedwell, ivyleaf | Wild radish |
| Pineapple weed | Speedwell, persian | Wild rape |
| Plains coreopsis (*tickseed) | Speedwell, purslane | Wild strawberry |
| Plantain, blackseed | Speedwell, slender | Wild sweet potato |
| Plantain, bracted | Speedwell, snow | Wild violet ¹ |
| Plantain, broadleaf | Speedwell, thymeleaf | Woodsorrel, creeping |
| Plantain, buckhorn | Spiny amaranth | Woodsorrel, violet |
| Plantain, hoary | Spurge, prostrate | Woodsorrel, yellow (*oxalis) |
| Plantain, narrowleaf | Spurge, spotted | Woolly croton |
| Plantain, slender | Spurweed | Wormseed |
| Plantain, woolly | Star of Bethlehem | Yarrow, common |
| | Strawberry, India mock | Yarrow, western |
| | Tansy ragwort | Yarrow, yellow |
| | Tanweed | Yellow rocket |
| | Thistle, blessed | |
| | Thistle, bull | |
| Grass Weeds | | |
| Crabgrass ^{1,2} (large and smooth) | Foxtail ¹ (green, yellow and giant) | Japanese stiltgrass ³ |
| | | Signalgrass ¹ (Broadleaf) |
| Sedges, Kyllinga, and Rushes (including ALS resistant annual sedges) | | |
| Yellow Nutsedge (*watergrass) | Green Kyllinga | Cockscomb Kyllinga |
| Purple Nutsedge | False Green Kyllinga | Path Rush |

¹Synonyms

¹ Follow-up application in 30 days may be required.

² Biotypes of large and smooth crabgrass in California have shown varied response to quinlorac. If weed control is not achieved with a sequential (or follow-up) application, do not reapply this product.

³ Not for use in California

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Keep from freezing.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

LIMITED WARRANTY AND DISCLAIMER

FOR USE ONLY AS DIRECTED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE HEREBY EXPRESSLY DISCLAIMED. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO CASE SHALL THE MANUFACTURER BE LIABLE FOR INCIDENTAL, CONSEQUENTIAL, OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. If these terms are not acceptable, return this product unopened immediately to the point of purchase, and the purchase price will be refunded in full. The terms of this LIMITED WARRANTY AND DISCLAIMER cannot be varied by any written or verbal statements or agreements at the point of sale or elsewhere.

For questions or comments call toll-free 800.884.3179 Mon.-Fri. 8:00 a.m. – 4:30 p.m. CST or visit: GordonsUSA.com

Gordon's® logo and Trimec® are registered trademarks of PBI-Gordon Corporation.

89777-2023 AP071423
EPA REG. NO. 2217-1078



MANUFACTURED BY
PBI/GORDON CORPORATION
P.O. BOX 860350
SHAWNEE, KANSAS 66286
GordonsUSA.com

ATTENTION: This specimen label is provided for informational use only. This product may not yet be available for sale in your state or area. The information found in this label may differ from the information found on the product label you are using. Always follow the instructions for use and precautions on the label of the product you are using.