

PYRAFLUFEN-ETHYL GROUP 14 HERBICIDE



VENUE[®] MAX

HERBICIDE

ACTIVE INGREDIENT:

Pyraflufen-ethyl: Acetic acid, [2-chloro-5-[4-chloro-5-(difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-4-fluorophenoxy]-, ethyl ester 4.0%

OTHER INGREDIENTS: 96.0%

TOTAL **100.0%**

Contains 0.35 lb pyraflufen-ethyl per gallon

(CB)

(E)

(AYE)

EPA Reg. No. 71711-69

EPA Est. No. 70815-GA-002

39578-TX-1

54675-MEX-001

superscript corresponds to lot number

KEEP OUT OF REACH OF CHILDREN CAUTION

See inside booklet for First Aid, Precautionary Statements, and Directions for Use

NET CONTENTS: 1 quart

 **NICHINO**

920102 02/24

FIRST AID

If on skin or clothing

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product, including human health concerns and medical emergencies, call 1-800-348-5832. In case of fire or spills, information may be obtained by calling 1-800-424-9300.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using the toilet.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, and/or Viton™ \geq 14 mils

User Safety Requirements

Follow the manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

ENGINEERING CONTROLS

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

User Safety Recommendations

Users should:

- Wear long sleeved shirt and long pants, shoes plus socks, and chemical and/or water-resistant gloves.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. This product may contaminate water through drift of spray in wind or via runoff events. Use care when applying in areas adjacent to any body of water. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate. **DO NOT** apply when weather conditions favor drift from treated areas.

Nontarget Organism Advisory

This product is toxic to plants and may adversely impact the forage and habitat of nontarget organisms, including pollinators, in areas adjacent to the treated site. Protect the forage and habitat of nontarget organisms by following label directions intended to minimize spray drift.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the state or tribal agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

For early entry into treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated including plants, soil, or water, wear:

- Coveralls
- Shoes plus socks
- Chemical-resistant gloves made of barrier laminate, butyl rubber \geq 14 mils, nitrile rubber \geq 14 mils, neoprene rubber \geq 14 mils, natural rubber \geq 14 mils, polyethylene, polyvinyl chloride (PVC) \geq 14 mils, and/or Viton \geq 14 mils

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For other uses, including interiorscapes and other nonagricultural uses, **DO NOT** enter treated areas without protective clothing until sprays have dried.

PRODUCT INFORMATION

VENUE[®] MAX Herbicide is a contact herbicide and requires thorough coverage for complete broadleaf weed control.

VENUE MAX Herbicide must be tank mixed with another foliar active broadleaf herbicide for complete control of most broadleaf weeds.

DO NOT apply **VENUE MAX** Herbicide through any type of irrigation system.

VENUE MAX Herbicide is rainfast one hour after application.

ROTATIONAL CROP RESTRICTIONS

Crop/Crop Group	Rotational/Plantback Intervals
<p> Corn Cottonseed Subgroup (Crop Subgroup 20C) Hops Pome Fruit (Crop Group 11-10) Pomegranate Small Fruit Vine Climbing Subgroup Except Fuzzy Kiwifruit (Crop Group 13-07F) Soybean Stone Fruit (Crop Group 12-12) Tree Nut (Crop Group 14-12) Triticale; Wheat Tropical and Subtropical, Small Fruit, Edible Peel (Crop Subgroup 23A) Tuberous and Corm Vegetables (Crop Subgroup 1C) </p>	<p>0 days following application</p>
<p> Brassica (Cole) Leafy Vegetables (Crop Group 5) Bulb Vegetables (Crop Group 3) Cereal Grains (Crop Group 15 except corn, wheat, triticale); see 0-day plantback interval above Cucurbit Vegetables (Crop Group 9) Fruiting Vegetables (Crop Group 8-10) Leafy Vegetables (Crop Group 4-16) Legume Vegetables, Succulent or Dried (Crop Group 6 except soybean); see 0-day plantback interval above </p>	<p>1 day following preplant burndown application</p>

ROTATIONAL CROP RESTRICTIONS *(continued)*

Crop/Crop Group	Rotational/Plantback Intervals
Oilseed (Crop Group 20 except Cottonseed Crop Subgroup 20C) see 0-day plantback interval above Root and Tuber Vegetables (Crop Group 1 except Tuberos and Corm Vegetables Crop Subgroup 1C); see 0-day plantback interval above Sugarcane	1 day following preplant burndown application
All Other Rotational Crops	DO NOT plant for 30 days following the last application of VENUE MAX Herbicide.

WEEDS CONTROLLED

The following broadleaf weed species can be controlled or suppressed up to 4 inches in height or less or rosettes of 3 inches in diameter or less. Tank mixtures of **VENUE MAX** Herbicide with other labeled broadleaf herbicides may be needed for control of some weed species. Control may be reduced with weeds larger than 4 inches in height or 3 inches in diameter.

(continued)

Alkaliweed* Amaranth, Palmer* Bedstraw Beggartick, hairy Beggarweed, Florida Bindweed, field Buckwheat, wild Canola Carpetweed Celery, wild Chickweed Clover, white Cocklebur	Cotton, volunteer (conventional, GMO varieties) Dandelion, common Dock, curly Dollarweed Eclipta Eveningprimrose, cutleaf Fleabane* Geranium, Carolina Henbit Horsenettle* Kochia	Ladysthumb Lambsquarters, common Lettuce, prickly Mallow, common Malva spp. Marestail* Milkthistle Morningglory species Mustard, wild* Nettle, stinging Nightshade, black Nightshade, silverleaf Panicle willowweed	Pigweed, redroot Pigweed, smooth Pineapple-weed Poinsettia, wild Poison-ivy Potato, volunteer Prickly sida (teaweed) Purslane, common Radish, wild Ragweed, common Ragweed, giant Redmaid Rocket, London Sesbania, hemp	Smartweed, Pennsylvania Smellmelon Sowthistle, annual Spurge, leafy Sunflower, common Tansymustard, Western Thistle, Canada Thistle, Russian Toadflax, Dalmatian Velvetleaf Virginia-creeper Waterhemp, common Waterhemp, tall
*suppression				

WEED RESISTANCE MANAGEMENT

For resistance management, **VENUE MAX** Herbicide is a Group 14 herbicide. Any weed population may contain or develop plants naturally resistant to **VENUE MAX** Herbicide and other Group 14 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Appropriate resistance management strategies must be followed.

Proactively implement diversified weed control strategies to minimize selection for weed populations resistant to one or more herbicides. A diversified weed management program may include the use of multiple herbicides with different modes of action and overlapping weed spectrum with or without tillage operations and/or other cultural practices. Research has demonstrated that using the labeled rate and application instructions is important to delay the selection for resistance. Scouting after a herbicide application is important because it can facilitate the early identification of weed shifts and/or weed resistance and thus provide direction on future weed management practices. One of the best ways to contain resistant populations is to implement measures to avoid allowing weeds to reproduce by seed or to proliferate vegetatively. Cleaning equipment between sites and avoiding movement of plant material between sites will greatly aid in retarding the spread of resistant weed seed.

To delay herbicide resistance, take one or more of the following steps:

- Rotate the use of **VENUE MAX** Herbicide or other Group 14 herbicides within a growing season sequence or among growing seasons with different herbicide groups that control the same weeds in a field.
- Use tank mixtures with herbicides from a different group if such use is permitted; where information on resistance in target weed species is available, use the less resistance-prone partner at a rate that will control the target weed(s) equally as well as the more resistance-prone partner. Consult your local extension service or certified crop advisor if you are unsure as to which active ingredient is currently less prone to resistance.
- Adopt an integrated weed management program for herbicide use that includes scouting and uses historical information related to herbicide use and crop rotation and that considers tillage (or other mechanical control methods), cultural (e.g., higher crop seeding rates; precision fertilizer application method and timing to favor the crop and not the weeds), biological (weed-competitive crops or varieties), and other management practices.
- Scout after herbicide application to monitor weed populations for early signs of resistance development. Indicators of possible herbicide resistance include: (1) failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds; (2) a spreading patch of noncontrolled plants of a particular weed species; (3) surviving plants mixed with controlled individuals of the same species. If resistance is suspected, prevent weed seed production in the affected area by an alternative herbicide from a different group or by a mechanical method (for example, hoeing or tillage). Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment when moving between fields and planting clean seed.
- If a weed pest population continues to progress after treatment with this product, discontinue use of this product, and switch to another management strategy or herbicide with a different mode of action, if available.
- Contact your local extension specialist or certified crop advisors for additional pesticide resistance management and/or integrated weed management recommendations for specific crops and weed biotypes.
- Report lack of performance to registrant or their representative.

Suspected herbicide resistant weeds may be identified by these indicators:

- Failure to control a weed species normally controlled by the herbicide at the dose applied, especially if control is achieved on adjacent weeds;
- A spreading patch of noncontrolled plants of a particular weed species; and
- Surviving plants mixed with controlled individuals of the same species.

Contact your local sales representative, certified crop advisor or pest control advisor, or extension agent to find out if suspected

resistant weeds to this MOA have been found in your region. If resistant biotypes of target weeds have been reported, use the application rates of this product specified for your local conditions. Tank mix products so that there are multiple effective mechanisms of action for each target weed.

MANDATORY SPRAY DRIFT MANAGEMENT

Aerial Applications

- **DO NOT** release spray at a height greater than 10 feet above the ground or vegetative canopy unless a greater application height is necessary for pilot safety.
- Applicators are required to use a fine or coarser droplet size (ASABE S572.1).
- The boom length must not exceed 75% of the wingspan for airplanes or 90% of the rotor blade diameter for helicopters.
- Applicators must use ½ swath displacement upwind at the downwind edge of the field.
- Nozzles must be oriented so the spray is directed toward the back of the aircraft.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Ground Boom Applications

- User must only apply with the release height specified by the manufacturer but no more than 3 feet above the ground or crop canopy.
- Applicators are required to use fine or coarser droplet size (ASABE S572.1).
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

Boomless Ground Applications

- Applicators are required to use fine or coarser droplet size (ASABE S572.1) for all applications.
- **DO NOT** apply when wind speeds exceed 10 miles per hour at the application site.
- **DO NOT** apply during temperature inversions.

SPRAY DRIFT ADVISORIES

THE APPLICATOR IS RESPONSIBLE FOR AVOIDING OFF-SITE SPRAY DRIFT.

BE AWARE OF NEARBY NONTARGET SITES AND ENVIRONMENTAL CONDITIONS.

Importance of Droplet Size

An effective way to reduce spray drift is to apply large droplets. Use the largest droplets that provide target pest control. While applying larger droplets will reduce spray drift, the potential for drift will be greater if applications are made improperly or under unfavorable environmental conditions.

Controlling Droplet Size - Ground Boom

- Volume - Increasing the spray volume so that larger droplets are produced will reduce spray drift. Use the highest practical spray volume for the application. If a greater spray volume is needed, consider using a nozzle with a higher flow rate.
- Pressure - Use the lowest spray pressure directed for the nozzle to produce the target spray volume and droplet size.
- Spray Nozzle - Use a spray nozzle that is designed for the intended application. Consider using nozzles designed to reduce drift.

Controlling Droplet Size - Aircraft

Adjust Nozzles - Follow nozzle manufacturers' specifications for setting up nozzles. To reduce fine droplets, orient nozzles parallel with the airflow in flight.

Boom Height - Ground Boom

Use the lowest boom height that is compatible with the spray nozzles that will provide uniform coverage. For ground equipment, the boom must remain level with the crop and have minimal bounce.

Release Height - Aircraft

Higher release heights increase the potential for spray drift. When applying aerially to crops, **DO NOT** release spray at a height greater than 10 feet above the crop canopy unless a greater application height is necessary for pilot safety.

Shielded Sprayers

Shielding the boom or individual nozzles can reduce spray drift. Consider using shielded sprayers. Verify that the shields are not interfering with the uniform deposition of the spray on the target area.

Temperature and Humidity

When making applications in hot and dry conditions, use larger droplets to reduce effects of evaporation.

Temperature Inversions

Drift potential is high during a temperature inversion. Temperature inversions are characterized by increasing temperature with altitude and are common on nights with limited cloud cover and light to no wind. The presence of an inversion can be indicated by ground fog or by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing. Avoid applications during temperature inversions.

Wind

Drift potential increases with wind speed. **AVOID APPLICATIONS DURING GUSTY WIND CONDITIONS.** Applicators need to be familiar with local wind patterns and terrain that could affect spray drift.

Boomless Ground Applications

Setting nozzles at the lowest effective height will help to reduce the potential for spray drift.

Handheld Technology Applications

Take precautions to minimize spray drift.

TANK MIXTURES

It is the pesticide user's responsibility to ensure that all products are registered for the intended use. Read and follow the applicable restrictions and limitations and application instructions on all product labels involved in tank mixing. Users must follow the most restrictive application instructions and precautionary statements of each product in the tank mixture.

VENUE MAX Herbicide may be applied as a tankmix or in sequential application with other herbicide, fungicide, or insecticide products. Weather, crop conditions, or the presence of certain weeds, crop damaging insects, or diseases will indicate the inclusion of other pesticides in the application.

Note: Test compatibility of **VENUE MAX** Herbicide in any tankmix combination before use. To determine the physical compatibility with other products, use a jar test, as described below:

Using a quart jar, add the proportionate amounts of the products to 1 quart of water. Add wettable powders and water-dispersible granular products first, then liquid flowables, and emulsifiable concentrates last. After thoroughly mixing, let stand for at least 5 minutes. If the combination remains mixed or can be remixed readily, it is physically compatible. Once compatibility has been proven, use the same procedure for adding required ingredients to the spray tank.

Read and follow all label directions for each tankmix product. Always use in accordance with the most restrictive of label precautions and limitations.

MIXING DIRECTIONS

VENUE MAX Herbicide Alone: Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application, and then turn on agitation. Pour the specified amount of product on the surface of the water in the spray tank. Add the remaining water volume to the spray tank with agitation running. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

VENUE MAX Herbicide in Tank Mixtures: Begin with clean equipment. Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application, and turn on agitation. If using a buffering agent, add after filling the tank with $\frac{3}{4}$ amount of water. Add the specified amount of tankmix products in the following order while maintaining agitation:

- 1) products in water-soluble packets
- 2) wettable powders
- 3) water-dispersible granulars and/or soluble powders
- 4) flowable liquids (including **VENUE MAX** Herbicide)
- 5) emulsifiable concentrates
- 6) adjuvants and/or oils
- 7) remaining amount of water to achieve the desired level

Always follow the labeled mixing instructions of any partner products. Keep agitation running during filling and spraying operations. If spraying must be stopped before emptying the sprayer, resume agitation before spraying the remainder of the load. Mix only as much spray solution as can be sprayed within four hours. Storage and use of the previous day's spray mix may result in reduced activity.

Use an approved agricultural buffering agent, buffering to pH 7.5 or less if using **VENUE MAX** Herbicide in a water source greater than or equal to pH 7.5. Always buffer the water source BEFORE adding **VENUE MAX** Herbicide to the spray tank.

SPOT TREATMENT

For spot treatment to listed broadleaf weeds or for sucker management, refer to the information below to determine the amount of **VENUE MAX** Herbicide to add to a tank. Spray using a pressure (pump-up) sprayer (or similar application equipment) until

wet but prior to runoff. Use information for rates, concentrations, water volumes, and timing and frequency of application can be found in the **Rate/Acre** and **Application Instructions** columns in the **Crop Use Directions** tables. Please refer to and follow all precautions and restrictions under **Application Instructions** for the crop to be treated.

Fluid Ounces of VENUE MAX Herbicide to Add to Sprayer Tank			
Sprayer tank capacity	Spray volume (gallons/A)	fl oz VENUE MAX to add to spray tank for an application rate of	
		0.5 fl oz/A (0.0013 lb ai/A)	1.9 fl oz/A (0.0053 lb ai/A)
1	20	0.03	0.10
3	20	0.08	0.29
5	20	0.13	0.48
10	20	0.25	0.95

Formula
 Fluid Ounce **VENUE MAX** Herbicide to add to sprayer tank =

$$\frac{\text{Application Rate} \times \text{Sprayer Tank Capacity}}{\text{Spray Volume}}$$

(continued)

**Fluid Ounces of VENUE MAX Herbicide
to Add to Sprayer Tank** *(continued)*

Example Calculation for 1 gallon sprayer tank capacity

Fluid Ounce **VENUE MAX** Herbicide to add to sprayer tank =
1.9 fl oz/A x 1 gallon

$$20 \text{ gallons/A} = 0.10 \text{ fl oz}$$

where: Application rate = 1.9 fl oz/A
Sprayer tank capacity = 1 gallon
Spray volume = 20 gallons/A

Example Calculation for 5 gallon sprayer tank capacity

Fluid Ounce **VENUE MAX** Herbicide to add to sprayer tank =
1.9 fl oz/A x 5 gallons

$$20 \text{ gallons/A} = 0.48 \text{ fl oz}$$

where: Application rate = 1.9 fl oz/A
Sprayer tank capacity = 5 gallons
Spray volume = 20 gallons/A

EQUIPMENT CLEANING

DO NOT allow the spray solution to dry in the application equipment. After application and before using the sprayer equipment for any other applications, the sprayer must be thoroughly cleaned. Applicators must ensure proper equipment clean-out for any other products mixed with **VENUE MAX** Herbicide as provided on the other product label(s). Immediately following application, clean all equipment thoroughly with detergent or a spray tank cleaner and water as described below. Residues of **VENUE MAX** Herbicide remaining in inadequately cleaned equipment and released in subsequent applications can cause injury to crops.

1. Drain sprayer tank, hoses, and spray boom, and thoroughly rinse with clean water the inside of the spray tank, sprayer hoses, boom, and nozzles to remove any sediment or residues.
2. Fill the tank ½ full with clean water, add the appropriate detergent (follow manufacturer's directions for use). Fill tank to capacity, and operate the sprayer with agitation for 15 minutes to flush hoses, boom, and nozzles.
3. Drain the sprayer tank, lines, and booms. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray nozzles, tips, and screens.
4. Dispose of all cleaning solutions, rinsate, and washwaters in accordance with federal, state, and local regulations.

CROP USE DIRECTIONS

Nonbearing Only			
Date; Feijoa; Fig; Kiwifruit; Mango; Persimmon			
Application	Pest	Rate/Acre	Applications Per Year
In-Season	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed a combined total of 2 applications of VENUE MAX Herbicide per year for these uses.
	Sucker Management		
Postharvest Dormant Prebloom	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed 3 applications of VENUE MAX Herbicide per year for this use.
	Sucker Management		DO NOT exceed 2 applications of VENUE MAX Herbicide per year for this use.

(continued)

CROP USE DIRECTIONS *(continued)*

Bearing and Nonbearing

Date; Feijoa; Fig; Kiwifruit; Mango; Persimmon

Application Instructions

- Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds and sucker growth.
- Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions.
- COC adjuvants are advised, although other adjuvants may be used.
- Avoid contact with green, uncallused bark of young trees or vines established less than one year unless protected from spray contact by non-porous wraps, grow tubes, or waxed containers.
- Use the higher rate for hard-to-control weeds.
- For the management of undesirable sucker growth on the basal portion of trunks, root sprouts, and tree/vine trunks, growth must be controlled when the tissue is young, immature, and/or not hardened off.

USE RESTRICTIONS

- **DO NOT** apply by air for this use.
- **DO NOT** allow spray to drift onto desirable fruit, foliage, vines, or trees as damage will occur.
- **DO NOT** exceed 1.9 fl oz (0.0053 lb ai) of **VENUE MAX** Herbicide per acre in a single application for these uses.
- **DO NOT** exceed 3.3 fl oz (0.0090 lb ai) of **VENUE MAX** Herbicide per acre per year for all in-season applications combined.
- **DO NOT** exceed 3.3 fl oz (0.0090 lb ai) of **VENUE MAX** Herbicide per acre per year for all postharvest, dormant, and pre-bloom applications combined.
- Allow a minimum of 30 days between applications for this use.

(continued)

CROP USE DIRECTIONS *(continued)*

Bearing and Nonbearing

Pome Fruit Group (Crop Group 11-10) apple; azarole; crabapple; loquat; mayhaw; medlar; pear; pear, Asian; quince; quince, Chinese; quince, Japanese; tejocote; cultivars, varieties, and/or hybrids of these

Pomegranate

Small Fruit Vine Climbing Subgroup Except Fuzzy Kiwifruit (Crop Group 13-07F) Amur river grape; gooseberry; grape; kiwifruit, hardy; Maypop; schisandra berry; cultivars, varieties, and/or hybrids of these

Stone Fruit Group (Crop Group 12-12) apricot; apricot, Japanese; capulin; cherry, black; cherry, Nanking; cherry, sweet; cherry, tart; Jujube, Chinese; nectarine; peach; plum; plum, American; plum, beach; plum, Canada; plum, cherry; plum, Chick-asaw; plum, Damson; plum, Japanese; plum, Klamath; plum, prune; plumcot; sloe; cultivars, varieties, and/or hybrids of these

Tree Nut Group (Crop Group 14-12) African nut-tree; almond; beechnut; Brazil nut; Brazilian pine; bunya; bur oak; butternut; Cajou nut; candlenut; cashew; chestnut; chinquapin; coconut; coquito nut; dika nut; ginkgo; Guiana chestnut; hazelnut (Filbert); heartnut; hickory nut; Japanese horse-chestnut; macadamia nut; mongongo nut; monkey-pot; monkey puzzle nut; Okari nut; Pachira nut; peach palm nut; pecan; pequi; Pili nut; pine nut; pistachio; Sapucaia nut; tropical almond; walnut, black; walnut, English; yellowhorn; cultivars, varieties, and/or hybrids of these

Tropical and Subtropical, Small Fruit, Edible Peel Subgroup (Crop Subgroup 23A) acerola; African plum; agritos; al-mondette; appleberry; arbutus berry; bayberry, red; bignay; breadnut; cabeluda; carandas-plum; Ceylon iron wood; Ceylon olive; cherry-of-the-Rio-Grande; Chinese olive, black; Chinese olive, white; chirauli-nut; cocoplum; desert-date; false sandal-wood; fragrant manjack; gooseberry, abyssinian; gooseberry, Ceylon; gooseberry, otaheite; governor's plum; grumichama; guabiroba; guava berry; guava, Brazilian; guava, Costa Rican; guayabillo; illawarra plum; Indian-plum; Jamaica-cherry; jambolan; kaffir-plum; kakadu plum; kapundung; karanda; lemon aspen; mombin, yellow; monos plum; mountain cherry; olive; persimmon, black; pitomba; plum-of-Martinique; rukam; rumberry; sea grape; sete-capotes; silver aspen; water apple; water pear; water berry; wax jambu; cultivars, varieties, and hybrids of these commodities

(continued)

CROP USE DIRECTIONS *(continued)*

Bearing and Nonbearing
Pome Fruit Group (Crop Group 11-10); Pomegranate; Small Fruit Vine Climbing Subgroup Except Fuzzy Kiwifruit (Crop Group 13-07F); Stone Fruit Group (Crop Group 12-12); Tree Nut Group (Crop Group 14-12); Tropical and Subtropical, Small Fruit, Edible Peel Subgroup (Crop Subgroup 23A)

Application	Pest	Rate/Acre	Applications Per Year
Postharvest Dormant Prebloom	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed 3 applications of VENUE MAX Herbicide per year for this use.
	Sucker Management		DO NOT exceed 2 applications of VENUE MAX Herbicide per year for this use.
In-Season	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed a combined total of 3 applications of VENUE MAX Herbicide per year for these uses.
	Sucker Management		

(continued)

CROP USE DIRECTIONS *(continued)*

Bearing and Nonbearing

Pome Fruit Group (Crop Group 11-10); Pomegranate; Small Fruit Vine Climbing Subgroup Except Fuzzy Kiwifruit (Crop Group 13-07F); Stone Fruit Group (Crop Group 12-12); Tree Nut Group (Crop Group 14-12); Tropical and Subtropical, Small Fruit, Edible Peel Subgroup (Crop Subgroup 23A)

Application Instructions

- Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds and sucker growth.
- Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions.
- COC adjuvants are advised, although other adjuvants may be used.
- Use the higher rate for hard-to-control weeds.
- For the management of undesirable sucker growth on the basal portion of trunks, root sprouts, and vine trunks, growth must be controlled when the tissue is young, immature, and/or not hardened off.
- Avoid contact with green, uncalled bark of young trees or vines established less than one year unless protected from spray contact by nonporous wraps, grow tubes, or waxed containers.

USE RESTRICTIONS

- **DO NOT** apply by air for this use.
- **DO NOT** allow spray to drift onto desirable fruit, foliage, vines, or trees as damage will occur.
- **DO NOT** exceed 1.9 fl oz (0.0053 lb ai) of **VENUE MAX** Herbicide per acre in a single application for these uses.
- **DO NOT** exceed 3.3 fl oz (0.0090 lb ai) of **VENUE MAX** Herbicide per acre per year for all postharvest, dormant, and pre-bloom applications combined.
- **DO NOT** exceed 5.8 fl oz (0.0159 lb ai) of **VENUE MAX** Herbicide per acre per year for all in-season applications combined.
- Allow a minimum of 30 days between applications for this use.
- Preharvest Interval (PHI): 0 days

(continued)

CROP USE DIRECTIONS *(continued)*

Nonbearing Only

Citrus Fruit Group (Crop Group 10-10)

Australian desert lime; Australian finger lime; Australian round lime; Brown River finger lime; calamondin; citron; citrus hybrids; grapefruit; Japanese summer grapefruit; kumquat; lemon; lime; Mediterranean mandarin; Mount White lime; New Guinea wild lime; orange, sour; orange, sweet; pummelo; Russell River lime; satsuma mandarin; sweet lime; tachibana orange; Tahiti lime; tangelo; tangerine (mandarin, clementine); tangor; trifoliate orange; unqi fruit; cultivars, varieties, and/or hybrids of these

Application	Pest	Rate/Acre	Applications Per Year
Dormant In-Season	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	DO NOT exceed a combined total of 3 applications of VENUE MAX Herbicide per year for this use.

Application Instructions

- Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds.
- The addition of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions.
- COC adjuvants are advised, though other adjuvants may be used.
- Avoid spray contact with any foliage or low hanging branches (tree skirts) and with green, uncalledus bark of young trees.
- Use the higher rate for hard-to-control weeds.

USE RESTRICTIONS

- **DO NOT** apply by air for this use.
- **DO NOT** exceed 1.9 fl oz/acre (0.0053 lb ai/acre) of **VENUE MAX** Herbicide in a single application for these uses.
- **DO NOT** exceed 3.3 fl oz/acre (0.0090 lb ai/acre) of **VENUE MAX** Herbicide per year for dormant applications.
- **DO NOT** exceed 5.8 fl oz/acre (0.0159 lb ai/acre) of **VENUE MAX** Herbicide per year for all in-season applications combined.
- Allow a minimum of 30 days between applications for this use.

(continued)

CROP USE DIRECTIONS *(continued)*

Hops (Not for Use in California)		
Application	Pest	Rate/Acre
In-Season	Listed Broadleaf Weeds Sucker Management	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)
Application Instructions		
<ul style="list-style-type: none"> • Apply in a minimum of 20 gallons spray solution per acre by ground equipment to target weeds and sucker growth. • Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. • COC adjuvants are advised, although other adjuvants may be used. • Use the higher rate for hard-to-control weeds. • For the management of undesirable sucker growth, growth must be controlled when the tissue is young, immature, and/or not hardened off. • Direct application to the lower 12-18 inches of the hops plant and extend out to approximately 20-40 inches from each side of the row. • Calculate the rate on a treated acre basis as follows: Assuming a plot length of 50 feet: Calculations would be 1.5 feet (if 18-inch extension), 2 feet (if 24-inch extension) x 50 feet x 2 passes (one on each side of the vine). <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • DO NOT apply by air for this use. • DO NOT allow spray to drift onto desirable foliage or cones as damage will occur. • DO NOT exceed 1.9 fl oz (0.0053 lb ai) of VENUE MAX Herbicide per acre in a single application for this use. • DO NOT exceed a combined total of 2 applications per year for these uses. • DO NOT exceed 3.8 fl oz (0.0106 lb ai) of VENUE MAX Herbicide per acre per year for all applications combined. • Allow a minimum of 30 days between applications for this use. • Preharvest Interval (PHI): 30 days 		

(continued)

CROP USE DIRECTIONS *(continued)*

Pasture and Rangeland		
Pest	Rate/Acre	Application Instructions
Listed Broadleaf Weeds	1.0 to 1.7 fl oz/acre (0.0027 to 0.0046 lb ai/acre)	<ul style="list-style-type: none"> • Apply in a minimum of 2 gallons water per acre by air or 10 gallons water per acre by ground for this application. • Use of a crop oil or spray tank adjuvant at a concentration of 0.5% to 1.0% is directed for optimum weed control. • Livestock may graze treated areas as soon as the spray solution has dried on the foliage. • Refer to Rotational Crop Restrictions table. • Use the higher rate for hard-to-control weeds.
Application Instructions		
<p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • DO NOT exceed 1.7 fl oz (0.0046 lb ai) of VENUE MAX Herbicide per acre in a single application for this use. • DO NOT make more than 2 applications or exceed 3.4 fl oz (0.0093 lb ai) of VENUE MAX Herbicide per acre per year for this use. • Allow a minimum of 14 days between applications for this use. • Preharvest Interval (PHI): 0 days 		

(continued)

CROP USE DIRECTIONS *(continued)*

Fallow Bed and Crop Stubble		
Application	Pest	Rate/Acre
Preplant Burndown	Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)
Application Instructions		
<ul style="list-style-type: none"> • Apply in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground. • Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. • COC adjuvants are advised, although other adjuvants may be used. • Refer to Rotational Crop Restrictions table. • Use the higher rate for hard-to-control weeds. <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • DO NOT exceed 1.9 fl oz (0.0053 lb ai) of VENUE MAX Herbicide per acre in a single application for this use. • DO NOT make more than 3 applications or exceed 3.3 fl oz (0.0090 lb ai) of VENUE MAX Herbicide per acre per year. • Allow a minimum of 30 days between applications for this use. 		

(continued)

CROP USE DIRECTIONS *(continued)*

Noncropland; Uncultivated Agricultural Areas; Conservation Reserve Program Land/Federal Set-Aside Acreage* (Non-Food Producing)

Pest	Rate/Acre	Application Instructions
Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	<ul style="list-style-type: none"> • Apply in a minimum of 5 gallons spray solution per acre by air or 10 gallons spray solution per acre by ground. • Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. • COC adjuvants are advised, although other adjuvants may be used.

Application Instructions

- Refer to Rotational Crop Restrictions table.
- Use the higher rate for hard-to-control weeds.
- Applications to noncropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

USE RESTRICTIONS

- **DO NOT** exceed 1.9 fl oz (0.0053 lb ai) of **VENUE MAX** Herbicide per acre in a single application for this use.
- **DO NOT** make more than 3 applications or exceed 3.3 fl oz (0.0090 lb ai) of **VENUE MAX** Herbicide per acre per year.
- Allow a minimum of 30 days between applications for this use.

*Follow federal, state, and local rules for use on grass and hay.

(continued)

CROP USE DIRECTIONS *(continued)*

Noncrop Weed Control

airports and airfields; commercial plants; storage and lumber yards; fence lines and fence rows; farmyards and farm buildings; barrier strips and firebreaks; equipment areas; railroads; roadside and utility rights-of-way; fuel tank farms and pumping stations; dry ditches and ditchbanks; vacant lots; and similar agricultural and industrial noncrop sites

Pest	Rate/Acre	Application Instructions
Listed Broadleaf Weeds	1.5 to 1.9 fl oz/acre (0.0040 to 0.0053 lb ai/acre)	<ul style="list-style-type: none"> • Apply in a minimum of 20 to 40 gallons spray solution per acre by ground. • Avoid contact with desirable vegetation. • Use of a COC adjuvant at a concentration of 1% to 2% is directed for optimum weed control. Use the higher COC rate for larger labeled weed species or in low moisture conditions. • COC adjuvants are advised, although other adjuvants may be used. • Use the higher rate for hard-to-control weeds. <p>USE RESTRICTIONS</p> <ul style="list-style-type: none"> • DO NOT exceed 1.9 fl oz (0.0053 lb ai) of VENUE MAX Herbicide per acre in a single application for this use. • DO NOT make more than 3 applications or exceed 5.8 fl oz (0.0159 lb ai) of VENUE MAX Herbicide per acre per year. • Allow a minimum of 30 days between applications for this use.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container, and keep tightly closed when not in use. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

IMPORTANT: READ BEFORE USE

By using this product, user or buyer accepts the following conditions, warranty, disclaimer of warranties, and limitations of liability.

CONDITIONS: The directions for use of this product are believed to be accurate and must be followed carefully. However, because of extreme weather and soil conditions, use methods and other factors beyond the control of Nichino America, Inc. (NAI), it is impossible for NAI to eliminate all risks associated with the use of this product. As a result, crop injury or ineffectiveness is always possible. To the extent consistent with applicable law, all such risks are assumed by the user or buyer.

DISCLAIMER OF WARRANTIES: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THERE ARE NO WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, WHICH EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of NAI is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, NAI disclaims any liability whatsoever for incidental or consequential damages, including, but not limited to, liability arising out of breach of contract, express or implied warranty (including warranties of merchantability and fitness for a particular purpose), tort, negligence, strict liability or otherwise.

LIMITATIONS OF LIABILITY: TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES, OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT THE ELECTION OF NICHINO AMERICA, THE REPLACEMENT OF PRODUCT.

©2022 Nichino America, Inc. Venue is a registered trademark of Nichino America, Inc. The Fujjichi-mark is a trademark of Nihon Nohyaku Co., Ltd. Viton is a trademark of The Chemours Company.

Formulated and Packaged for
Nichino America, Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700

518-3 06302021
06/09/2022



VENUE[®] MAX
HERBICIDE

ACTIVE INGREDIENT:

Pyraflufen-ethyl: Acetic acid, [2-chloro-5-[4-chloro-5-(difluoromethoxy)-1-methyl-1H-pyrazol-3-yl]-4-fluorophenoxy]-, ethyl ester 4.0%

OTHER INGREDIENTS: 96.0%

TOTAL 100.0%

Contains 0.35 lb pyraflufen-ethyl per gallon

EPA Reg. No. 71711- 69

CB

E

AYE

EPA Est. No. 70815-GA-002 39578-TX-1 54675-MEX-001

superscript corresponds to lot number

KEEP OUT OF REACH OF CHILDREN CAUTION

FIRST AID: **If on skin or clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For additional information on this pesticide product, including human health concerns and medical emergencies, call 1-800-348-5832. In case of fire or spills, information may be obtained by calling 1-800-424-9300.

PYRAFLUFEN-ETHYL

GROUP

14

HERBICIDE

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container, and keep tightly closed when not in use. Store in a cool, dry place.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

CONTAINER HANDLING: Nonrefillable container. **DO NOT** reuse or refill this container. Clean container promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available, or reconditioning if appropriate, or puncture and dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

See attached booklet for First Aid, Precautionary Statements, and Directions for Use

Formulated and Packaged for
Nichino America Inc.
4550 Linden Hill Road, Suite 501
Wilmington, DE 19808
888-740-7700

NET CONTENTS: 1 quart

920102 02/24